


Balaji

Advanced Problems in Mathematics Chapter 10 to 26

for IIT JEE Main and Advanced

by

Vikas Gupta and Pankaj Joshi



**श्री**  
**Balaji**

**Advanced Problems *in***  
**MATHEMATICS**

*for*  
**JEE (MAIN & ADVANCED)**

*by :*

**Vikas Gupta**  
Director  
Vibrant Academy India(P) Ltd.  
KOTA (Rajasthan)

**Pankaj Joshi**  
Director  
Vibrant Academy India(P) Ltd.  
KOTA (Rajasthan)

# CONTENTS

## CALCULUS

1. Function	3 – 29
2. Limit	30 – 44
3. Continuity, Differentiability and Differentiation	45 – 74
4. Application of Derivatives	75 – 97
5. Indefinite and Definite Integration	98 – 127
6. Area Under Curves	128 – 134
7. Differential Equations	135 – 144

## ALGEBRA

8. Quadratic Equations	147 – 176
9. Sequence and Series	177 – 197
10. Determinants	198 – 206
11. Complex Numbers	207 – 216
12. Matrices	217 – 224
13. Permutation and Combinations	225 – 233
14. Binomial Theorem	234 – 242
15. Probability	243 – 251
16. Logarithms	252 – 264



## **CO-ORDINATE GEOMETRY**

17. Straight Lines	267 – 280
18. Circle	281 – 295
19. Parabola	296 – 302
20. Ellipse	303 – 307
21. Hyperbola	308 – 312

## **TRIGONOMETRY**

22. Compound Angles	315 – 334
23. Trigonometric Equations	335 – 343
24. Solution of Triangles	344 – 359
25. Inverse Trigonometric Functions	360 – 370

## **VECTOR & 3DIMENSIONAL GEOMETRY**

26. Vector & 3Dimensional Geometry	373 – 389
------------------------------------	-----------

 **Exercise-1 : Single Choice Problems**

1. If  $\begin{vmatrix} 1 & \cos \alpha & \cos \beta \\ \cos \alpha & 1 & \cos \gamma \\ \cos \beta & \cos \gamma & 1 \end{vmatrix} = \begin{vmatrix} 0 & \cos \alpha & \cos \beta \\ \cos \alpha & 0 & \cos \gamma \\ \cos \beta & \cos \gamma & 0 \end{vmatrix}$  then the value of  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma$  is :
- (a) 1                      (b)  $\frac{3}{2}$                       (c)  $\frac{3}{8}$                       (d)  $\frac{9}{4}$
2. Let the following system of equations
- $$\begin{aligned} kx + y + z &= 1 \\ x + ky + z &= k \\ x + y + kz &= k^2 \end{aligned}$$
- has no solution. Find  $|k|$ .
- (a) 0                      (b) 1                      (c) 2                      (d) 3
3. If  $\begin{vmatrix} a & a^2 & 1+a^3 \\ b & b^2 & 1+b^3 \\ c & c^2 & 1+c^3 \end{vmatrix} = 0$  and vectors  $(1, a, a^2)$ ,  $(1, b, b^2)$  and  $(1, c, c^2)$  are non-coplanar, then the product  $abc$  equals :
- (a) 2                      (b) -1                      (c) 1                      (d) 0
4. If the system of linear equations
- $$\begin{aligned} x + 2ay + az &= 0 \\ x + 3by + bz &= 0 \\ x + 4cy + cz &= 0 \end{aligned}$$
- has a non-zero solution, then  $a, b, c$  :
- (a) are in A.P.                      (b) are in G.P.  
(c) are in H.P.                      (d) satisfy  $a + 2b + 3c = 0$
5. If the number of quadratic polynomials  $ax^2 + 2bx + c$  which satisfy the following conditions :
- (i)  $a, b, c$  are distinct

(ii)  $a, b, c \in \{1, 2, 3, \dots, 2001, 2002\}$

(iii)  $x + 1$  divides  $ax^2 + 2bx + c$

is equal to  $1000\lambda$ , then find the value of  $\lambda$ .

- (a) 2002                      (b) 2001                      (c) 2003                      (d) 2004

6. If the system of equations  $2x + ay + 6z = 8$ ,  $x + 2y + z = 5$ ,  $2x + ay + 3z = 4$  has a unique solution then 'a' cannot be equal to :

- (a) 2                              (b) 3                              (c) 4                              (d) 5

7. If one of the roots of the equation  $\begin{vmatrix} 7 & 6 & x^2 - 13 \\ 2 & x^2 - 13 & 2 \\ x^2 - 13 & 3 & 7 \end{vmatrix} = 0$  is  $x = 2$ , then sum of all other

five roots is :

- (a) -2                              (b) 0                              (c)  $2\sqrt{5}$                       (d)  $\sqrt{15}$

8. The system of equations

$$kx + (k + 1)y + (k - 1)z = 0$$

$$(k + 1)x + ky + (k + 2)z = 0$$

$$(k - 1)x + (k + 2)y + kz = 0$$

has a nontrivial solution for :

- (a) Exactly three real values of  $k$ .                      (b) Exactly two real values of  $k$ .  
 (c) Exactly one real value of  $k$ .                      (d) Infinite number of values of  $k$ .

9. If  $a_1, a_2, a_3, \dots, a_n$  are in G.P. and  $a_i > 0$  for each  $i$ , then the determinant

$$\Delta = \begin{vmatrix} \log a_n & \log a_{n+2} & \log a_{n+4} \\ \log a_{n+6} & \log a_{n+8} & \log a_{n+10} \\ \log a_{n+12} & \log a_{n+14} & \log a_{n+16} \end{vmatrix}$$
 is equal to :

- (a) 0                              (b)  $\log \left( \sum_{i=1}^{n^2+n} a_i \right)$                       (c) 1                              (d) 2

10. If  $D_1 = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$  and  $D_2 = \begin{vmatrix} a_1 + 2a_2 + 3a_3 & 2a_3 & 5a_2 \\ b_1 + 2b_2 + 3b_3 & 2b_3 & 5b_2 \\ c_1 + 2c_2 + 3c_3 & 2c_3 & 5c_2 \end{vmatrix}$  then  $\frac{D_2}{D_1}$  is equal to :

- (a) 10                              (b) -10                              (c) 20                              (d) -20

11. If  $\Delta_1 = \begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix}$  and  $\Delta_2 = \begin{vmatrix} 1 & bc & a \\ 1 & ac & b \\ 1 & ab & c \end{vmatrix}$  then :

- (a)  $\Delta_1 = \Delta_2$                       (b)  $\Delta_1 = 2\Delta_2$                       (c)  $\Delta_1 + \Delta_2 = 0$                       (d)  $\Delta_1 + 2\Delta_2 = 0$

12. The value of the determinant  $\begin{vmatrix} 1 & 0 & -1 \\ a & 1 & 1-a \\ b & a & 1+a-b \end{vmatrix}$  depends on :

- (a) only  $a$                       (b) only  $b$                       (c) neither  $a$  nor  $b$                       (d) both  $a$  and  $b$

13. Sum of solutions of the equation  $\begin{vmatrix} 1 & 2 & x \\ 2 & 3 & x^2 \\ 3 & 5 & 2 \end{vmatrix} = 10$  is :
- (a) 1 (b) -1 (c) 2 (d) 4
14. If  $D = \begin{vmatrix} x+d & x+e & x+f \\ x+d+1 & x+e+1 & x+f+1 \\ x+a & x+b & x+c \end{vmatrix}$  then  $D$  does not depend on :
- (a)  $a$  (b)  $e$  (c)  $d$  (d)  $x$
15. The value of the determinant  $\begin{vmatrix} x-y-z & 2x & 2x \\ 2y & y-z-x & 2y \\ 2z & 2z & z-x-y \end{vmatrix} =$
- (a)  $xyz(x+y+z)^2$  (b)  $(x+y-z)(x+y+z)^2$   
 (c)  $(x+y+z)^3$  (d)  $(x+y+z)^2$
16. A rectangle  $ABCD$  is inscribed in a circle. Let  $PQ$  be the diameter of the circle parallel to the side  $AB$ . If  $\angle BPC = 30^\circ$ , then the ratio of the area of rectangle to the area of circle is :
- (a)  $\frac{\sqrt{3}}{\pi}$  (b)  $\frac{\sqrt{3}}{2\pi}$  (c)  $\frac{3}{\pi}$  (d)  $\frac{\sqrt{3}}{9\pi}$
17. Let  $ab = 1$ ,  $\Delta = \begin{vmatrix} 1+a^2-b^2 & 2ab & -2b \\ 2ab & 1-a^2+b^2 & 2a \\ 2b & -2a & 1-a^2-b^2 \end{vmatrix}$  then the minimum value of  $\Delta$  is :
- (a) 3 (b) 9 (c) 27 (d) 81
18. The determinant  $\begin{vmatrix} 2 & a+b+c+d & ab+cd \\ a+b+c+d & 2(a+b)(c+d) & ab(c+d)+cd(a+b) \\ ab+cd & ab(c+d)+cd(a+b) & 2abcd \end{vmatrix} = 0$  for
- (a)  $a+b+c+d = 0$  (b)  $ab+cd = 0$   
 (c)  $ab(c+d)+cd(a+b) = 0$  (d) any  $a, b, c, d$
19. Let  $\det A = \begin{vmatrix} l & m & n \\ p & q & r \\ 1 & 1 & 1 \end{vmatrix}$  and if  $(l-m)^2 + (p-q)^2 = 9$ ,  $(m-n)^2 + (q-r)^2 = 16$ ,  $(n-l)^2 + (r-p)^2 = 25$ , then the value of  $(\det A)^2$  equals :
- (a) 36 (b) 100 (c) 144 (d) 169
20. The number of distinct real values of  $K$  such that the system of equations  $x+2y+z=1$ ,  $x+3y+4z=K$ ,  $x+5y+10z=K^2$  has infinitely many solutions is :
- (a) 0 (b) 4 (c) 2 (d) 3

21. If  $\begin{vmatrix} (x+1) & (x+1)^2 & (x+1)^3 \\ (x+2) & (x+2)^2 & (x+2)^3 \\ (x+3) & (x+3)^2 & (x+3)^3 \end{vmatrix}$  is expressed as a polynomial in  $x$ , then the term independent of

$x$  is :

- (a) 0 (b) 2 (c) 12 (d) 16

22. If  $A, B, C$  are the angles of triangle  $ABC$ , then the minimum value of  $\begin{vmatrix} -2 & \cos C & \cos B \\ \cos C & -1 & \cos A \\ \cos B & \cos A & -1 \end{vmatrix}$  is

equal to :

- (a) 0 (b) -1 (c) 1 (d) -2

23. If the system of linear equations

$$x + 2ay + az = 0$$

$$x + 3by + bz = 0$$

$$x + 4cy + cz = 0$$

has a non-zero solution then  $a, b, c$  are in

- (a) A.P (b) G.P (c) H.P (d) None of these

24. If  $a, b$  and  $c$  are the roots of the equation  $x^3 + 2x^2 + 1 = 0$ , find  $\begin{vmatrix} a & b & x \\ b & c & a \\ c & a & b \end{vmatrix}$ .

- (a) 8 (b) -8 (c) 0 (d) 2

25. The system of homogeneous equation  $\lambda x + (\lambda + 1)y + (\lambda - 1)z = 0$ ,

$(\lambda + 1)x + \lambda y + (\lambda + 2)z = 0, (\lambda - 1)x + (\lambda + 2)y + \lambda z = 0$  has non-trivial solution for :

- (a) exactly three real values of  $\lambda$  (b) exactly two real values of  $\lambda$   
 (c) exactly three real value of  $\lambda$  (d) infinitely many real value of  $\lambda$

26. If one of the roots of the equation  $\begin{vmatrix} 7 & 6 & x^2 - 13 \\ 2 & x^2 - 13 & 2 \\ x^2 - 13 & 3 & 7 \end{vmatrix} = 0$  is  $x = 2$ , then sum of all other


five roots is :

- (a) -2 (b) 0 (c)  $2\sqrt{5}$  (d)  $\sqrt{15}$

### Answers

1.	(a)	2.	(c)	3.	(b)	4.	(c)	5.	(a)	6.	(c)	7.	(a)	8.	(c)	9.	(a)	10.	(b)
11.	(c)	12.	(c)	13.	(b)	14.	(d)	15.	(c)	16.	(a)	17.	(c)	18.	(d)	19.	(c)	20.	(c)
21.	(c)	22.	(b)	23.	(c)	24.	(a)	25.	(c)	26.	(a)								




**Exercise-2 : One or More than One Answer is/are Correct**

1. Let  $f(a, b) = \begin{vmatrix} a & a^2 & 0 \\ 1 & (2a+b) & (a+b)^2 \\ 0 & 1 & (2a+3b) \end{vmatrix}$ , then

(a)  $(2a+b)$  is a factor of  $f(a, b)$

(b)  $(a+2b)$  is a factor of  $f(a, b)$

(c)  $(a+b)$  is a factor of  $f(a, b)$

(d)  $a$  is a factor of  $f(a, b)$

2. If  $\begin{vmatrix} 1 + \cos^2 \theta & \sin^2 \theta & 2\sqrt{3} \tan \theta \\ \cos^2 \theta & 1 + \sin^2 \theta & 2\sqrt{3} \tan \theta \\ \cos^2 \theta & \sin^2 \theta & 1 + 2\sqrt{3} \tan \theta \end{vmatrix} = 0$  then  $\theta$  may be :

(a)  $\frac{\pi}{6}$

(b)  $\frac{5\pi}{6}$

(c)  $\frac{7\pi}{6}$

(d)  $\frac{11\pi}{6}$

3. Let  $\Delta = \begin{vmatrix} a & a+d & a+3d \\ a+d & a+2d & a \\ a+2d & a & a+d \end{vmatrix}$  then :

(a)  $\Delta$  depends on  $a$

(b)  $\Delta$  depends on  $d$

(c)  $\Delta$  is independent of  $a, d$

(d)  $\Delta = 0$

4. The value(s) of  $\lambda$  for which the system of equations

$$(1-\lambda)x + 3y - 4z = 0$$

$$x - (3+\lambda)y + 5z = 0$$

$$3x + y - \lambda z = 0$$

possesses non-trivial solutions.

(a)  $-1$

(b)  $0$

(c)  $1$

(d)  $2$

5. Let  $D(x) = \begin{vmatrix} x^2 + 4x - 3 & 2x + 4 & 13 \\ 2x^2 + 5x - 9 & 4x + 5 & 26 \\ 8x^2 - 6x + 1 & 16x - 6 & 104 \end{vmatrix} = \alpha x^3 + \beta x^2 + \gamma x + \delta$  then :

(a)  $\alpha + \beta = 0$

(b)  $\beta + \gamma = 0$

(c)  $\alpha + \beta + \gamma + \delta = 0$

(d)  $\alpha + \beta + \gamma = 0$

6. Let  $D(x) = \begin{vmatrix} x^2 + 4x - 3 & 2x + 4 & 13 \\ 2x^2 + 5x - 9 & 4x + 5 & 26 \\ 8x^2 - 6x + 1 & 16x - 6 & 104 \end{vmatrix} = \alpha x^3 + \beta x^2 + \gamma x + \delta$  then :

(a)  $\alpha + \beta = 0$

(b)  $\beta + \gamma = 0$

(c)  $\alpha + \beta + \gamma + \delta = 0$

(d)  $\alpha + \beta + \gamma = 0$

7. If the system of equations

$$ax + y + 2z = 0$$

$$x + 2y + z = b$$

$$2x + y + az = 0$$

has no solution then  $(a+b)$  can be equals to :

(a)  $-1$

(b)  $2$

(c)  $3$

(d)  $4$





**Exercise-4 : Subjective Type Problems**

1. If  $3^n$  is a factor of the determinant  $\begin{vmatrix} 1 & 1 & 1 \\ {}^n C_1 & {}^{n+3} C_1 & {}^{n+6} C_1 \\ {}^n C_2 & {}^{n+3} C_2 & {}^{n+6} C_2 \end{vmatrix}$  then the maximum value of  $n$  is

.....

2. Find the value of  $\lambda$  for which  $\begin{vmatrix} 2a_1 + b_1 & 2a_2 + b_2 & 2a_3 + b_3 \\ 2b_1 + c_1 & 2b_2 + c_2 & 2b_3 + c_3 \\ 2c_1 + a_1 & 2c_2 + a_2 & 2c_3 + a_3 \end{vmatrix} = \lambda \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$

3. Find the co-efficient of  $x$  in the expansion of the determinant  $\begin{vmatrix} (1+x)^2 & (1+x)^4 & (1+x)^6 \\ (1+x)^3 & (1+x)^6 & (1+x)^9 \\ (1+x)^4 & (1+x)^8 & (1+x)^{12} \end{vmatrix}$ .

4. If  $x, y, z \in R$  and  $\begin{vmatrix} x & y^2 & z^3 \\ x^4 & y^5 & z^6 \\ x^7 & y^8 & z^9 \end{vmatrix} = 2$  then find the value of

$$\begin{vmatrix} y^5 z^6 (z^3 - y^3) & x^4 z^6 (x^3 - z^3) & x^4 y^5 (y^3 - x^3) \\ y^2 z^3 (y^6 - z^6) & xz^3 (z^6 - x^6) & xy^2 (x^6 - y^6) \\ y^2 z^3 (z^3 - y^3) & xz^3 (x^3 - z^3) & xy^2 (y^3 - x^3) \end{vmatrix}$$

5. If the system of equations :

$$\begin{aligned} 2x + 3y - z &= 0 \\ 3x + 2y + kz &= 0 \\ 4x + y + z &= 0 \end{aligned}$$

have a set of non-zero integral solutions then, find the smallest positive value of  $z$ .

6. Find  $a \in R$  for which the system of equations  $2ax - 2y + 3z = 0$ ;  $x + ay + 2z = 0$  and  $2x + az = 0$  also have a non-trivial solution.

7. If three non-zero distinct real numbers form an arithmetic progression and the squares of these numbers taken in the same order constitute a geometric progression. Find the sum of all possible common ratios of the geometric progression.

8. Let  $\Delta_1 = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$ ,  $\Delta_2 = \begin{vmatrix} 6a_1 & 2a_2 & 2a_3 \\ 3b_1 & b_2 & b_3 \\ 12c_1 & 4c_2 & 4c_3 \end{vmatrix}$  and  $\Delta_3 = \begin{vmatrix} 3a_1 + b_1 & 3a_2 + b_2 & 3a_3 + b_3 \\ 3b_1 & 3b_2 & 3b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{vmatrix}$

then  $\Delta_3 - \Delta_2 = k\Delta_1$ , find  $k$ .

9. The minimum value of determinant  $\Delta = \begin{vmatrix} 1 & \cos \theta & 1 \\ -\cos \theta & 1 & \cos \theta \\ -1 & -\cos \theta & 2 \end{vmatrix} \forall \theta \in R$  is :

10. For a unique value of  $\mu$  &  $\lambda$ , the system of equations given by

$$x + y + z = 6$$

$$x + 2y + 3z = 14$$

$$2x + 5y + \lambda z = \mu$$

has infinitely many solutions, then  $\frac{\mu - \lambda}{4}$  is equal to

11. Let  $\lim_{n \rightarrow \infty} n \sin(2\pi e \lfloor \frac{n}{e} \rfloor) = k\pi$ , where  $n \in \mathbb{N}$ . Find  $k$ :

12. If the system of linear equations

$$(\cos \theta)x + (\sin \theta)y + \cos \theta = 0$$

$$(\sin \theta)x + (\cos \theta)y + \sin \theta = 0$$

$$(\cos \theta)x + (\sin \theta)y - \cos \theta = 0$$

is consistent, then the number of possible values of  $\theta$ ,  $\theta \in [0, 2\pi]$  is :

### Answers

1.	3	2.	9	3.	0	4.	4	5.	5	6.	2	7.	6
8.	3	9.	3	10.	7	11.	2	12.	2				

□□□


### Exercise-1 : Single Choice Problems

- Let  $t_1, t_2, t_3$  be three distinct points on circle  $|t|=1$ . If  $\theta_1, \theta_2$  and  $\theta_3$  be the arguments of  $t_1, t_2, t_3$  respectively then  $\cos(\theta_1 - \theta_2) + \cos(\theta_2 - \theta_3) + \cos(\theta_3 - \theta_1)$ 
  - $\geq -\frac{3}{2}$
  - $\leq -\frac{3}{2}$
  - $\geq \frac{3}{2}$
  - $\leq 2$
- The number of points of intersection of the curves represented by  $\arg(z - 2 - 7i) = \cot^{-1}(2)$  and  $\arg\left(\frac{z - 5i}{z + 2 - i}\right) = \pm\frac{\pi}{2}$ 
  - 0
  - 1
  - 2
  - None of these
- All three roots of  $az^3 + bz^2 + cz + d = 0$ , have negative real part, ( $a, b, c \in R$ ) then :
  - All  $a, b, c, d$  have the same sign
  - $a, b, c$  have same sign
  - $a, b, d$  have same sign
  - $b, c, d$  have same sign
- Let  $z_1$  and  $z_2$  be two roots of the equation  $z^2 + az + b = 0$ ,  $z$  being complex number. Further, assume that the origin,  $z_1$  and  $z_2$  form an equilateral triangle, then :
  - $a^2 = b$
  - $a^2 = 2b$
  - $a^2 = 3b$
  - $a^2 = 4b$
- If  $z$  and  $\omega$  are two non-zero complex numbers such that  $|z\omega| = 1$ , and  $\arg(z) - \arg(\omega) = \frac{\pi}{2}$ , then  $\bar{z}\omega$  is equal to :
  - 1
  - 1
  - $i$
  - $-i$
- If  $\omega$  be an imaginary  $n^{\text{th}}$  root of unity, then  $\sum_{r=1}^n (ar + b)\omega^{r-1}$  is equal to :
  - $\frac{n(n+1)a}{2\omega}$
  - $\frac{nb}{1-n}$
  - $\frac{na}{\omega-1}$
  - None of these

7. If  $\alpha, \beta$  are complex numbers then the maximum value of  $\frac{\alpha\bar{\beta} + \bar{\alpha}\beta}{|\alpha\beta|}$  is equal to :
- (a) 1 (b) 2 (c) greater than 2 (d) less than 1
8. Let  $z_1, z_2, z_3$  and  $z_4$  be the roots of the equation  $z^4 + z^3 + 2 = 0$ , then the value of  $\prod_{r=1}^4 (2z_r + 1)$  is equal to :
- (a) 28 (b) 29 (c) 30 (d) 31
9. If  $\arg\left(\frac{z-6-3i}{z-3-6i}\right) = \frac{\pi}{4}$ , then :
- (a) minimum value of  $|z|$  is  $6\sqrt{2} - 3$  (b) Maximum value of  $|z|$  is  $6\sqrt{2} + 3$   
(c) minimum value of  $|z|$  is  $15\sqrt{2} - 6$  (d) Maximum value of  $|z|$  is  $15\sqrt{2} + 6$
10. If  $z_1 \neq -z_2$  and  $|z_1 + z_2| = \left|\frac{1}{z_1} + \frac{1}{z_2}\right|$  then :
- (a) at least one of  $z_1, z_2$  is unimodular (b) both  $z_1, z_2$  are unimodular  
(c)  $z_1 \cdot z_2$  is unimodular (d)  $z_1 - z_2$  is unimodular
11. If  $|z - i| \leq 2$  and  $z_1 = 5 + 3i$ , then the maximum value of  $|iz + z_1|$  is :
- (a)  $5 + \sqrt{13}$  (b)  $5 + \sqrt{2}$  (c) 7 (d) 8
12. If  $z_1, z_2, z_3$  are vertices of a triangle such that  $|z_1 - z_2| = |z_1 - z_3|$  then  $\arg\left(\frac{2z_1 - z_2 - z_3}{z_3 - z_2}\right)$  is :
- (a)  $\pm\frac{\pi}{3}$  (b) 0 (c)  $\pm\frac{\pi}{2}$  (d)  $\pm\frac{\pi}{6}$
13. It is given that complex numbers  $z_1$  and  $z_2$  satisfy  $|z_1| = 2$  and  $|z_2| = 3$ . If the included angle of their corresponding vectors is  $60^\circ$ , then  $\left|\frac{z_1 + z_2}{z_1 - z_2}\right|$  can be expressed as  $\frac{\sqrt{n}}{7}$ , where 'n' is a natural number then  $n =$
- (a) 126 (b) 119 (c) 133 (d) 19
14. If all the roots of  $z^3 + az^2 + bz + c = 0$  are of unit modulus, then :
- (a)  $|a| \leq 3$  (b)  $|b| \leq 3$  (c)  $|c| = 1$  (d) All of the above
15. Let  $z$  be a complex number satisfying  $\frac{1}{2} \leq |z| \leq 4$ , then sum of greatest and least values of  $\left|z + \frac{1}{z}\right|$  is :
- (a)  $\frac{65}{4}$  (b)  $\frac{65}{16}$  (c)  $\frac{17}{4}$  (d) 17
16. If  $|z - 2i| \leq \sqrt{2}$ , then the maximum value of  $|3 + i(z - 1)|$  is :
- (a)  $\sqrt{2}$  (b)  $2\sqrt{2}$  (c)  $2 + \sqrt{2}$  (d)  $3 + 2\sqrt{2}$





 **Exercise-2 : One or More than One Answer is/are Correct**

1. Let  $Z_1$  and  $Z_2$  are two non-zero complex number such that  $|Z_1 + Z_2| = |Z_1| = |Z_2|$ , then  $\frac{Z_1}{Z_2}$  may be :
- (a)  $1 + \omega$  (b)  $1 + \omega^2$   
 (c)  $\omega$  (d)  $\omega^2$
2. Let  $z_1, z_2$  and  $z_3$  be three distinct complex numbers, satisfying  $|z_1| = |z_2| = |z_3| = 1$ . Which of the following is/are true :
- (a) If  $\arg\left(\frac{z_1}{z_2}\right) = \frac{\pi}{2}$  then  $\arg\left(\frac{z - z_1}{z - z_2}\right) > \frac{\pi}{4}$  where  $|z| > 1$   
 (b)  $|z_1 z_2 + z_2 z_3 + z_3 z_1| = |z_1 + z_2 + z_3|$   
 (c)  $\operatorname{Im}\left(\frac{(z_1 + z_2)(z_2 + z_3)(z_3 + z_1)}{z_1 \cdot z_2 \cdot z_3}\right) = 0$   
 (d) If  $|z_1 - z_2| = \sqrt{2}|z_1 - z_3| = \sqrt{2}|z_2 - z_3|$ , then  $\operatorname{Re}\left(\frac{z_3 - z_1}{z_3 - z_2}\right) = 0$
3. The triangle formed by the complex numbers  $z, iz, i^2 z$  is :
- (a) equilateral (b) isosceles  
 (c) right angled (d) isosceles but not right angled
4. If  $A(z_1), B(z_2), C(z_3), D(z_4)$  lies on  $|z| = 4$  (taken in order), where  $z_1 + z_2 + z_3 + z_4 = 0$  then:
- (a) Max. area of quadrilateral  $ABCD = 32$   
 (b) Max. area of quadrilateral  $ABCD = 16$   
 (c) The triangle  $\Delta ABC$  is right angled  
 (d) The quadrilateral  $ABCD$  is rectangle
5. Let  $z_1, z_2$  and  $z_3$  be three distinct complex numbers satisfying  $|z_1| = |z_2| = |z_3| = 1$ . Which of the following is/are true ?
- (a) If  $\arg\left(\frac{z_1}{z_2}\right) = \frac{\pi}{2}$  then  $\arg\left(\frac{z - z_1}{z - z_2}\right) > \frac{\pi}{4}$  where  $|z| > 1$   
 (b)  $|z_1 z_2 + z_2 z_3 + z_3 z_1| = |z_1 + z_2 + z_3|$   
 (c)  $\operatorname{Im}\left(\frac{(z_1 + z_2)(z_2 + z_3)(z_3 + z_1)}{z_1 \cdot z_2 \cdot z_3}\right) = 0$   
 (d) If  $|z_1 - z_2| = \sqrt{2}|z_1 - z_3| = \sqrt{2}|z_2 - z_3|$ , then  $\operatorname{Re}\left(\frac{z_3 - z_1}{z_3 - z_2}\right) = 0$

6. If  $z_1 = a + ib$  and  $z_2 = c + id$  are two complex numbers where  $a, b, c, d \in \mathbb{R}$  and  $|z_1| = |z_2| = 1$  and  $\text{Im}(z_1 \bar{z}_2) = 0$ . If  $w_1 = a + ic$  and  $w_2 = b + id$ , then :
- (a)  $\text{Im}(w_1 \bar{w}_2) = 0$  (b)  $\text{Im}(\bar{w}_1 w_2) = 0$   
 (c)  $\text{Im}\left(\frac{w_1}{w_2}\right) = 0$  (d)  $\text{Re}\left(\frac{w_1}{w_2}\right) = 0$
7. The solutions of the equation  $z^4 + 4iz^3 - 6z^2 - 4iz - i = 0$  represent vertices of a convex polygon in the complex plane. The area of the polygon is :  
 (a)  $2^{1/2}$  (b)  $2^{3/2}$  (c)  $2^{5/2}$  (d)  $2^{5/4}$
8. Least positive argument of the 4<sup>th</sup> root of the complex number  $2 - i\sqrt{12}$  is :  
 (a)  $\frac{\pi}{6}$  (b)  $\frac{\pi}{12}$  (c)  $\frac{5\pi}{12}$  (d)  $\frac{7\pi}{12}$
9. Let  $\omega$  be the imaginary cube root of unity and  $(a + b\omega + c\omega^2)^{2015} = (a + b\omega^2 + c\omega)$  where  $a, b, c$  are unequal real numbers. Then the value of  $a^2 + b^2 + c^2 - ab - bc - ca$  equals :  
 (a) 0 (b) 1 (c) 2 (d) 3
10. Let  $n$  be a positive integer and a complex number with unit modulus is a solution of the equation  $z^n + z + 1 = 0$  then the value of  $n$  can be :  
 (a) 62 (b) 155 (c) 221 (d) 196

### Answers

1.	(c, d)	2.	(b, c, d)	3.	(b, c)	4.	(a, c, d)	5.	(b, c, d)	6.	(a, b, c)
7.	(d)	8.	(c)	9.	(b)	10.	(a, b, c)				

### Exercise-3 : Comprehension Type Problems

#### Paragraph for Question Nos. 1 to 2

Let  $f(z)$  is of the form  $\alpha z + \beta$ , where  $\alpha, \beta$  are constants and  $\alpha, \beta, z$  are complex numbers such that  $|\alpha| \neq |\beta|$ .  $f(z)$  satisfies following properties :

- (i) If imaginary part of  $z$  is non zero, then  $f(z) + \overline{f(z)} = f(\bar{z}) + \overline{f(\bar{z})}$   
 (ii) If real part of  $z$  is zero, then  $f(z) + \overline{f(z)} = 0$   
 (iii) If  $z$  is real, then  $\overline{f(z)} f(z) > (z+1)^2 \forall z \in R$ .

1.  $\frac{4x^2}{(f(1) - f(-1))^2} + \frac{y^2}{(f(0))^2} = 1, x, y \in R$ , in  $(x, y)$  plane will represent :  
 (a) hyperbola (b) circle (c) ellipse (d) pair of line
2. Consider ellipse  $S : \frac{x^2}{(\operatorname{Re}(\alpha))^2} + \frac{y^2}{(\operatorname{Im}(\beta))^2} = 1, x, y \in R$  in  $(x, y)$  plane, then point  $(1, 1)$  will lie :  
 (a) outside the ellipse  $S$  (b) inside the ellipse  $S$   
 (c) on the ellipse  $S$  (d) none of these

#### Paragraph for Question Nos. 3 to 5

Let  $z_1$  and  $z_2$  be complex numbers, such what  $z_1^2 - 4z_2 = 16 + 20i$ . Also suppose that roots  $\alpha$  and  $\beta$  of  $t^2 + z_1 t + z_2 + m = 0$  for some complex number  $m$  satisfy  $|\alpha - \beta| = 2\sqrt{7}$ , then :

3. The complex number ' $m$ ' lies on :  
 (a) a square with side 7 and centre  $(4, 5)$  (b) a circle with radius 7 and centre  $(4, 5)$   
 (c) a circle with radius 7 and centre  $(-4, 5)$  (d) a square with side 7 and centre  $(-4, 5)$
4. The greatest value of  $|m|$  is :  
 (a)  $5\sqrt{21}$  (b)  $5 + \sqrt{23}$  (c)  $7 + \sqrt{43}$  (d)  $7 + \sqrt{41}$
5. The least value of  $|m|$  is :  
 (a)  $7 - \sqrt{41}$  (b)  $7 - \sqrt{43}$  (c)  $5 - \sqrt{23}$  (d)  $5 + \sqrt{21}$

#### Paragraph for Question Nos. 6 to 7

Let  $z_1 = 3$  and  $z_2 = 7$  represent two points  $A$  and  $B$  respectively on complex plane. Let the curve  $C_1$  be the locus of point  $P(z)$  satisfying  $|z - z_1|^2 + |z - z_2|^2 = 10$  and the curve  $C_2$  be the locus of point  $P(z)$  satisfying  $|z - z_1|^2 + |z - z_2|^2 = 16$ .

6. Least distance between curves  $C_1$  and  $C_2$  is :  
 (a) 4 (b) 3 (c) 2 (d) 1

7. The locus of point from which tangents drawn to  $C_1$  and  $C_2$  are perpendicular, is :  
 (a)  $|z - 5| = 4$       (b)  $|z - 3| = 2$       (c)  $|z - 5| = 3$       (d)  $|z - 5| = \sqrt{5}$

**Paragraph for Question Nos. 8 to 9**

In the Argand plane  $Z_1, Z_2$  and  $Z_3$  are respectively the vertices of an isosceles triangle  $ABC$  with  $AC = BC$  and  $\angle CAB = \theta$ . If  $I(Z_4)$  is the incentre of triangle, then :

8. The value of  $\left(\frac{AB}{IA}\right)^2 \left(\frac{AC}{AB}\right)$  is equal to :

(a)  $\left| \frac{(Z_2 - Z_1)(Z_1 - Z_3)}{(Z_4 - Z_1)} \right|$

(b)  $\left| \frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)} \right|$

(c)  $\left| \frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)^2} \right|$

(d)  $\left| \frac{(Z_2 + Z_1)(Z_3 + Z_1)}{(Z_4 + Z_1)} \right|$

9. The value of  $(Z_4 - Z_1)^2(1 + \cos\theta) \sec\theta$  is :

(a)  $(Z_2 - Z_1)(Z_3 - Z_1)$


(b)  $\frac{(Z_2 - Z_1)(Z_3 - Z_1)}{Z_4 - Z_1}$

(c)  $\frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)^2}$

(d)  $(Z_2 - Z_1)(Z_3 - Z_1)^2$

**Answers**

1.	(a)	2.	(b)	3.	(b)	4.	(d)	5.	(a)	6.	(d)	7.	(d)	8.	(c)	9.	(a)
----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----


**Exercise-4 : Matching Type Problems**

1. In a  $\Delta ABC$ , the side lengths  $BC$ ,  $CA$  and  $AB$  are consecutive positive integers in increasing order.

Column-I		Column-II
(A)	If $z_1, z_2$ and $z_3$ be the affixes of vertices $A, B$ and $C$ respectively in argand plane, such that $\left  \arg \left( \frac{z_1 - z_3}{z_2 - z_3} \right) \right  = \left  2 \arg \left( \frac{z_3 - z_1}{z_2 - z_1} \right) \right $ , then biggest side of the triangle is	(P) 2
(B)	Let $\vec{a}, \vec{b}$ and $\vec{c}$ be the position vectors of vertices $A, B$ and $C$ respectively. If $(\vec{c} - \vec{a}) \cdot (\vec{b} - \vec{c}) = 0$ then the value of $ \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} $ equals to	(Q) 3
(C)	Let the equations $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$ represent the lines $AB$ and $AC$ respectively and $\frac{a_1b_2 - a_2b_1}{a_1a_2 + b_1b_2} = \frac{4}{3}$ then the value of $s - c$ (where $s$ is the semiperimeter) $a = BC, b = CA, c = AB$	(R) 4
(D)	If the altitudes of $\Delta ABC$ are in harmonic progression then the side length ' $b$ ' can be	(S) 6
		(T) 12

2. Let  $ABCDEF$  is a regular hexagon  $A(z_1), B(z_2), C(z_3), D(z_4), E(z_5), F(z_6)$  in argand plane where  $A, B, C, D, E$  and  $F$  are taken in anticlockwise manner. If  $z_1 = -2, z_3 = 1 - \sqrt{3}i$ .

Column-I		Column-II
(A)	If $z_2 = a + ib$ , then $2a^2 + b^2$ is equal to	(P) 8
(B)	The square of the inradius of hexagon is	(Q) 7
(C)	The area of region formed by point $P(z)$ lying inside the incircle of hexagon and satisfying $\frac{\pi}{3} \leq \arg(z) \leq \frac{5\pi}{6}$ is $\frac{m}{n} \pi$ , where $m, n$ are relatively prime natural numbers, then $m + n$ is equal to	(R) 5
(D)	The value of $z_4^2 - z_1^2 - z_2^2 - z_3^2 - z_5^2 - z_6^2$ is equal to	(S) 3
		(T) 2

3.

Column-I		Column-II	
(A)	Let $\omega$ be a non real cube root of unity then the number of distinct elements in the set $\{(1 + \omega + \omega^2 + \dots + \omega^n)^m; n, m \in N\}$ is :	(P)	3
(B)	Let $\omega$ and $\omega^2$ be non real cube root of unity. The least possible degree of a polynomial with real co-efficients having roots $2\omega, (2 + 3\omega), (2 + 3\omega)^2, (2 - \omega - \omega^2)$ is	(Q)	4
(C)	Let $\alpha = 6 + 4i$ and $\beta = 2 + 4i$ are two complex numbers on Argand plane. A complex number $z$ satisfying $\text{amp} \left( \frac{z - \alpha}{z - \beta} \right) = \frac{\pi}{6}$ moves on a major segment of a circle whose radius is	(R)	5
(D)	Let $z_1, z_2, z_3$ are complex numbers denoting the vertices of an equilateral triangle $ABC$ having circumradius equals to unity. If $P$ denotes any arbitrary point on its circumcircle then the value of $\frac{1}{2}((PA)^2 + (PB)^2 + (PC)^2)$ equals to	(S)	7

### Answers

1.  $A \rightarrow S; B \rightarrow T; C \rightarrow S; D \rightarrow Q, R, S, T$
2.  $A \rightarrow R; B \rightarrow S; C \rightarrow Q; D \rightarrow P$
3.  $A \rightarrow S; B \rightarrow R; C \rightarrow Q; D \rightarrow P$

### Exercise-5 : Subjective Type Problems

- Let complex number 'z' satisfy the inequality  $2 \leq |z| \leq 4$ . A point P is selected in this region at random. The probability that argument of P lies in the interval  $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$  is  $\frac{1}{K}$ , then  $K =$
- Let  $z$  be a complex number satisfying  $|z - 3| \leq |z - 1|$ ,  $|z - 3| \leq |z - 5|$ ,  $|z - i| \leq |z + i|$  and  $|z - i| \leq |z - 5i|$ . Then the area of region in which  $z$  lies is A square units, where  $A =$
- Complex number  $z_1$  and  $z_2$  satisfy  $z + \bar{z} = 2|z - 1|$  and  $\arg(z_1 - z_2) = \frac{\pi}{4}$ . Then the value of  $\operatorname{Im}(z_1 + z_2)$  is :
- If  $|z_1| = 1$ ,  $|z_2| = 2$ ,  $|z_3| = 3$  and  $|9z_1z_2 + 4z_1z_3 + z_2z_3| = 36$ , then  $|z_1 + z_2 + z_3|$  is equal to :
- If  $|z_1|$  and  $|z_2|$  are the distances of points on the curve  $5z\bar{z} - 2i(z^2 - \bar{z}^2) - 9 = 0$  which are at maximum and minimum distance from the origin, then the value of  $|z_1| + |z_2|$  is equal to :
- Let  $\frac{1}{a_1 + \omega} + \frac{1}{a_2 + \omega} + \frac{1}{a_3 + \omega} + \dots + \frac{1}{a_n + \omega} = i$  where  $a_1, a_2, a_3, \dots, a_n \in R$  and  $\omega$  is imaginary cube root of unity, then evaluate  $\sum_{r=1}^n \frac{2a_r - 1}{a_r^2 - a_r + 1}$ .
- If  $|z_1| = 2$ ,  $|z_2| = 3$ ,  $|z_3| = 4$  and  $|2z_1 + 3z_2 + 4z_3| = 9$ , then value of  $|8z_2z_3 + 27z_3z_1 + 64z_1z_2|^{1/3}$  is :
- The sum of maximum and minimum modulus of a complex number  $z$  satisfying  $|z - 25i| \leq 15$ ,  $i = \sqrt{-1}$  is  $S$ , then  $\frac{S}{10}$  is :

### Answers

1.	4	2.	6	3.	2	4.	6	5.	4	6.	0	7.	6
8.	5												

□□□


**Exercise-1 : Single Choice Problems**

1. Let  $A = BB^T + CC^T$ , where  $B = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$ ,  $C = \begin{bmatrix} \sin \theta \\ -\cos \theta \end{bmatrix}$ ;  $\theta \in R$ . Then  $A$  is :
- (a)  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$       (b)  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$       (c)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$       (d)  $\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$
2. Let  $A = \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$ . The only correct statement about the matrix  $A$  is :
- (a)  $A$  is a zero matrix      (b)  $A^2 = I$ , where  $I$  is a unit matrix  
(c)  $A^{-1}$  does not exist      (d)  $A = (-1)I$ , where  $I$  is a unit matrix
3. Let  $A = [a_{ij}]_{3 \times 3}$  be such that  $a_{ij} = \begin{cases} 3; & \text{when } i = j \\ 0; & i \neq j \end{cases}$ , then  $\left\{ \frac{\det(\text{adj}(\text{adj } A))}{5} \right\}$  equals :
- (where  $\{ \cdot \}$  denotes fractional part function)
- (a)  $\frac{2}{5}$       (b)  $\frac{1}{5}$       (c)  $\frac{2}{3}$       (d)  $\frac{1}{3}$
4. If  $A^{-1} = \begin{bmatrix} \sin^2 \alpha & 0 & 0 \\ 0 & \sin^2 \beta & 0 \\ 0 & 0 & \sin^2 \gamma \end{bmatrix}$  and  $B^{-1} = \begin{bmatrix} \cos^2 \alpha & 0 & 0 \\ 0 & \cos^2 \beta & 0 \\ 0 & 0 & \cos^2 \gamma \end{bmatrix}$  where  $\alpha, \beta, \gamma$  are any real numbers and  $C = (A^{-5} + B^{-5}) + 5A^{-1}B^{-1}(A^{-3} + B^{-3}) + 10A^{-2}B^{-2}(A^{-1} + B^{-1})$  then find  $|C|$ .
- (a) 0      (b) 1      (c) 2      (d) 3
5. If  $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ ; then  $A^{-1} =$
- (a)  $A$       (b)  $A^2$       (c)  $A^3$       (d)  $A^4$
6. Let  $M = [a_{ij}]_{3 \times 3}$  where  $a_{ij} \in \{-1, 1\}$ . Find the maximum possible value of  $\det(M)$ .
- (a) 3      (b) 4      (c) 5      (d) 6



7. Let matrix  $A = \begin{bmatrix} x & 3 & 2 \\ 1 & y & 4 \\ 2 & 2 & z \end{bmatrix}$ ; if  $xyz = 2\lambda$  and  $8x + 4y + 3z = \lambda + 28$ , then  $(\text{adj } A)A$  equals :

(a)  $\begin{bmatrix} \lambda+1 & 0 & 0 \\ 0 & \lambda+1 & 0 \\ 0 & 0 & \lambda+1 \end{bmatrix}$

(b)  $\begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$

(c)  $\begin{bmatrix} \lambda^2 & 0 & 0 \\ 0 & \lambda^2 & 0 \\ 0 & 0 & \lambda^2 \end{bmatrix}$

(d)  $\begin{bmatrix} \lambda+2 & 0 & 0 \\ 0 & \lambda+2 & 0 \\ 0 & 0 & \lambda+2 \end{bmatrix}$

8. If the trace of matrix  $A = \begin{pmatrix} x-2 & e^x & -\sin x \\ \cos x^2 & x^2-x+3 & \ln|x| \\ 0 & \tan^{-1}x & x-7 \end{pmatrix}$  is zero, then  $x$  is equal to :

(a) -2 or 3

(b) -3 or -2

(c) -3 or 2

(d) 2 or 3

9. If  $A = [a_{ij}]_{2 \times 2}$  where  $a_{ij} = \begin{cases} i+j, & i \neq j \\ i^2-2j, & i = j \end{cases}$  then  $A^{-1}$  is equal to :

(a)  $\frac{1}{9} \begin{bmatrix} 0 & 3 \\ 3 & 1 \end{bmatrix}$

(b)  $\frac{1}{9} \begin{bmatrix} 0 & -3 \\ 3 & -1 \end{bmatrix}$

(c)  $\frac{1}{9} \begin{bmatrix} 0 & -3 \\ -3 & -1 \end{bmatrix}$

(d)  $\frac{1}{3} \begin{bmatrix} 0 & 3 \\ 3 & 1 \end{bmatrix}$

10. If  $\begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^{-1} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ , then :

(a)  $a = b = 1$

(b)  $a = \cos 2\theta, b = \sin 2\theta$

(c)  $a = \sin 2\theta, b = \cos 2\theta$

(d)  $a = 1, b = \sin 2\theta$

11. A square matrix  $P$  satisfies  $P^2 = I - P$ , where  $I$  is identity matrix. If  $P^n = 5I - 8P$ , then  $n$  is :

(a) 4

(b) 5

(c) 6

(d) 7

12. Let matrix  $A = \begin{bmatrix} x & y & -z \\ 1 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$  where  $x, y, z \in N$ . If  $\det. (\text{adj. } A) = 2^8 \cdot 3^4$  then the number

of such matrices  $A$  is :

[Note :  $\text{adj. } A$  denotes adjoint of square matrix  $A$ .]

(a) 220

(b) 45

(c) 55

(d) 110

13. If  $A$  is a  $2 \times 2$  non singular matrix, then  $\text{adj} (\text{adj } A)$  is equal to :

(a)  $A^2$

(b)  $A$

(c)  $A^{-1}$

(d)  $(A^{-1})^2$

14.  $A = \begin{bmatrix} a & b \\ b & -a \end{bmatrix}$  and  $MA = A^{2m}$ ,  $m \in N, a, b \in R$ , for some matrix  $M$ , then which one of the following is correct :

(a)  $M = \begin{bmatrix} a^{2m} & b^{2m} \\ b^{2m} & -a^{2m} \end{bmatrix}$

(b)  $M = (a^2 + b^2)^m \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(c)  $M = (a^m + b^m) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(d)  $M = (a^2 + b^2)^{m-1} \begin{bmatrix} a & b \\ b & -a \end{bmatrix}$



### Exercise-2 : One or More than One Answer is/are Correct

1. If  $A$  and  $B$  are two orthogonal matrices of order  $n$  and  $\det(A) + \det(B) = 0$ , then which of the following must be correct ?
- (a)  $\det(A+B) = \det(A) + \det(B)$       (b)  $\det(A+B) = 0$   
 (c)  $A$  and  $B$  both are singular matrices      (d)  $A+B=0$
2. Let  $M$  be a  $3 \times 3$  matrix satisfying  $M^3 = 0$ . Then which of the following statement(s) are true:
- (a)  $\left| \frac{1}{2}M^2 + M + I \right| \neq 0$       (b)  $\left| \frac{1}{2}M^2 - M + I \right| = 0$   
 (c)  $\left| \frac{1}{2}M^2 + M + I \right| = 0$       (d)  $\left| \frac{1}{2}M^2 - M + I \right| \neq 0$
3. Let  $A_\alpha = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$ , then :
- (a)  $A_{\alpha+\beta} = A_\alpha A_\beta$       (b)  $A_\alpha^{-1} = A_{-\alpha}$   
 (c)  $A_\alpha^{-1} = -A_\alpha$       (d)  $A_\alpha^2 = -I$
4.  $A^3 - 2A^2 - A + 2I = 0$  if  $A =$
- (a)  $I$       (b)  $2I$       (c)  $\begin{bmatrix} 2 & -1 & 2 \\ -1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$       (d)  $\begin{bmatrix} 2 & 1 & -2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$
5. Let  $A$  be a  $3 \times 3$  symmetric invertible matrix with real positive elements. Then the number of zero elements in  $A^{-1}$  are less than or equal to :
- (a) 0      (b) 1      (c) 2      (d) 3

### Answers

1.	(a, b)	2.	(a, d)	3.	(a, b)	4.	(a, b, c, d)	5.	(d)
----	--------	----	--------	----	--------	----	--------------	----	-----

**Exercise-3 : Matching Type Problems**

1. Consider a square matrix  $A$  of order 2 which has its elements as 0, 1, 2 and 4. Let  $N$  denotes the number of such matrices.

	Column-I		Column-II
(A)	Possible non-negative value of $\det(A)$ is	(P)	2
(B)	Sum of values of determinants corresponding to $N$ matrices is	(Q)	4
(C)	If absolute value of $(\det(A))$ is least, then possible value of $ \text{adj}(\text{adj}(\text{adj} A)) $	(R)	-2
(D)	If $\det(A)$ is least, then possible value of $\det(4A^{-1})$ is	(S)	0
		(T)	8

2.

	Column-I		Column-II
(A)	If $A$ is an idempotent matrix and $I$ is an identify matrix of the same order, then the value of $n$ , such that $(A + I)^n = I + 127A$ is	(P)	9
(B)	If $(I - A)^{-1} = I + A + A^2 + \dots + A^7$ , then $A^n = O$ where $n$ is	(Q)	10
(C)	If $A$ is matrix such that $a_{ij} = (i + j)(i - j)$ , then $A$ is singular if order of matrix is	(R)	7
(D)	If a non-singular matrix $A$ is symmetric, such that $A^{-1}$ is also symmetric, then order of $A$ can be	(S)	8

3.

	Column-I		Column-II
(A)	Number of ordered pairs $(x, y)$ of real numbers satisfying $\sin x + \cos y = 0$ , $\sin^2 x + \cos^2 y = \frac{1}{2}$ , $0 < x < \pi$ and $0 < y < \pi$ , is equal to	(P)	0
(B)	Given $\vec{a}$ , $\vec{b}$ and $\vec{c}$ are three vectors such that $\vec{b}$ and $\vec{c}$ are unit like vectors and $ \vec{a}  = 4$ . If $\vec{a} + \lambda \vec{c} = 2\vec{b}$ then the sum of all possible values of $\lambda$ is equal to	(Q)	2

(C)	If $P = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix}$ , $10Q = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & t \\ 1 & -2 & 3 \end{bmatrix}$ and $Q = P^{-1}$ , then the value of $t$ is equal to	(R)	4
(D)	If $y = \tan u$ where $u = v - \frac{1}{v}$ and $v = \ln x$ , then the value of $\frac{dy}{dx}$ at $x = e$ is equal to $\lambda$ then $[\lambda]$ is equal to (where $[\ ]$ denotes greatest integer function)	(S)	5

4.

Column-I		Column-II	
(A)	If $P$ and $Q$ are variable points on $C_1 : x^2 + y^2 = 4$ and $C_2 : x^2 + y^2 - 8x - 6y + 24 = 0$ respectively then the maximum value of $PQ$ , is equal to	(P)	1
(B)	Let $P, Q, R$ be invertible matrices of second order such that $A = PQ^{-1}, B = QR^{-1}, C = RP^{-1}$ , then the value of $\det. (ABC + BCA + CAB)$ is equal to	(Q)	2
(C)	The perpendicular distance of the point whose position vector is $(1, 3, 5)$ from the line $\vec{r} = \hat{i} + 2\hat{j} + 3\hat{k} + \lambda(\hat{i} + 2\hat{j} + 2\hat{k})$ is equal to	(R)	9
(D)	Let $f(x)$ be a continuous function in $[-1, 1]$ such that $f(x) = \begin{cases} \frac{\ln(px^2 + qx + r)}{x^2} & ; -1 \leq x < 0 \\ 1 & ; x = 0 \\ \frac{\sin(e^{x^2} - 1)}{x^2} & ; 0 < x \leq 1 \end{cases}$ then the value of $(p + q + r)$ , is equal to	(S)	8

5.

Column-I		Column-II	
(A)	$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \left( 1 + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \dots + \frac{1}{\sqrt{n}} \right)$ has the value equal to	(P)	1

<p><b>(B)</b> Let <math>A = [a_{ij}]</math> be a <math>3 \times 3</math> matrix where</p> $a_{ij} = \begin{cases} 2 \cos t; & \text{if } i = j \\ 1; & \text{if }  i - j  = 1 \\ 0; & \text{otherwise} \end{cases}$ <p>then maximum value of <math>\det(A)</math> is</p>	<p><b>(Q)</b> 2</p>
<p><b>(C)</b> Let <math>f(x) = x^3 + px^2 + qx + 6</math>; where <math>p, q \in R</math> and <math>f'(x) &lt; 0</math> in largest possible interval <math>\left(-\frac{5}{3}, -1\right)</math> then value of <math>q - p</math> is</p>	<p><b>(R)</b> 3</p>
<p><b>(D)</b> If <math>4^x - 2^{x+2} + 5 +   b - 1  - 3  =  \sin y </math>; <math>x, y, b \in R</math> then the sum of the possible values of <math>b</math> is <math>\lambda</math> then <math>(\lambda + 1)</math> equals</p>	<p><b>(S)</b> 4</p>

### Answers

1.  $A \rightarrow P, Q, T; B \rightarrow S; C \rightarrow P, R; D \rightarrow R$
2.  $A \rightarrow R; B \rightarrow P, Q, S; C \rightarrow P, R; D \rightarrow P, Q, R, S$
3.  $A \rightarrow Q; B \rightarrow R; C \rightarrow S; D \rightarrow P$
4.  $A \rightarrow S; B \rightarrow R; C \rightarrow P; D \rightarrow Q$
5.  $A \rightarrow Q; B \rightarrow S; C \rightarrow P; D \rightarrow R$

### Exercise-4 : Subjective Type Problems

1.  $A$  and  $B$  are two square matrices. Such that  $A^2B = BA$  and if  $(AB)^{10} = A^k \cdot B^{10}$ . Find the value of  $k - 1020$ .
2. Let  $A_n$  and  $B_n$  be square matrices of order 3, which are defined as :  
 $A_n = [a_{ij}]$  and  $B_n = [b_{ij}]$  where  $a_{ij} = \frac{2i+j}{3^{2n}}$  and  $b_{ij} = \frac{3i-j}{2^{2n}}$  for all  $i$  and  $j$ ,  $1 \leq i, j \leq 3$ .  
 If  $l = \lim_{n \rightarrow \infty} \text{Tr.} (3A_1 + 3^2A_2 + 3^3A_3 + \dots + 3^nA_n)$  and  
 $m = \lim_{n \rightarrow \infty} \text{Tr.} (2B_1 + 2^2B_2 + 2^3B_3 + \dots + 2^nB_n)$ , then find the value of  $\frac{l+m}{3}$   
 [Note : Tr. (P) denotes the trace of matrix P]
3. Let  $A$  be a  $2 \times 3$  matrix whereas  $B$  be a  $3 \times 2$  matrix. If  $\det. (AB) = 4$ , then the value of  $\det. (BA)$ , is :
4. Find the maximum value of the determinant of an arbitrary  $3 \times 3$  matrix  $A$ , each of whose entries  $a_{ij} \in \{-1, 1\}$ .
5. The set of natural numbers is divided into array of rows and columns in the form of matrices as  
 $A_1 = [1], A_2 = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}, A_3 = \begin{bmatrix} 6 & 7 & 8 \\ 9 & 10 & 11 \\ 12 & 13 & 14 \end{bmatrix}$  and so on. Let the trace of  $A_{10}$  be  $\lambda$ . Find unit digit of  $\lambda$ ?

### Answers

1.	3	2.	7	3.	0	4.	4	5.	5				
----	---	----	---	----	---	----	---	----	---	--	--	--	--

□□□

## Exercise-1 : Single Choice Problems

- The number of 3-digit numbers containing the digit 7 exactly once :  
 (a) 225                      (b) 220                      (c) 200                      (d) 180
- Let  $A = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$ ,  $B = \{y_1, y_2, y_3, y_4\}$ . The total number of function  $f : A \rightarrow B$  that are onto and there are exactly three elements  $x$  in  $A$  such that  $f(x) = y_1$  is :  
 (a) 11088                      (b) 10920                      (c) 13608                      (d) None of these
- The number of arrangements of the word "IDIOTS" such that vowels are at the places which form three consecutive terms of an A.P is :  
 (a) 36                      (b) 72                      (c) 24                      (d) 108
- Consider all the 5 digit numbers where each of the digits is chosen from the set  $\{1, 2, 3, 4\}$ . Then the number of numbers, which contain all the four digits is :  
 (a) 240                      (b) 244                      (c) 586                      (d) 781
- How many ways are there to arrange the letters of the word "GARDEN" with the vowels in alphabetical order ?  
 (a) 120                      (b) 480                      (c) 360                      (d) 240
- If  $\alpha \neq \beta$  but  $\alpha^2 = 5\alpha - 3$  and  $\beta^2 = 5\beta - 3$  then the equation having  $\alpha/\beta$  and  $\beta/\alpha$  as its roots is :  
 (a)  $3x^2 - 19x + 3 = 0$                       (b)  $3x^2 + 19x - 3 = 0$   
 (c)  $3x^2 - 19x - 3 = 0$                       (d)  $x^2 - 5x + 3 = 0$
- A student is to answer 10 out of 13 questions in an examination such that he must choose at least 4 from the first five questions. The number of choices available to him is :  
 (a) 140                      (b) 196                      (c) 280                      (d) 346
- Let set  $A = \{1, 2, 3, \dots, 22\}$ . Set  $B$  is a subset of  $A$  and  $B$  has exactly 11 elements, find the sum of elements of all possible subsets  $B$ .  
 (a)  $252 \cdot {}^{21}C_{11}$                       (b)  $230 \cdot {}^{21}C_{10}$   
 (c)  $253 \cdot {}^{21}C_9$                       (d)  $253 \cdot {}^{21}C_{10}$



9. The value of  $\left[ \frac{2009! + 2006!}{2008! + 2007!} \right] =$   
 ([.] denotes greatest integer function.)  
 (a) 2009 (b) 2008 (c) 2007 (d) 1
10. If  $p_1, p_2, p_3, \dots, p_{m+1}$  are distinct prime numbers. Then the number of factors of  $p_1^n p_2 p_3 \dots p_{m+1}$  is :  
 (a)  $m(n+1)$  (b)  $(n+1)2^m$  (c)  $n \cdot 2^m$  (d)  $2^{nm}$
11. A basket ball team consists of 12 pairs of twin brothers. On the first day of training, all 24 players stand in a circle in such a way that all pairs of twin brothers are neighbours. Number of ways this can be done is :  
 (a)  $(12)! 2^{11}$  (b)  $(11)! 2^{12}$  (c)  $(12)! 2^{12}$  (d)  $(11)! 2^{11}$
12. Let 'm' denotes the number of four digit numbers such that the left most digit is odd, the second digit is even and all four digits are different and 'n' denotes the number of four digit numbers such that left most digit is even, second digit is odd and all four digits are different. If  $m = nk$ , then k equals :  
 (a)  $\frac{4}{5}$  (b)  $\frac{3}{4}$  (c)  $\frac{5}{4}$  (d)  $\frac{4}{3}$
13. The number of three digit numbers of the form  $xyz$  such that  $x < y$  and  $z \leq y$  is :  
 (a) 156 (b) 204 (c) 240 (d) 276
14. A and B are two sets and their intersection has 3 elements. If A has 1920 more subsets than B has, then the number of elements of A union B is :  
 (a) 12 (b) 14 (c) 15 (d) 16
15. All possible 120 permutations of WDSMC are arranged in dictionary order, as if each were an ordinary five-letter word. The last letter of the 86<sup>th</sup> word in the list, is :  
 (a) W (b) D (c) M (d) C
16. The number of permutation of all the letters AAAABBBBC in which the A's appear together in a block of 4 letters or the B's appear together in a block of 3 letters is :  
 (a) 44 (b) 50 (c) 60 (d) 89
17. Number of zero's at the ends of  $\prod_{n=5}^{30} (n)^{n+1}$  is :  
 (a) 111 (b) 147 (c) 137 (d) None of these
18. The number of positive integral pairs  $(x, y)$  satisfying the equation  $x^2 - y^2 = 3370$  is :  
 (a) 0 (b) 1 (c) 2 (d) 4
19. The number of ways of selecting 'n' things out of '3n' things of which 'n' are of one kind and alike and 'n' are of second kind and alike and the rest unlike is :  
 (a)  $n 2^{n-1}$  (b)  $(n-1) 2^{n-1}$  (c)  $(n+1) 2^{n-1}$  (d)  $(n+2) 2^{n-1}$

20. If  $x, y, z$  are three natural numbers in A.P such that  $x + y + z = 30$ , then the possible number of ordered triplet  $(x, y, z)$  is :  
 (a) 18 (b) 19 (c) 20 (d) 21
21. A dice is rolled 4 times, the numbers appearing are listed. The number of different throws, such that the largest number appearing in the list is not 4, is : :  
 (a) 175 (b) 625 (c) 1040 (d) 1121
22. Let  $m$  denotes the number of ways in which 5 boys and 5 girls can be arranged in a line alternately and  $n$  denotes the number of ways in which 5 boys and 5 girls can be arranged in a circle so that no two boys are together. if  $m = kn$  then the value of  $k$  is :  
 (a) 2 (b) 5 (c) 6 (d) 10
23. Number of ways in which 4 students can sit in 7 chair in a row, if there is no empty chair between any two students is :  
 (a) 24 (b) 28 (c) 72 (d) 96
24. Number of zero's at the ends of  $\prod_{n=5}^{30} (n)^{n+1}$  is :  
 (a) 111 (b) 147 (c) 137 (d) None
25. The number of words of four letters consisting of equal number of vowels and consonants (of english language) with repetition permitted is :  
 (a) 51030 (b) 50030 (c) 63050 (d) 66150
26. Ten different letters of an alphabet are given. Words with five letters are formed with these given letters. Then the number of words which have atleast one letter repeated is :  
 (a) 30240 (b) 69760 (c) 69780 (d) 99784
27. Number of four digit numbers in which at least one digit occurs more than once, is:  
 (a) 4464 (b) 4644 (c) 4446 (d) 6444
28. In a game of minesweeper, a number on a square denotes the number of mines that share at least one vertex with that square. A square with a number may not have a mine, and the blank squares are undetermined. In how many ways can the mines be placed in the given configuration on blank squares:

	2		1		2

- (a) 120 (b) 105 (c) 95 (d) 100
29. Let the product of all the divisors of 1440 be  $P$ . If  $P$  is divisible by  $24^x$ , then the maximum value of  $x$  is :  
 (a) 28 (b) 30 (c) 32 (d) 36


- 30.** Let  $N$  be the number of 4-digit numbers which contain not more than 2 different digits. The sum of the digits of  $N$  is :  
 (a) 18 (b) 19 (c) 20 (d) 21
- 31.** The number of different permutations of all the letters of the word PERMUTATION such that any two consecutive letters in the arrangement are neither both vowels nor both identical is :  
 (a)  $63 \times \underline{6} \times \underline{5}$  (b)  $8 \times \underline{6} \times \underline{5}$  (c)  $57 \times \underline{5} \times \underline{5}$  (d)  $7 \times \underline{7} \times \underline{5}$
- 32.** A batsman can score 0, 1, 2, 3, 4 or 6 runs from a ball. The number of different sequences in which he can score exactly 30 runs in an over of six balls :  
 (a) 4 (b) 72 (c) 56 (d) 71
- 33.** A batsman can score 0, 2, 3, or 4 runs for each ball he receives. If  $N$  is the number of ways of scoring a total of 20 runs in one over of six balls, then  $N$  is divisible by:  
 (a) 5 (b) 7 (c) 14 (d) 16
- 34.** The number of non-negative integral solutions of the equation  $x + y + z = 5$  is :  
 (a) 20 (b) 19 (c) 21 (d) 25
- 35.** The number of solutions of the equation  $x_1 + x_2 + x_3 + x_4 + x_5 = 101$ , where  $x_i$ 's are odd natural numbers is :  
 (a)  $^{105}C_4$  (b)  $^{52}C_5$  (c)  $^{52}C_4$  (d)  $^{50}C_4$
- 36.** An ordinary dice is rolled 4 times, numbers appearing on them are listed. The number of different throws, such that the largest number appearing on them is NOT 4, is :  
 (a) 175 (b) 625 (c) 1121 (d) 1040
- 37.** Number of four letter words can be formed using the letters of word VIBRANT if letter V is must included, are :  
 (a) 840 (b) 480 (c) 120 (d) 240
- 38.** The number of rectangles that can be obtained by joining four of the twelve vertices of a 12-sided regular polygon is :  
 (a) 66 (b) 30 (c) 24 (d) 15
- 39.** Number of five digit integers, with sum of the digits equal to 43 are :  
 (a) 5 (b) 10 (c) 15 (d) 35

## Answers

<b>1.</b>	(a)	<b>2.</b>	(d)	<b>3.</b>	(d)	<b>4.</b>	(a)	<b>5.</b>	(c)	<b>6.</b>	(a)	<b>7.</b>	(b)	<b>8.</b>	(d)	<b>9.</b>	(b)	<b>10.</b>	(b)
<b>11.</b>	(b)	<b>12.</b>	(c)	<b>13.</b>	(d)	<b>14.</b>	(c)	<b>15.</b>	(b)	<b>16.</b>	(a)	<b>17.</b>	(c)	<b>18.</b>	(a)	<b>19.</b>	(d)	<b>20.</b>	(b)
<b>21.</b>	(d)	<b>22.</b>	(d)	<b>23.</b>	(d)	<b>24.</b>	(c)	<b>25.</b>	(d)	<b>26.</b>	(b)	<b>27.</b>	(a)	<b>28.</b>	(c)	<b>29.</b>	(b)	<b>30.</b>	(a)
<b>31.</b>	(c)	<b>32.</b>	(d)	<b>33.</b>	(d)	<b>34.</b>	(c)	<b>35.</b>	(c)	<b>36.</b>	(c)	<b>37.</b>	(b)	<b>38.</b>	(d)	<b>39.</b>	(c)		

**Exercise-2 : One or More than One Answer is/are Correct**

- The number of 5 letter words formed with the letters of the word CALCULUS is divisible by :  
 (a) 2                      (b) 3                      (c) 5                      (d) 7
- The coefficient of  $x^{50}$  in the expansion of  $\sum_{k=0}^{100} {}^{100}C_k (x-2)^{100-k} 3^k$  is also equal to :  
 (a) Number of ways in which 50 identical books can be distributed in 100 students, if each student can get atmost one book.  
 (b) Number of ways in which 100 different white balls and 50 identical red balls can be arranged in a circle, if no two red balls are together.  
 (c) Number of dissimilar terms in  $(x_1 + x_2 + x_3 + \dots + x_{50})^{51}$ .  
 (d)  $\frac{2 \cdot 6 \cdot 10 \cdot 14 \dots 198}{50!}$
- Number of ways in which the letters of the word "NATION" can be filled in the given figure such that no row remains empty and each box contains not more than one letter, are :




- (a)  $11 \mid 6$                       (b)  $12 \mid 6$                       (c)  $13 \mid 6$                       (d)  $14 \mid 6$
- Let  $a, b, c, d$  be non zero distinct digits. The number of 4 digit numbers  $abcd$  such that  $ab + cd$  is even is divisible by :  
 (a) 3                      (b) 4                      (c) 7                      (d) 11

**Answers**

1.	(a, b, c)	2.	(a, d)	3.	(c)	4.	(a, b, d)			
----	-----------	----	--------	----	-----	----	-----------	--	--	--



 **Exercise-4 : Matching Type Problems**

1. All letters of the word BREAKAGE are to be jumbled. The number of ways of arranging them so that :

Column-I		Column-II	
(A)	The two A's are not together	(P)	720
(B)	The two E's are together but not two A's	(Q)	1800
(C)	Neither two A's nor two E's are together	(R)	5760
(D)	No two vowels are together	(S)	6000
		(T)	7560

2. Consider the letters of the word MATHEMATICS. Set of repeating letters = { M, A, T }, set of non repeating letters = { H, E, I, C, S }:


Column-I		Column-II	
(A)	The number of words taking all letters of the given word such that atleast one repeating letter is at odd position is	(P)	$28 \cdot (7!)$
(B)	The number of words formed taking all letters of the given word in which no two vowels are together is	(Q)	$\frac{(11)!}{(2!)^3}$
(C)	The number of words formed taking all letters of the given word such that in each word both M's are together and both T's are together but both A's are not together is	(R)	$210(7!)$
(D)	The number of words formed taking all letters of the given word such that relative order of vowels and consonants does not change is	(S)	$840(7!)$
		(T)	$\frac{4!7!}{(2!)^3}$

### Answers

- |    |                            |
|----|----------------------------|
| 1. | A → T; B → Q; C → R; D → P |
| 2. | A → Q; B → R; C → P; D → T |

### Exercise-5 : Subjective Type Problems

- The number of ways in which eight digit number can be formed using the digits from 1 to 9 without repetition if first four places of the numbers are in increasing order and last four places are in decreasing order is  $N$ , then find the value of  $\frac{N}{70}$ .
- Number of ways in which the letters of the word DECISIONS be arranged so that letter N be somewhere to the right of the letter "D" is  $\frac{9}{\lambda}$ . Find  $\lambda$ .
- There are 10 stations enroute. A train has to be stopped at 3 of them. Let  $N$  be the ways in which the train can be stopped if atleast two of the stopping stations are consecutive. Find the value of  $\sqrt{N}$ .
- There are 10 girls and 8 boys in a class room including Mr. Ravi, Ms. Rani and Ms. Radha. A list of speakers consisting of 8 girls and 6 boys has to be prepared. Mr. Ravi refuses to speak if Ms. Rani is a speaker. Ms. Rani refuses to speak if Ms. Radha is a speaker. The number of ways the list can be prepared is a 3 digit number  $n_1 n_2 n_3$ , then  $|n_3 + n_2 - n_1| =$
- Nine people sit around a round table. The number of ways of selecting four of them such that they are not from adjacent seats, is
- Let the number of arrangements of all the digits of the numbers 12345 such that atleast 3 digits will not come in it's original position is  $N$ . Then the unit digit of  $N$  is
- The number of triangles with each side having integral length and the longest side is of 11 units is equal to  $k^2$ , then the value of ' $k$ ' is equal to
- 8 clay targets are arranged as shown. If  $N$  be the number of different ways they can be shot (one at a time) if no target can be shot until the target(s) below it have been shot. Find the ten's digit of  $N$ .



- There are  $n$  persons sitting around a circular table. They start singing a 2 minute song in pairs such that no two persons sitting together will sing together. This process is continued for 28 minutes. Find  $n$ .
- The number of ways to choose 7 distinct natural numbers from the first 100 natural numbers such that any two chosen numbers differ atleast by 7 can be expressed as  ${}^n C_7$ . Find the number of divisors of  $n$ .
- Four couples (husband and wife) decide to form a committee of four members. The number of different committees that can be formed in which no couple finds a place is  $\lambda$ , then the sum of digits of  $\lambda$  is :


13. The number of ways in which  $2n$  objects of one type,  $2n$  of another type and  $2n$  of a third type can be divided between 2 persons so that each may have  $3n$  objects is  $\alpha n^2 + \beta n + \gamma$ . Find the value of  $(\alpha + \beta + \gamma)$ .
14. Let  $N$  be the number of integral solution of the equation  $x + y + z + w = 15$  where  $x \geq 0, y > 5, z \geq 2$  and  $w \geq 1$ . Find the unit digit of  $N$ .

## Answers

1.	9	2.	8	3.	8	4.	8	5.	5	6.	9	7.	9
8.	6	9.	6	10.	7	11.	7	12.	7	13.	7	14.	4

□□□




**Exercise-1 : Single Choice Problems**

- Let  $N = 2^{1224} - 1$ ,  $\alpha = 2^{153} + 2^{77} + 1$  and  $\beta = 2^{408} - 2^{204} + 1$ . Then which of the following statement is correct ?
  - $\alpha$  divides  $N$  but  $\beta$  does not
  - $\beta$  divides  $N$  but  $\alpha$  does not
  - $\alpha$  and  $\beta$  both divide  $N$
  - neither  $\alpha$  nor  $\beta$  divides  $N$
- If  $(1 + x + x^2)^n = \sum_{r=0}^{2n} a_r x^r$ , then  $a_r - {}^n C_1 \cdot a_{r-1} + {}^n C_2 a_{r-2} - {}^n C_3 a_{r-3} + \dots + (-1)^r {}^n C_r a_0$  is equal to : ( $r$  is not multiple of 3)
  - 0
  - ${}^n C_r$
  - $a_r$
  - 1
- The coefficient of the middle term in the binomial expansion in powers of  $x$  of  $(1 + \alpha x)^4$  and of  $(1 - \alpha x)^6$  is the same if  $\alpha$  equals :
  - $-\frac{5}{3}$
  - $\frac{3}{5}$
  - $-\frac{3}{10}$
  - $\frac{10}{3}$
- If  $(1 + x)^{2010} = C_0 + C_1 x + C_2 x^2 + \dots + C_{2010} x^{2010}$  then the sum of series  $C_2 + C_5 + C_8 + \dots + C_{2009}$  equals to :
  - $\frac{1}{2}(2^{2010} - 1)$
  - $\frac{1}{3}(2^{2010} - 1)$
  - $\frac{1}{2}(2^{2009} - 1)$
  - $\frac{1}{3}(2^{2009} - 1)$
- Let  $\alpha_n = (2 + \sqrt{3})^n$ . Find  $\lim_{n \rightarrow \infty} (\alpha_n - [\alpha_n])$  ( $[\cdot]$  denotes greatest integer function)
  - 1
  - $\frac{1}{2}$
  - $\frac{1}{3}$
  - $\frac{2}{3}$
- The number  $N = {}^{20}C_7 - {}^{20}C_8 + {}^{20}C_9 - {}^{20}C_{10} + \dots - {}^{20}C_{20}$  is not divisible by :
  - 3
  - 7
  - 11
  - 19

7. The value of the expression  $\log_2 \left( 1 + \frac{1}{2} \sum_{k=1}^{11} {}^{12}C_k \right)$ :
- (a) 11                      (b) 12                      (c) 13                      (d) 14
8. The constant term in the expansion of  $\left( x + \frac{1}{x^3} \right)^{12}$  is :
- (a) 26                      (b) 169                      (c) 260                      (d) 220
9. If  $\frac{3}{4!} + \frac{4}{5!} + \frac{5}{6!} + \dots + 50$  term  $= \frac{1}{3!} - \frac{1}{(k+3)!}$ , then sum of coefficients in the expansion  $(1 + 2x_1 + 3x_2 + \dots + 100x_{100})^k$  is :
- (where  $x_1, x_2, x_3, \dots, x_{100}$  are independent variables)
- (a)  $(5050)^{49}$                       (b)  $(5050)^{51}$   
(c)  $(5050)^{52}$                       (d)  $(5050)^{50}$
10. **Statement-1:** The remainder when  $(128)^{(128)^{128}}$  is divided by 7 is 3.  
**because**  
**Statement-2:**  $(128)^{128}$  when divided by 3 leaves the remainder 1.
- (a) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
(b) Statement-1 is true, statement-2 is true and statement-2 is not the correct explanation for statement-1.  
(c) Statement-1 is true, statement-2 is false.  
(d) Statement-1 is false, statement-2 is true.
11. If  $n > 3$ , then  $xyz {}^n C_0 - (x-1)(y-1)(z-1) {}^n C_1 + (x-2)(y-2)(z-2) {}^n C_2 - (x-3)(y-3)(z-3) {}^n C_3 + \dots + (-1)^n (x-n)(y-n)(z-n) {}^n C_n$  equals :
- (a)  $xyz$                       (b)  $x+y+z$   
(c)  $xy+yz+zx$                       (d) 0
12. If  $\alpha_1, \alpha_2, \dots, \alpha_n$  are the  $n; n^{\text{th}}$  roots of unity,  $\alpha_r = e^{\frac{i2(r-1)\pi}{n}}, r = 1, 2, \dots, n$  then  ${}^n C_1 \alpha_1 + {}^n C_2 \alpha_2 + \dots + {}^n C_n \alpha_n$  is equal to:
- (a)  $\left( 1 + \frac{\alpha_2}{\alpha_1} \right)^n - 1$       (b)  $\frac{\alpha_1}{2} [(1 + \alpha_1)^n - 1]$       (c)  $\frac{\alpha_1 + \alpha_{n-1} - 1}{2}$       (d)  $(\alpha_1 + \alpha_{n-1})^n - 1$
13. The remainder when  $2^{30} \cdot 3^{20}$  is divided by 7 is :
- (a) 1                      (b) 2                      (c) 4                      (d) 6
14.  ${}^{26}C_0 + {}^{26}C_1 + {}^{26}C_2 + \dots + {}^{26}C_{13}$  is equal to :
- (a)  $2^{25} - \frac{1}{2} \cdot {}^{26}C_{13}$       (b)  $2^{25} + \frac{1}{2} \cdot {}^{26}C_{13}$       (c)  $2^{13}$                       (d)  $2^{26} + \frac{1}{2} \cdot {}^{26}C_{13}$




**Exercise-2 : One or More than One Answer is/are Correct**

- The number  $N = {}^{20}C_7 - {}^{20}C_8 + {}^{20}C_9 - {}^{20}C_{10} + \dots - {}^{20}C_{20}$  is divisible by :  
 (a) 3 (b) 4 (c) 7 (d) 19
- If  $(1 + x + x^2 + x^3)^{100} = a_0 + a_1x + a_2x^2 + \dots + a_{300}x^{300}$  then which of the following statement(s) is/are correct ?  
 (a)  $a_1 = 100$   
 (b)  $a_0 + a_1 + a_2 + \dots + a_{300}$  is divisible by 1024  
 (c) coefficients equidistant from beginning and end are equal  
 (d)  $a_0 + a_2 + a_4 + \dots + a_{300} = a_1 + a_3 + a_5 + \dots + a_{299}$
- $\sum_{r=0}^4 (-1)^r {}^{16}C_r$  is divisible by :  
 (a) 5 (b) 7 (c) 11 (d) 13
- The expansion of  $\left(\sqrt{x} + \frac{1}{2\sqrt[4]{x}}\right)^n$  is arranged in decreasing powers of  $x$ . If coefficient of first three terms form an A.P. then in expansion, the integral powers of  $x$  are :  
 (a) 0 (b) 2 (c) 4 (d) 8
- Let  $(1 + x^2)^2(1 + x)^n = \sum_{k=0}^{n+4} a_k x^k$ . If  $a_1, a_2, a_3$  are in AP then  $n$  is (given that  ${}^nC_r = 0$ , if  $n < r$ ):  
 (a) 6 (b) 4 (c) 3 (d) 2
- $\sum_{i=0}^n \sum_{j=0}^n \sum_{k=0}^n \binom{n}{i} \binom{n}{j} \binom{n}{k}, \binom{n}{r} = {}^nC_r$  :  
 (a) is less than 500 if  $n = 3$  (b) is greater than 600 if  $n = 3$   
 (c) is less than 5000 if  $n = 4$  (d) is greater than 4000 if  $n = 4$
- If  ${}^{100}C_6 + 4 \cdot {}^{100}C_7 + 6 \cdot {}^{100}C_8 + 4 \cdot {}^{100}C_9 + {}^{100}C_{10}$  has the value equal to  ${}^xC_y$ ; then the possible value(s) of  $x + y$  can be :  
 (a) 112 (b) 114 (c) 196 (d) 198
- If the co-efficient of  $x^{2r}$  is greater than half of the co-efficient of  $x^{2r+1}$  in the expansion of  $(1 + x)^{15}$ ; then the possible value of 'r' equal to :  
 (a) 5 (b) 6 (c) 7 (d) 8
- Let  $f(x) = 1 + x^{111} + x^{222} + x^{333} + \dots + x^{999}$  then  $f(x)$  is divisible by  
 (a)  $x + 1$  (b)  $x$   
 (c)  $x - 1$  (d)  $1 + x^{222} + x^{444} + x^{666} + x^{888}$

**Answers**

1.	(a, b, c, d)	2.	(a, b, c, d)	3.	(a, b, d)	4.	(a, c, d)	5.	(b, c, d)	6.	(c, d)
7.	(b, d)	8.	(a, b, c)	9.	(a, d)						


**Exercise-3 : Matching Type Problems**

1.

	Column-I		Column-II
(A)	If ${}^{n-1}C_r = (k^2 - 3){}^nC_{r+1}$ and $k \in R^+$ , then least value of $5[k]$ is (where $[\cdot]$ represents greatest integer function)	(P)	10
(B)	$\sum_{i=0}^m {}^{20}C_i \cdot {}^{40}C_{m-i}$ , where ${}^nC_r = 0$ if $r > n$ , is maximum when $\frac{m}{5}$ is	(Q)	5
(C)	Number of non-negative integral solutions of inequation $x + y + z \leq 4$ is	(R)	35
(D)	Let $A = \{1, 2, 3, 4, 5\}$ , $f : A \rightarrow A$ , The number of onto functions such that $f(x) = x$ for atleast 3 distinct $x \in A$ , is not a multiple of	(S)	6
		(T)	12

2.

	Column-I		Column-II
(A)	Number of real solution of $(x^2 + 6x + 7)^2 + 6(x^2 + 6x + 7) + 7 = x$ is/are	(P)	15
(B)	If $P = \sum_{r=0}^n {}^nC_r$ ; $q = \sum_{r=0}^m {}^mC_r$ , $(15)^r$ ( $m, n \in N$ ) and if $P = q$ and $m, n$ are least then $m + n =$	(Q)	5
(C)	Remainder when $1! + 3! + 5! + \dots + 2011!$ is divided by 56 is	(R)	3
(D)	Inequality $\left  1 - \frac{ x }{1+ x } \right  \geq \frac{1}{2}$ holds for $x$ , then number of integral values of 'x' is/are	(S)	0

3. Match the following

	Column-I		Column-II
(A)	If the sum of first 84 terms of the series $\frac{4 + \sqrt{3}}{1 + \sqrt{3}} + \frac{8 + \sqrt{15}}{\sqrt{3} + \sqrt{5}} + \frac{12 + \sqrt{35}}{\sqrt{5} + \sqrt{7}} + \dots$ is $549k$ , then $k$ is equal to	(P)	3

<b>(B)</b>	If $x, y \in \mathbb{R}$ , $x^2 + y^2 - 6x + 8y + 24 = 0$ , the greatest value of $\frac{16}{5} \cos^2(\sqrt{x^2 + y^2}) - \frac{24}{5} \sin(\sqrt{x^2 + y^2})$ is	<b>(Q)</b>	2
<b>(C)</b>	If $(\sqrt{3} + 1)^6 + (\sqrt{3} - 1)^6 = 416$ , if $xyz = [(\sqrt{3} + 1)^6]$ , $x, y, z \in \mathbb{N}$ , (where $[ ]$ denotes the greatest integer function), then the number of ordered triplets $(x, y, z)$ is	<b>(R)</b>	5
<b>(D)</b>	If $(1+x)(1+x^2)(1+x^4)\dots(1+x^{128}) = \sum_{r=0}^n x^r$ , then $\frac{n}{85}$ is equal to	<b>(S)</b>	9

### Answers

1.  $A \rightarrow Q$ ;  $B \rightarrow S$ ;  $C \rightarrow R$ ;  $D \rightarrow P, Q, R, S, T$
2.  $A \rightarrow S$ ;  $B \rightarrow Q$ ;  $C \rightarrow P$ ;  $D \rightarrow R$
3.  $A \rightarrow Q$ ;  $B \rightarrow R$ ;  $C \rightarrow S$ ;  $D \rightarrow P$

### Exercise-4 : Subjective Type Problems

- The sum of the series  $3 \cdot {}^{2007}C_0 - 8 \cdot {}^{2007}C_1 + 13 \cdot {}^{2007}C_2 - 18 \cdot {}^{2007}C_3 + \dots$  upto 2008 terms is  $K$ , then  $K$  is :
- In the polynomial function  $f(x) = (x-1)(x^2-2)(x^3-3)\dots(x^{11}-11)$  the coefficient of  $x^{60}$  is :
- If  $\sum_{r=0}^{3n} a_r (x-4)^r = \sum_{r=0}^{3n} A_r (x-5)^r$  and  $a_k = 1 \forall k \geq 2n$  and  $\sum_{r=0}^{3n} d_r (x-8)^r = \sum_{r=0}^{3n} B_r (x-9)^r$  and  $\sum_{r=0}^{3n} d_r (x-12)^r = \sum_{r=0}^{3n} D_r (x-13)^r$  and  $d_k = 1 \forall k \geq 2n$ . The find the value of  $\frac{A_{2n} + D_{2n}}{B_{2n}}$ .
- If  $3^{101} - 2^{100}$  is divided by 11, the remainder is
- Find the hundred's digit in the co-efficient of  $x^{17}$  in the expansion of  $(1 + x^5 + x^7)^{20}$ .
- Let  $x = (3\sqrt{6} + 7)^{89}$ . If  $\{x\}$  denotes the fractional part of ' $x$ ' then find the remainder when  $x\{x\} + (x\{x\})^2 + (x\{x\})^3$  is divided by 31.
- Let  $n \in N$ ;  $S_n = \sum_{r=0}^{3n} ({}^{3n}C_r)$  and  $T_n = \sum_{r=0}^n ({}^{3n}C_{3r})$ . Find  $|S_n - 3T_n|$ .
- Find the sum of possible real values of  $x$  for which the sixth term of  $\left( 3^{\log_3 \sqrt{9^{|x-2|}}} + 7^{\frac{1}{5} \log_7 (3^{|x-2|-9})} \right)^7$  equal 567 :
- Let  $q$  be a positive integer with  $q \leq 50$ .  
If the sum  ${}^{98}C_{30} + 2 \cdot {}^{97}C_{30} + 3 \cdot {}^{96}C_{30} + \dots + 68 \cdot {}^{31}C_{30} + 69 \cdot {}^{30}C_{30} = {}^{100}C_q$   
Find the sum of the digits of  $q$ .
- The remainder when  $\left( \sum_{k=1}^5 {}^{20}C_{2k-1} \right)^6$  is divided by 11, is :
- Let  $a = 3^{\frac{1}{223}} + 1$  and for all  $n \geq 3$ , let  
 $f(n) = {}^nC_0 \cdot a^{n-1} - {}^nC_1 \cdot a^{n-2} + {}^nC_2 \cdot a^{n-3} - \dots + (-1)^{n-1} {}^nC_{n-1} \cdot a^0$ .  
If the value of  $f(2007) + f(2008) = 3^7 k$  where  $k \in N$  then find  $k$
- In the polynomial  $(x-1)(x^2-2)(x^3-3)\dots(x^{11}-11)$ , the coefficient of  $x^{60}$  is :
- Let the sum of all divisors of the form  $2^p \cdot 3^q$  (with  $p, q$  positive integers) of the number  $19^{88} - 1$  be  $\lambda$ . Find the unit digit of  $\lambda$ .

14. Find the sum of possible real values of  $x$  for which the sixth term of  $\left(3^{\log_3 \sqrt{9^{x-2}}} + 7^{\left(\frac{1}{5}\right) \log_7 (3^{x-2}-9)}\right)^7$  equals 567.
15. Let  $1 + \sum_{r=1}^{10} (3^r \cdot {}^{10}C_r + r \cdot {}^{10}C_r) = 2^{10} (\alpha \cdot 4^5 + \beta)$  where  $\alpha, \beta \in N$  and  $f(x) = x^2 - 2x - k^2 + 1$ . If  $\alpha, \beta$  lies between the roots of  $f(x) = 0$ . Then find the smallest positive integral value of  $k$ .
16. Let  $S_n = {}^nC_0 {}^nC_1 + {}^nC_1 {}^nC_2 + \dots + {}^nC_{n-1} {}^nC_n$  if  $\frac{S_{n+1}}{S_n} = \frac{15}{4}$ ; find the sum of all possible values of  $n$  ( $n \in N$ )

### Answers


<b>1.</b>	0	<b>2.</b>	1	<b>3.</b>	2	<b>4.</b>	2	<b>5.</b>	4	<b>6.</b>	0	<b>7.</b>	2
<b>8.</b>	4	<b>9.</b>	5	<b>10.</b>	3	<b>11.</b>	9	<b>12.</b>	(1)	<b>13.</b>	(4)	<b>14.</b>	(4)
<b>15.</b>	5	<b>16.</b>	6										

□□□





7. Three different numbers are selected at random from the set  $A = \{1, 2, 3, \dots, 10\}$ . Then the probability that the product of two numbers equal to the third number is  $\frac{p}{q}$ , where  $p$  and  $q$  are relatively prime positive integers then the value of  $(p + q)$  is :
- (a) 39                      (b) 40                      (c) 41                      (d) 42
8. Mr. A's T.V. has only 4 channels ; all of them quite boring so he naturally desires to switch (change) channel after every one minute. The probability that he is back to his original channel for the first time after 4 minutes can be expressed as  $\frac{m}{n}$ ; where  $m$  and  $n$  are relatively prime numbers. Then  $(m + n)$  equals :
- (a) 27                      (b) 31                      (c) 23                      (d) 33
9. Letters of the word TITANIC are arranged to form all the possible words. What is the probability that a word formed starts either with a T or a vowel ?
- (a)  $\frac{2}{7}$                       (b)  $\frac{4}{7}$                       (c)  $\frac{3}{7}$                       (d)  $\frac{5}{7}$
10. A mapping is selected at random from all mappings  $f : A \rightarrow A$  where set  $A = \{1, 2, 3, \dots, n\}$
- If the probability that mapping is injective is  $\frac{3}{32}$ , then the value of  $n$  is :
- (a) 3                      (b) 4                      (c) 8                      (d) 16
11. A 4 digit number is randomly picked from all the 4 digit numbers, then the probability that the product of its digit is divisible by 3 is :
- (a)  $\frac{107}{125}$                       (b)  $\frac{109}{125}$   
 (c)  $\frac{111}{125}$                       (d) None of these
12. To obtain a gold coin; 6 men, all of different weight, are trying to build a human pyramid as shown in the figure. Human pyramid is called "stable" if some one not in the bottom row is "supported by" each of the two closest people beneath him and no body can be supported by anybody of lower weight. Formation of 'stable' pyramid is the only condition to get a gold coin. What is the probability that they will get gold coin ?
- (a)  $\frac{1}{45}$                       (b)  $\frac{2}{45}$   
 (c)  $\frac{4}{45}$                       (d)  $\frac{1}{5}$
13. From a pack of 52 playing cards; half of the cards are randomly removed without looking at them. From the remaining cards, 3 cards are drawn randomly. The probability that all are king.



(a)  $\frac{1}{(25)(17)(13)}$

(b)  $\frac{1}{(25)(15)(13)}$

(c)  $\frac{1}{(52)(17)(13)}$

(d)  $\frac{1}{(13)(51)(17)}$

14. A bag contains 10 white and 3 black balls. Balls are drawn one by one without replacement till all the black balls are drawn. The probability that the procedure of drawing balls will come to an end at the seventh draw is :

(a)  $\frac{15}{286}$

(b)  $\frac{105}{286}$

(c)  $\frac{35}{286}$

(d)  $\frac{7}{286}$

15. Let  $S$  be the set of all function from the set  $\{1, 2, \dots, 10\}$  to itself. One function is selected from  $S$ , the probability that the selected function is one-one onto is :

(a)  $\frac{9!}{10^9}$

(b)  $\frac{1}{10}$

(c)  $\frac{100}{10!}$

(d)  $\frac{9!}{10^{10}}$

16. Two friends visit a restaurant randomly during 5 pm to 6 pm. Among the two, whoever comes first waits for 15 min and then leaves. The probability that they meet is :

(a)  $\frac{1}{4}$

(b)  $\frac{1}{16}$

(c)  $\frac{7}{16}$

(d)  $\frac{9}{16}$

17. Three numbers are randomly selected from the set  $\{10, 11, 12, \dots, 100\}$ . Probability that they form a Geometric progression with integral common ratio greater than 1 is :

(a)  $\frac{15}{{}_{91}C_3}$

(b)  $\frac{16}{{}_{91}C_3}$

(c)  $\frac{17}{{}_{91}C_3}$

(d)  $\frac{18}{{}_{91}C_3}$

## Answers

1.	(a)	2.	(c)	3.	(c)	4.	(c)	5.	(c)	6.	(d)	7.	(c)	8.	(b)	9.	(d)	10.	(b)
11.	(a)	12.	(a)	13.	(a)	14.	(a)	15.	(a)	16.	(c)	17.	(d)						

### Exercise-2 : One or More than One Answer is/are Correct

1. A consignment of 15 record players contain 4 defectives. The record players are selected at random, one by one and examined. The one examined is not put back. Then :
- (a) Probability of getting exactly 3 defectives in the examination of 8 record players is  $\frac{{}^4C_3 \cdot {}^{11}C_5}{{}^{15}C_8}$ .
- (b) Probability that 9<sup>th</sup> one examined is the last defective is  $\frac{8}{197}$ .
- (c) Probability that 9<sup>th</sup> examined record player is defective, given that there are 3 defectives in first 8 players examined is  $\frac{1}{7}$ .
- (d) Probability that 9<sup>th</sup> one examined is the last defective is  $\frac{8}{195}$ .
2. If  $A_1, A_2, A_3, \dots, A_{1006}$  be independent events such that  $P(A_i) = \frac{1}{2i}$  ( $i = 1, 2, 3, \dots, 1006$ ) and probability that none of the events occurs be  $\frac{\alpha!}{2^\alpha (\beta!)^2}$ , then :
- (a)  $\beta$  is of form  $4k + 2$ ,  $k \in I$  (b)  $\alpha = 2\beta$   
(c)  $\beta$  is a composite number (d)  $\alpha$  is of form  $4k$ ,  $k \in I$
3. A bag contains four tickets marked with 112, 121, 211, 222 one ticket is drawn at random from the bag. let  $E_i$  ( $i = 1, 2, 3$ ) denote the event that  $i^{\text{th}}$  digit on the ticket is 2. Then :
- (a)  $E_1$  and  $E_2$  are independent (b)  $E_2$  and  $E_3$  are independent  
(c)  $E_3$  and  $E_1$  are independent (d)  $E_1, E_2, E_3$  are independent
4. For two events  $A$  and  $B$  let,  $P(A) = \frac{3}{5}, P(B) = \frac{2}{3}$ , then which of the following is/are correct ?
- (a)  $P(A \cap \bar{B}) \leq \frac{1}{3}$  (b)  $P(A \cup B) \geq \frac{2}{3}$   
(c)  $\frac{4}{15} \leq P(A \cap B) \leq \frac{3}{5}$  (d)  $\frac{1}{10} \leq P(\bar{A}/B) \leq \frac{3}{5}$

### Answers

1.	(a, c, d)	2.	(a, b, c, d)	3.	(a, b, c)	4.	(a, b, c, d)				
----	-----------	----	--------------	----	-----------	----	--------------	--	--	--	--






### Exercise-4 : Matching Type Problems

1. A is a set containing  $n$  elements, A subset  $P$  (may be void also) is selected at random from set  $A$  and the set  $A$  is then reconstructed by replacing the elements of  $P$ . A subset  $Q$  (may be void also) of  $A$  is again chosen at random. The probability that

Column-I		Column-II	
(A)	Number of elements in $P$ is equal to the number of elements in $Q$ is	(P)	$\frac{{}^{2n}C_n}{4^n}$
(B)	The number of elements in $P$ is more than that in $Q$ is	(Q)	$\frac{(2^{2n} - 2^n C_n)}{2^{2n+1}}$
(C)	$P \cap Q = \phi$ is	(R)	$\frac{{}^{2n}C_{n+1}}{4^n}$
(D)	$Q$ is a subset of $P$ is	(S)	$\left(\frac{3}{4}\right)^n$
		(T)	$\frac{{}^{2n}C_n}{4^{n-1}}$

### Answers

1.  $A \rightarrow P; B \rightarrow Q; C \rightarrow S; D \rightarrow S$


**Exercise-5 : Subjective Type Problems**

- Mr. A writes an article. The article originally is error free. Each day Mr. B introduces one new error into the article. At the end of the day, Mr. A checks the article and has  $\frac{2}{3}$  chance of catching each individual error still in the article. After 3 days, the probability that the article is error free can be expressed as  $\frac{p}{q}$  where  $p$  and  $q$  are relatively prime positive integers. Let  $\lambda = q - p$ , then find the sum of the digits of  $\lambda$ .
- India and Australia play a series of 7 one-day matches. Each team has equal probability of winning a match. No match ends in a draw. If the probability that India wins atleast three consecutive matches can be expressed as  $\frac{p}{q}$  where  $p$  and  $q$  are relatively prime positive integers. Find the unit digit of  $p$ .
- Two hunters A and B set out to hunt ducks. Each of them hits as often as he misses when shooting at ducks. Hunter A shoots at 50 ducks and hunter B shoots at 51 ducks. The probability that B bags more ducks than A can be expressed as  $\frac{p}{q}$  in its lowest form. Find the value of  $(p + q)$ .
- If  $a, b, c \in N$ , the probability that  $a^2 + b^2 + c^2$  is divisible by 7 is  $\frac{m}{n}$  where  $m, n$  are relatively prime natural numbers, then  $m + n$  is equal to :
- A fair coin is tossed 10 times. If the probability that heads never occur on consecutive tosses be  $\frac{m}{n}$  (where  $m, n$  are coprime and  $m, n \in N$ ), then the value of  $(n - 7m)$  equals to :
- A bag contains 2 red, 3 green and 4 black balls. 3 balls are drawn randomly and exactly 2 of them are found to be red. If  $p$  denotes the chance that one of the three balls drawn is green ; find the value of  $7p$ .
- There are 3 different pairs (i.e., 6 units say  $a, a, b, b, c, c$ ) of shoes in a lot. Now three person come and pick the shoes randomly (each gets 2 units). Let  $p$  be the probability that no one is able to wear shoes (i.e., no one gets a correct pair), then the value of  $\frac{13p}{4-p}$ , is :
- A fair coin is tossed 12 times. If the probability that two heads do not occur consecutively is  $p$ , then the value of  $\frac{[\sqrt{4096p-1}]}{2}$  is, where  $[ ]$  denotes greatest integer function :
- $X$  and  $Y$  are two weak students in mathematics and their chances of solving a problem correctly are  $\frac{1}{8}$  and  $\frac{1}{12}$  respectively. They are given a question and they obtain the same answer. If the probability of common mistake is  $\frac{1}{1001}$ , then probability that the answer was correct is  $a/b$  ( $a$  and  $b$  are coprimes). Then  $|a - b| =$



- 10.** Seven digit numbers are formed using digits 1, 2, 3, 4, 5, 6, 7, 8, 9 without repetition. The probability of selecting a number such that product of any 5 consecutive digits is divisible by either 5 or 7 is  $P$ . Then  $12P$  is equal to
- 11.** Assume that for every person the probability that he has exactly one child, exactly 2 children and exactly 3 children are  $\frac{1}{4}$ ,  $\frac{1}{2}$  and  $\frac{1}{4}$  respectively. The probability that a person will have 4 grand children can be expressed as  $\frac{p}{q}$  where  $p$  and  $q$  are relatively prime positive integers. Find the value of  $5p - q$ .
- 12.** Mr. B has two fair 6-sided dice, one whose faces are numbered 1 to 6 and the second whose faces are numbered 3 to 8. Twice, he randomly picks one of dice (each dice equally likely) and rolls it. Given the sum of the resulting two rolls is 9. The probability he rolled same dice twice is  $\frac{m}{n}$  where  $m$  and  $n$  are relatively prime positive integers. Find  $(m + n)$ .

### Answers

1.	7	2.	7	3.	3	4.	8	5.	1	6.	3	7.	2
8.	9	9.	1	10.	7	11.	7	12.	7				

□□□

### Exercise-1 : Single Choice Problems

- Solution set of the inequality  $\log_{10^2} x - 3(\log_{10} x)(\log_{10}(x-2)) + 2\log_{10^2}(x-2) < 0$ , is :  
 (a) (0, 4)                      (b)  $(-\infty, 1)$                       (c) (4,  $\infty$ )                      (d) (2, 4)
- The number of real solution/s of the equation  $9^{\log_3(\log_e x)} = \log_e x - (\log_e x)^2 + 1$  is :  
 (a) 0                      (b) 1                      (c) 2                      (d) 3
- If  $a, b, c$  are positive numbers such that  $a^{\log_3 7} = 27, b^{\log_7 11} = 49, c^{\log_{11} 25} = \sqrt{11}$ , then the sum of digits of  $S = a^{(\log_3 7)^2} + b^{(\log_7 11)^2} + c^{(\log_{11} 25)^2}$  is :  
 (a) 15                      (b) 17                      (c) 19                      (d) 21
- Least positive integral value of 'a' for which  $\log\left(\frac{x+1}{x}\right)(a^2 - 3a + 3) > 0; (x > 0)$  :  
 (a) 1                      (b) 2                      (c) 3                      (d) 4
- Let  $P = \frac{5}{\frac{1}{\log_2 x} + \frac{1}{\log_3 x} + \frac{1}{\log_4 x} + \frac{1}{\log_5 x}}$  and  $(120)^P = 32$ , then the value of  $x$  be :  
 (a) 1                      (b) 2                      (c) 3                      (d) 4
- If  $x, y, z$  be positive real numbers such that  $\log_{2x}(z) = 3, \log_{5y}(z) = 6$  and  $\log_{xy}(z) = \frac{2}{3}$  then the value of  $z$  is :  
 (a)  $\frac{1}{5}$                       (b)  $\frac{1}{10}$                       (c)  $\frac{3}{5}$                       (d)  $\frac{4}{9}$
- Sum of values of  $x$  and  $y$  satisfying  $\log_x(\log_3(\log_x y)) = 0$  and  $\log_y 27 = 1$  is :  
 (a) 27                      (b) 30                      (c) 33                      (d) 36
- $\log_{0.01} 1000 + \log_{0.1} 0.0001$  is equal to :  
 (a) -2                      (b) 3                      (c) -5/2                      (d) 5/2

9. If  $\log_{12} 27 = a$ , then  $\log_6 16 =$   
 (a)  $2\left(\frac{3-a}{3+a}\right)$  (b)  $3\left(\frac{3-a}{3+a}\right)$  (c)  $4\left(\frac{3-a}{3+a}\right)$  (d) None of these
10. If  $\log_2(\log_2(\log_3 x)) = \log_2(\log_3(\log_2 y)) = 0$  then the value of  $(x + y)$  is :  
 (a) 17 (b) 9 (c) 21 (d) 19
11. Suppose that  $a$  and  $b$  are positive real numbers such that  $\log_{27} a + \log_9 b = \frac{7}{2}$  and  $\log_{27} b + \log_9 a = \frac{2}{3}$ . Then the value of  $a \cdot b$  is :  
 (a) 81 (b) 243 (c) 27 (d) 729
12. If  $2^a = 5$ ,  $5^b = 8$ ,  $8^c = 11$  and  $11^d = 14$ , then the value of  $2^{abcd}$  is :  
 (a) 1 (b) 2 (c) 7 (d) 14
13. Which of the following conditions necessarily imply that the real number  $x$  is rational ?  
 (I)  $x^2$  is rational (II)  $x^3$  and  $x^5$  are rational (III)  $x^2$  and  $x^3$  are rational  
 (a) I and II only (b) I and III only (c) II and III only (d) III only
14. The value of  $\frac{\log_8 17}{\log_9 23} - \frac{\log_{2\sqrt{2}} 17}{\log_3 23}$  is equal to :  
 (a) -1 (b) 0 (c)  $\frac{\log_2 17}{\log_3 23}$  (d)  $\frac{4(\log_2 17)}{3(\log_3 23)}$
15. The true solution set of inequality  $\log_{(2x-3)}(3x-4) > 0$  is equal to :  
 (a)  $\left(\frac{4}{3}, \frac{5}{3}\right) \cup (2, \infty)$  (b)  $\left(\frac{3}{2}, \frac{5}{3}\right) \cup (2, \infty)$  (c)  $\left(\frac{4}{3}, \frac{3}{2}\right) \cup (2, \infty)$  (d)  $\left(\frac{2}{3}, \frac{4}{3}\right) \cup (2, \infty)$
16. If  $P$  is the number of natural numbers whose logarithm to the base 10 have the characteristic  $p$  and  $Q$  is the number of natural numbers logarithm of whose reciprocals to the base 10 have the characteristic  $-q$  then  $\log_{10} P - \log_{10} Q$  has the value equal to :  
 (a)  $p - q + 1$  (b)  $p - q$  (c)  $p + q - 1$  (d)  $p - q - 1$
17. If  $2^{2010} = a_n 10^n + a_{n-1} 10^{n-1} + \dots + a_2 10^2 + a_1 \cdot 10 + a_0$ , where  $a_i \in \{0, 1, 2, \dots, 9\}$  for all  $i = 0, 1, 2, 3, \dots, n$ , then  $n =$   
 (a) 603 (b) 604 (c) 605 (d) 606
18. The number of zeros after decimal before the start of any significant digit in the number  $N = (0.15)^{20}$  are :  
 (a) 15 (b) 16 (c) 17 (d) 18
19.  $\log_2[\log_4(\log_{10} 16^4 + \log_{10} 25^8)]$  simplifies to :  
 (a) an irrational (b) an odd prime  
 (c) a composite (d) unity
20. The sum of all the solutions to the equation  $2 \log x - \log(2x - 75) = 2$  :  
 (a) 30 (b) 350 (c) 75 (d) 200

21.  $x^{\log_x a \cdot \log_a y \cdot \log_y z}$  is equal to :  
 (a)  $x$  (b)  $y$  (c)  $z$  (d)  $x^z$
22. Number of solution(s) of the equation  $x^{x\sqrt{x}} = (x\sqrt{x})^x$  is/are :  
 (a) 0 (b) 1 (c) 2 (d) 3
23. The difference of roots of the equation  $(\log_{27} x^3)^2 = \log_{27} x^6$  is :  
 (a)  $\frac{2}{3}$  (b) 1 (c) 9 (d) 8
24. If  $\log_{10} x + \log_{10} y = 2$ ,  $x - y = 15$  then :  
 (a)  $(x, y)$  lies on the line  $y = 4x + 3$  (b)  $(x, y)$  lies on  $y^2 = 4x$   
 (c)  $(x, y)$  lies on  $x = 4y$  (d)  $(x, y)$  lies on  $4x = y$
25. Product of all values of  $x$  satisfying the equation  
 $\sqrt{2^x} \sqrt[3]{4^x} (0.125)^{1/x} = 4(\sqrt[3]{2})$  is :  
 (a)  $\frac{14}{5}$  (b) 3 (c)  $-\frac{1}{5}$  (d)  $-\frac{3}{5}$
26. Sum of all values of  $x$  satisfying the equation  
 $25^{(2x-x^2+1)} + 9^{(2x-x^2+1)} = 34(15^{(2x-x^2)})$  is :  
 (a) 1 (b) 2 (c) 3 (d) 4
27. If  $a^x = b^y = c^z = d^w$ , then  $\log_a(bcd) =$   
 (a)  $z\left(\frac{1}{x} + \frac{1}{y} + \frac{1}{w}\right)$  (b)  $y\left(\frac{1}{x} + \frac{1}{z} + \frac{1}{w}\right)$  (c)  $x\left(\frac{1}{y} + \frac{1}{z} + \frac{1}{w}\right)$  (d)  $\frac{xyz}{w}$
28. If  $x = \frac{4}{(\sqrt{5}+1)(\sqrt[4]{5}+1)(\sqrt[8]{5}+1)(\sqrt[16]{5}+1)}$ . Then the value of  $(1+x)^{48}$  is :  
 (a) 5 (b) 25 (c) 125 (d) 625
29. If  $\log_x \log_{18}(\sqrt{2} + \sqrt{8}) = \frac{1}{3}$ , then the value of  $32x =$   
 (a) 2 (b) 4 (c) 6 (d) 8
30. Let  $n \in N$ ,  $f(n) = \begin{cases} \log_8 n & \text{if } \log_8 n \text{ is integer} \\ 0 & \text{otherwise} \end{cases}$ , then the value of  $\sum_{n=1}^{2011} f(n)$  is :  
 (a) 2011 (b)  $2011 \times 1006$  (c) 6 (d)  $2^{2011}$
31. If the equation  $\frac{\log_{12}(\log_8(\log_4 x))}{\log_5(\log_4(\log_y(\log_2 x)))} = 0$  has a solution for 'x' when  $c < y < b$ ,  $y \neq a$ , where 'b' is as large as possible and 'c' is as small as possible, then the value of  $(a + b + c)$  is equals to :  
 (a) 18 (b) 19 (c) 20 (d) 21

32. If  $\log_{0.3}(x-1) < \log_{0.09}(x-1)$ , then  $x$  lies in the interval :  
 (a)  $(2, \infty)$  (b)  $(1, 2)$  (c)  $(-2, -1)$  (d)  $\left(1, \frac{3}{2}\right)$
33. The absolute integral value of the solution of the equation  $\sqrt{7^{2x^2-5x-6}} = (\sqrt{2})^{3\log_2 49}$   
 (a) 2 (b) 1 (c) 4 (d) 5
34. Let  $1 \leq x \leq 256$  and  $M$  be the maximum value of  $(\log_2 x)^4 + 16(\log_2 x)^2 \log_2 \left(\frac{16}{x}\right)$ . The sum of the digits of  $M$  is :  
 (a) 9 (b) 11 (c) 13 (d) 15
35. Let  $1 \leq x \leq 256$  and  $M$  be the maximum value of  $(\log_2 x)^4 + 16(\log_2 x)^2 \log_2 \left(\frac{16}{x}\right)$ . The sum of the digits of  $M$  is :  
 (a) 9 (b) 11 (c) 13 (d) 15
36. Number of real solution(s) of the equation  $9\log_3(\log nx) = \ln x - (\ln^2 x) + 1$  is :  
 (a) 0 (b) 1 (c) 2 (d) 3
37. The number of real values of the parameter  $\lambda$  for which  $(\log_{16} x)^2 - \log_{16} x + \log_{16} \lambda = 0$  with real coefficients will have exactly one solution is :  
 (a) 1 (b) 2 (c) 3 (d) 4
38. A rational number which is 50 times its own logarithm to the base 10 is :  
 (a) 1 (b) 10 (c) 100 (d) 1000
39. If  $x = \log_5(1000)$  and  $y = \log_7(2058)$ , then  
 (a)  $x > y$  (b)  $x < y$  (c)  $x = y$  (d) none of these
40.  $7 \log \left(\frac{16}{15}\right) + 5 \log \left(\frac{25}{24}\right) + 3 \log \left(\frac{81}{80}\right)$  is equal to :  
 (a) 0 (b) 1 (c)  $\log 2$  (d)  $\log 3$
41.  $\log_{10} \tan 1^\circ + \log_{10} \tan 2^\circ + \dots + \log_{10} \tan 89^\circ$  is equal to :  
 (a) 0 (b) 1 (c) 27 (d) 81
42.  $\log_7 \log_7 \sqrt{7\sqrt{7\sqrt{7}}}$  is equal to :  
 (a)  $3 \log_2 7$  (b)  $3 \log_7 2$  (c)  $1 - 3 \log_7 2$  (d)  $1 - 3 \log_2 7$
43. If  $(4)^{\log_9 3} + (9)^{\log_2 4} = (10)^{\log_x 83}$ , then  $x$  is equal to :  
 (a) 2 (b) 3 (c) 10 (d) 30
44.  $x^{\log_{10} \left(\frac{y}{z}\right)} \cdot y^{\log_{10} \left(\frac{z}{x}\right)} \cdot z^{\log_{10} \left(\frac{x}{y}\right)}$  is equal to :  
 (a) 0 (b) 1 (c) -1 (d) 2

45. The solution set of the equation :  $\log_x 2 \log_{2x} 2 = \log_{4x} 2$  is :  
 (a)  $\{2^{-\sqrt{2}}, 2^{\sqrt{2}}\}$  (b)  $\{1/2, 2\}$  (c)  $\{1/4, 2^2\}$  (d) none of these
46. The least value of the expression  $2 \log_{10} x - \log_x 0.01$  is ( $x > 1$ )  
 (a) 2 (b) 4 (c) 6 (d) 8
47. If  $\sqrt{\log_2 x} - 0.5 = \log_2 \sqrt{x}$ , then  $x$  equals to :  
 (a) odd integer (b) prime number  
 (c) composite number (d) irrational
48. If  $x_1$  and  $x_2$  are the roots of the equation  $e^{2x} x^{\ln x} = x^3$  with  $x_1 > x_2$ , then  
 (a)  $x_1 = 2x_2$  (b)  $x_1 = x_2^2$  (c)  $2x_1 = x_2^2$  (d)  $x_1^2 = x_2^2$
49. Let  $M$  denote  $\text{antilog}_{32} 0.6$  and  $N$  denote the value of  $49^{(1-\log_7 2)} + 5^{-\log_5 4}$ . Then  $M.N$  is :  
 (a) 100 (b) 400 (c) 50 (d) 200
50. If  $\log_2(\log_2(\log_3 x)) = \log_3(\log_3(\log_2 y)) = 0$ , then  $x - y$  is equal to :  
 (a) 0 (b) 1 (c) 8 (d) 9
51.  $\left| \log_{\frac{1}{2}} 10 + \left| \log_4 625 - \left| \log_{\frac{1}{2}} 5 \right| \right| \right| =$   
 (a)  $\log_{1/2} 2$  (b)  $\log_2 5$  (c)  $\log_2 2$  (d)  $\log_2 25$
52. If  $\log_4 5 = a$  and  $\log_5 6 = b$ , then  $\log_3 2$  is equal to :  
 (a)  $\frac{1}{2a+1}$  (b)  $\frac{1}{2b+1}$  (c)  $2ab+1$  (d)  $\frac{1}{2ab-1}$
53. If  $x = \log_a bc$ ;  $y = \log_b ac$  and  $z = \log_c ab$  then which of the following is equal to unity ?  
 (a)  $x+y+z$  (b)  $xyz$   
 (c)  $\frac{1}{1+x} + \frac{1}{1+y} + \frac{1}{1+z}$  (d)  $(1+x) + (1+y) + (1+z)$
54.  $x^{\log_x a \cdot \log_a y \cdot \log_y z}$  is equal to :  
 (a)  $x$  (b)  $y$  (c)  $z$  (d)  $a$
55. Number of value(s) of 'x' satisfying the equation  $x^{\log_{\sqrt{x}}(x-3)} = 9$  is/are  
 (a) 0 (b) 1 (c) 2 (d) 6
56.  $\log_{0.01} 1000 + \log_{0.1} 0.0001$  is equal to :  
 (a) -2 (b) 3 (c)  $-\frac{5}{2}$  (d)  $\frac{5}{2}$
57. If  $7 \log_a \frac{16}{15} + 5 \log_a \frac{25}{24} + 3 \log_a \frac{81}{80} = 8$ , then  $a =$   
 (a)  $2^{1/8}$  (b)  $(10)^{1/8}$  (c)  $(30)^{1/8}$  (d) 1

58.  $\log_8(128) - \log_9 \cot\left(\frac{\pi}{3}\right) =$
- (a)  $\frac{31}{12}$  (b)  $\frac{19}{12}$  (c)  $\frac{13}{12}$  (d)  $\frac{11}{12}$
59. The value of  $\left(\frac{1}{\sqrt{27}}\right)^2 \left(\frac{\log_5 16}{2 \log_5 9}\right)$  equals to :
- (a)  $\frac{5\sqrt{2}}{27}$  (b)  $\frac{\sqrt{2}}{27}$  (c)  $\frac{4\sqrt{2}}{27}$  (d)  $\frac{2\sqrt{2}}{27}$
60. The sum of all the roots of the equation  $\log_2(x-1) + \log_2(x+2) - \log_2(3x-1) = \log_2 4$
- (a) 12 (b) 2 (c) 10 (d) 11
61.  $\frac{(\log_{100} 10)(\log_2(\log_4 2))(\log_4 \log_2^2(256)^2)}{\log_4 8 + \log_8 4} =$
- (a)  $-\frac{6}{13}$  (b)  $-\frac{1}{2}$  (c)  $-\frac{8}{13}$  (d)  $-\frac{12}{13}$
62. Let  $\lambda = \log_5 \log_5(3)$ . If  $3^{k+5^{-\lambda}} = 405$ , then the value of  $k$  is :
- (a) 3 (b) 5 (c) 4 (d) 6
63. A circle has a radius  $\log_{10}(a^2)$  and a circumference of  $\log_{10}(b^4)$ . Then the value of  $\log_a b$  is equal to :
- (a)  $\frac{1}{4\pi}$  (b)  $\frac{1}{\pi}$  (c)  $2\pi$  (d)  $\pi$
64. If  $2^x = 3^y = 6^{-z}$ , the value of  $\frac{1}{x} + \frac{1}{y} + \frac{1}{z}$  is equal to :
- (a) 0 (b) 1 (c) 2 (d) 3
65. The value of  $\log_{(\sqrt{2}-1)}(5\sqrt{2}-7)$  is :
- (a) 0 (b) 1 (c) 2 (d) 3
66. The value of  $\log_{ab} \left(\frac{\sqrt[3]{a}}{\sqrt{b}}\right)$ , if  $\log_{ab} a = 4$  is equal to :
- (a) 2 (b)  $\frac{13}{6}$  (c)  $\frac{15}{6}$  (d)  $\frac{17}{6}$
67. Identify the correct option
- (a)  $\log_2 3 < \log_{1/4} 5$  (b)  $\log_5 7 < \log_8 3$
- (c)  $\log_{\sqrt[3]{2}} \sqrt{3} > \log_{\sqrt[3]{2}} \sqrt{5}$  (d)  $2^4 > \left(\frac{3}{2}\right)^{1/3}$
68. Sum of all values of  $x$  satisfying the system of equations  $5(\log_y x + \log_x y) = 26$ ,  $xy = 64$  is :
- (a) 42 (b) 34 (c) 32 (d) 2

69. The product of all values of  $x$  satisfying the equations  $\log_3 a - \log_x a = \log_{x/3} a$  is :
- (a) 3                      (b)  $\frac{3}{2}$                       (c) 18                      (d) 27
70. The value of  $x + y + z$  satisfying the system of equations
- $$\begin{aligned} \log_2 x + \log_4 y + \log_4 z &= 2 \\ \log_3 y + \log_9 z + \log_9 x &= 2 \\ \log_4 z + \log_{16} x + \log_{16} y &= 2 \end{aligned}$$
- (a)  $\frac{175}{12}$                       (b)  $\frac{349}{24}$                       (c)  $\frac{353}{24}$                       (d)  $\frac{112}{3}$
71.  $\left(\frac{1}{49}\right)^{1+\log_7 2} + 5^{-\log_1 7} =$
- (a)  $7\frac{1}{196}$                       (b)  $7\frac{3}{196}$                       (c)  $7\frac{5}{196}$                       (d)  $7\frac{1}{98}$
72. The number of real values of  $x$  satisfying the equation  $\log_2(3-x) - \log_2 \left( \frac{\sin\left(\frac{3\pi}{4}\right)}{(5-x)} \right) = \frac{1}{2} + \log_2(x+7)$  is :
- (a) 0                      (b) 1                      (c) 2                      (d) 3
73. If  $\log_k x \log_5 k = \log_x 5$ ,  $k \neq 1$ ,  $k > 0$ , then sum of all values of  $x$  is :
- (a) 5                      (b)  $\frac{24}{5}$                       (c)  $\frac{26}{5}$                       (d)  $\frac{37}{5}$
74. The product of all values of  $x$  satisfying the equation  $|x-1|^{\log_3 x^2 - 2\log_x 9} = (x-1)^7$ , is :
- (a) 162                      (b)  $\frac{162}{\sqrt{3}}$                       (c)  $\frac{81}{\sqrt{3}}$                       (d) 81
75. The number of values of  $x$  satisfying the equation  $\log_2(9^{x-1} + 7) = 2 + \log_2(3^{x-1} + 1)$  is :
- (a) 1                      (b) 2                      (c) 3                      (d) 0
76. Which is the correct order for a given number  $\alpha$ ,  $\alpha > 1$
- (a)  $\log_2 \alpha < \log_3 \alpha < \log_e \alpha < \log_{10} \alpha$                       (b)  $\log_{10} \alpha < \log_3 \alpha < \log_e \alpha < \log_2 \alpha$   
 (c)  $\log_{10} \alpha < \log_e \alpha < \log_2 \alpha < \log_3 \alpha$                       (d)  $\log_3 \alpha < \log_e \alpha < \log_2 \alpha < \log_{10} \alpha$
77. Let  $1 \leq x \leq 256$  and  $M$  be the maximum value of  $(\log_2 x)^4 + 16(\log_2 x)^2 \log_2 \left(\frac{16}{x}\right)$ . The sum of the digits of  $M$  is :
- (a) 9                      (b) 11                      (c) 13                      (d) 15





**Exercise-2 : One or More than One Answer is/are Correct**

1. The values of 'x' satisfies the equation  $\frac{1 - 2(\log x^2)^2}{\log x - 2(\log x)^2} = 1$  (is/are) :

(where log is logarithm to the base 10)

- (a)  $\frac{1}{\sqrt{10}}$       (b)  $\frac{1}{\sqrt{20}}$       (c)  $\sqrt[3]{10}$       (d)  $\sqrt{10}$

2. If  $\log_a x = b$  for permissible values of  $a$  and  $x$  then identify the statement(s) which can be correct?

- (a) If  $a$  and  $b$  are two irrational numbers then  $x$  can be rational.  
 (b) If  $a$  rational and  $b$  irrational then  $x$  can be rational.  
 (c) If  $a$  irrational and  $b$  rational then  $x$  can be rational.  
 (d) If  $a$  rational and  $b$  rational then  $x$  can be rational.

3. Consider the quadratic equation,  $(\log_{10} 8)x^2 - (\log_{10} 5)x = 2(\log_2 10)^{-1} - x$ . Which of the following quantities are irrational ?

- (a) Sum of the roots      (b) Product of the roots  
 (c) Sum of the coefficients      (d) Discriminant

4. Let  $A = \text{Minimum}(x^2 - 2x + 7)$ ,  $x \in R$  and  $B = \text{Minimum}(x^2 - 2x + 7)$ ,  $x \in [2, \infty)$ , then :

- (a)  $\log_{(B-A)}(A+B)$  is not defined      (b)  $A+B=13$   
 (c)  $\log_{(2B-A)} A < 1$       (d)  $\log_{(2A-B)} A > 1$

**Answers**

1.	(a, c)	2.	(a, b, c, d)	3.	(c, d)	4.	(a, b, c, d)				
----	--------	----	--------------	----	--------	----	--------------	--	--	--	--

**Exercise-3 : Comprehension Type Problems**

**Paragraph for Question Nos. 1 to 3**

Let  $\log_3 N = \alpha_1 + \beta_1$

$\log_5 N = \alpha_2 + \beta_2$

$\log_7 N = \alpha_3 + \beta_3$

where  $\alpha_1, \alpha_2$  and  $\alpha_3$  are integers and  $\beta_1, \beta_2, \beta_3 \in [0, 1)$ .

1. Number of integral values of  $N$  if  $\alpha_1 = 4$  and  $\alpha_2 = 2$ :  
 (a) 46                      (b) 45                      (c) 44                      (d) 47
2. Largest integral value of  $N$  if  $\alpha_1 = 5, \alpha_2 = 3$  and  $\alpha_3 = 2$ .  
 (a) 342                      (b) 343                      (c) 243                      (d) 242
3. Difference of largest and smallest integral values of  $N$  if  $\alpha_1 = 5, \alpha_2 = 3$  and  $\alpha_3 = 2$ .  
 (a) 97                      (b) 100                      (c) 98                      (d) 99

**Paragraph for Question Nos. 4 to 5**

If  $\log_{10}|x^3 + y^3| - \log_{10}|x^2 - xy + y^2| + \log_{10}|x^3 - y^3| - \log_{10}|x^2 + xy + y^2| = \log_{10} 221$ .

Where  $x, y$  are integers, then

4. If  $x = 111$ , then  $y$  can be :  
 (a)  $\pm 111$                       (b)  $\pm 2$                       (c)  $\pm 110$                       (d)  $\pm 109$
5. If  $y = 2$ , then value of  $x$  can be :  
 (a)  $\pm 111$                       (b)  $\pm 15$                       (c)  $\pm 2$                       (d)  $\pm 110$


**Paragraph for Question Nos. 6 to 7**

Given a right triangle  $ABC$  right angled at  $C$  and whose legs are given  $1 + 4\log_{p^2}(2p)$ ,  $1 + 2^{\log_2(\log_2 p)}$  and hypotenuse is given to be  $1 + \log_2(4p)$ . The area of  $\Delta ABC$  and circle circumscribing it are  $\Delta_1$  and  $\Delta_2$  respectively, then

6.  $\Delta_1 + \frac{4\Delta_2}{\pi}$  is equal to :  
 (a) 31                      (b) 28                      (c)  $3 + \frac{1}{\sqrt{2}}$                       (d) 199
7. The value of  $\sin\left(\frac{\pi(25p^2\Delta_1 + 2)}{6}\right) =$   
 (a)  $\frac{1}{2}$                       (b)  $\frac{1}{\sqrt{2}}$                       (c)  $\frac{\sqrt{3}}{2}$                       (d) 1

**Answers**

1.	(c)	2.	(a)	3.	(d)	4.	(c)	5.	(b)	6.	(a)	7.	(c)						
----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	--	--	--	--	--	--


**Exercise-4 : Matching Type Problems**

1.

	Column-I		Column-II
(A)	If $a = 3(\sqrt{8+2\sqrt{7}} - \sqrt{8-2\sqrt{7}})$ , $b = \sqrt{(42)(30) + 36}$ , then the value of $\log_a b$ is equal to	(P)	-1
(B)	If $a = (\sqrt{4+2\sqrt{3}} - \sqrt{4-2\sqrt{3}})$ , $b = \sqrt{11+6\sqrt{2}} - \sqrt{11-6\sqrt{2}}$ then the value of $\log_a b$ is equal to	(Q)	1
(C)	If $a = \sqrt{3+2\sqrt{2}}$ , $b = \sqrt{3-2\sqrt{2}}$ , then the value of $\log_a b$ is equal to	(R)	2
(D)	If $a = \sqrt{7+\sqrt{7^2-1}}$ , $b = \sqrt{7-\sqrt{7^2-1}}$ , then the value of $\log_a b$ is equal to	(S)	$\frac{3}{2}$
		(T)	None of these

**Answers**
**1.** A  $\rightarrow$  R ; B  $\rightarrow$  S ; C  $\rightarrow$  P ; D  $\rightarrow$  P

### Exercise-5 : Subjective Type Problems

- The number  $N = 6^{\log_{10} 40} \cdot 5^{\log_{10} 36}$  is a natural number. Then sum of digits of  $N$  is :
- The minimum value of 'c' such that  $\log_b(a^{\log_2 b}) = \log_a(b^{\log_2 b})$  and  $\log_a(c - (b - a)^2) = 3$ , where  $a, b \in N$  is :
- How many positive integers  $b$  have the property that  $\log_b 729$  is a positive integer ?
- The number of negative integral values of  $x$  satisfying the inequality  $\log_{\left(\frac{x+5}{2}\right)} \left(\frac{x-5}{2x-3}\right)^2 < 0$  is :
- $\frac{6}{5} a^{(\log_a x)(\log_{10} a)(\log_a 5)} - 3^{\log_{10} \left(\frac{x}{10}\right)} = 9^{\log_{100} x + \log_4 2}$  (where  $a > 0, a \neq 1$ ), then  $\log_3 x = \alpha + \beta$ ,  $\alpha$  is integer,  $\beta \in [0, 1)$ , then  $\alpha =$
- If  $\log_5 \left(\frac{a+b}{3}\right) = \frac{\log_5 a + \log_5 b}{2}$ , then  $\frac{a^4 + b^4}{a^2 b^2} =$
- Let  $a, b, c, d$  are positive integers such that  $\log_a b = \frac{3}{2}$  and  $\log_c d = \frac{5}{4}$ . If  $(a - c) = 9$ . Find the value of  $(b - d)$ .
- The number of real values of  $x$  satisfying the equation  $\log_{10} \sqrt{1+x} + 3 \log_{10} \sqrt{1-x} = 2 + \log_{10} \sqrt{1-x^2}$  is :
- The ordered pair  $(x, y)$  satisfying the equation  $x^2 = 1 + 6 \log_4 y$  and  $y^2 = 2^x y + 2^{2x+1}$  are  $(x_1, y_1)$  and  $(x_2, y_2)$ , then find the value of  $\log_2 |x_1 x_2 y_1 y_2|$ .
- If  $\log_7 \log_7 \sqrt{7\sqrt{7\sqrt{7}}} = 1 - a \log_7 2$  and  $\log_{15} \log_{15} \sqrt{15\sqrt{15\sqrt{15\sqrt{15}}}} = 1 - b \log_{15} 2$ , then  $a + b =$
- The number of ordered pair(s) of  $(x, y)$  satisfying the equations  $\log_{(1+x)}(1 - 2y + y^2) + \log_{(1-y)}(1 + 2x + x^2) = 4$  and  $\log_{(1+x)}(1 + 2y) + \log_{(1-y)}(1 + 2x) = 2$
- If  $\log_b n = 2$  and  $\log_n(2b) = 2$ , then  $nb =$
- If  $\log_y x + \log_x y = 2$ , and  $x^2 + y = 12$ , then the value of  $xy$  is :
- If  $x, y$  satisfy the equation,  $y^x = x^y$  and  $x = 2y$ , then  $x^2 + y^2 =$
- Find the number of real values of  $x$  satisfying the equation. 
$$\log_2(4^{x+1} + 4) \cdot \log_2(4^x + 1) = \log_{1/\sqrt{2}} \sqrt{\frac{1}{8}}$$
- If  $x_1, x_2 (x_1 > x_2)$  are the two solutions of the equation  $3^{\log_2 x} - 12(x^{\log_{16} 9}) = \log_3 \left(\frac{1}{3}\right)^{3^3}$ , then the value of  $x_1 - 2x_2$  is :

17. Find the number of real values of  $x$  satisfying the equation  $9^{2\log_9 x} + 4x + 3 = 0$ .

18. If  $\log_{16}(\log_{\sqrt[3]{3}}(\log_{\sqrt[5]{5}}(x))) = \frac{1}{2}$ ; find  $x$ .

19. The value  $\left[ \frac{1}{6} \left( \frac{2\log_{10}(1728)}{1 + \frac{1}{2}\log_{10}(0.36) + \frac{1}{3}\log_{10} 8} \right)^{1/2} \right]^{-1}$  is :

### Answers

1.	9	2.	8	3.	4	4.	0	5.	4	6.	47	7.	93
8.	0	9.	7	10.	7	11.	1	12.	2	13.	9	14.	20
15.	1	16.	8	17.	0	18.	5	19.	2				

# Co-ordinate Geometry

---

**17. Straight Lines**

**18. Circle**

**19. Parabola**

**20. Ellipse**

**21. Hyperbola**

Chapter 17 – Straight Lines

### Exercise-1 : Single Choice Problems

- The ratio in which the line segment joining  $(2, -3)$  and  $(5, 6)$  is divided by the  $x$ -axis is :
  - $3 : 1$
  - $1 : 2$
  - $\sqrt{3} : 2$
  - $\sqrt{2} : 3$
- If  $L$  is the line whose equation is  $ax + by = c$ . Let  $M$  be the reflection of  $L$  through the  $y$ -axis, and let  $N$  be the reflection of  $L$  through the  $x$ -axis. Which of the following must be true about  $M$  and  $N$  for all choices of  $a, b$  and  $c$  ?
  - The  $x$ -intercepts of  $M$  and  $N$  are equal
  - The  $y$ -intercepts of  $M$  and  $N$  are equal
  - The slopes of  $M$  and  $N$  are equal
  - The slopes of  $M$  and  $N$  are reciprocal
- The complete set of real values of ' $a$ ' such that the point  $P(a, \sin a)$  lies inside the triangle formed by the lines  $x - 2y + 2 = 0$ ;  $x + y = 0$  and  $x - y - \pi = 0$ , is :
  - $\left(0, \frac{\pi}{6}\right) \cup \left(\frac{\pi}{3}, \frac{\pi}{2}\right)$
  - $\left(\frac{\pi}{2}, \pi\right) \cup \left(\frac{2\pi}{2}, 2\pi\right)$
  - $(0, \pi)$
  - $\left(\frac{\pi}{3}, \frac{\pi}{2}\right)$
- Let  $m$  be a positive integer and let the lines  $13x + 11y = 700$  and  $y = mx - 1$  intersect in a point whose coordinates are integer. Then  $m$  equals to :
  - 4
  - 5
  - 6
  - 7
- If  $P = \left(\frac{1}{x_p}, p\right)$ ;  $Q = \left(\frac{1}{x_q}, q\right)$ ;  $R = \left(\frac{1}{x_r}, r\right)$   
 where  $x_k \neq 0$ , denotes the  $k^{\text{th}}$  terms of a H.P for  $k \in N$ , then:




- (a) ar.  $(\Delta PQR) = \frac{p^2 q^2 r^2}{2} \sqrt{(p-q)^2 + (q-r)^2 + (r-p)^2}$
- (b)  $\Delta PQR$  is a right angled triangle  
 (c) the points  $P, Q, R$  are collinear  
 (d) None of these
6. If the sum of the slopes of the lines given by  $x^2 - 2cxy - 7y^2 = 0$  is four times their product, then  $c$  has the value :  
 (a) 1 (b) -1 (c) 2 (d) -2
7. A piece of cheese is located at  $(12, 10)$  in a coordinate plane. A mouse is at  $(4, -2)$  and is running up the line  $y = -5x + 18$ . At the point  $(a, b)$ , the mouse starts getting farther from the cheese rather than closer to it. The value of  $(a + b)$  is:  
 (a) 6 (b) 10  
 (c) 18 (d) 14
8. The vertex of right angle of a right angled triangle lies on the straight line  $2x + y - 10 = 0$  and the two other vertices, at points  $(2, -3)$  and  $(4, 1)$  then the area of triangle in sq. units is:  
 (a)  $\sqrt{10}$  (b) 3 (c)  $\frac{33}{5}$  (d) 11
9. Given the family of lines,  $a(2x + y + 4) + b(x - 2y - 3) = 0$ . Among the lines of the family, the number of lines situated at a distance of  $\sqrt{10}$  from the point  $M(2, -3)$  is:  
 (a) 0 (b) 1  
 (c) 2 (d)  $\infty$
10. Point  $(0, \beta)$  lies on or inside the triangle formed by the lines  $y = 0, x + y = 8$  and  $3x - 4y + 12 = 0$ . Then  $\beta$  can be :  
 (a) 2 (b) 4 (c) 8 (d) 12
11. If the lines  $x + y + 1 = 0, 4x + 3y + 4 = 0$  and  $x + \alpha y + \beta = 0$ , where  $\alpha^2 + \beta^2 = 2$ , are concurrent then:  
 (a)  $\alpha = 1, \beta = -1$  (b)  $\alpha = 1, \beta = \pm 1$   
 (c)  $\alpha = -1, \beta = \pm 1$  (d)  $\alpha = \pm 1, \beta = 1$
12. A straight line through the origin 'O' meets the parallel lines  $4x + 2y = 9$  and  $2x + y = -6$  at points  $P$  and  $Q$  respectively. Then the point 'O' divides the segment  $PQ$  in the ratio :  
 (a) 1 : 2 (b) 4 : 3 (c) 2 : 1 (d) 3 : 4
13. If the points  $(2a, a), (a, 2a)$  and  $(a, a)$  enclose a triangle of area 72 units, then co-ordinates of the centroid of the triangle may be :  
 (a)  $(4, 4)$  (b)  $(-4, 4)$  (c)  $(12, 12)$  (d)  $(16, 16)$
14. Let  $g(x) = ax + b$ , where  $a < 0$  and  $g$  is defined from  $[1, 3]$  onto  $[0, 2]$  then the value of  $\cot(\cos^{-1}(|\sin x| + |\cos x|) + \sin^{-1}(-|\cos x| - |\sin x|))$  is equal to :  
 (a)  $g(1)$  (b)  $g(2)$  (c)  $g(3)$  (d)  $g(1) + g(3)$

15. If the distances of any point  $P$  from the points  $A(a + b, a - b)$  and  $B(a - b, a + b)$  are equal, then locus of  $P$  is :
- (a)  $ax + by = 0$       (b)  $ax - by = 0$       (c)  $bx + ay = 0$       (d)  $x - y = 0$
16. If the equation  $4y^3 - 8a^2yx^2 - 3ay^2x + 8x^3 = 0$  represent three straight lines, two of them are perpendicular then sum of all possible values of  $a$  is equal to :
- (a)  $\frac{3}{8}$       (b)  $-\frac{3}{4}$       (c)  $\frac{1}{4}$       (d)  $-2$
17. The orthocentre of the triangle formed by the lines  $x - 7y + 6 = 0$ ,  $2x - 5y - 6 = 0$  and  $7x + y - 8 = 0$  is :
- (a)  $(8, 2)$       (b)  $(0, 0)$       (c)  $(1, 1)$       (d)  $(2, 8)$
18. All the chords of the curve  $2x^2 + 3y^2 - 5x = 0$  which subtend a right angle at the origin are concurrent at :
- (a)  $(0, 1)$       (b)  $(1, 0)$       (c)  $(-1, 1)$       (d)  $(1, -1)$
19. From a point  $P \equiv (3, 4)$  perpendiculars  $PQ$  and  $PR$  are drawn to line  $3x + 4y - 7 = 0$  and a variable line  $y - 1 = m(x - 7)$  respectively, then maximum area of  $\Delta PQR$  is :
- (a) 10      (b) 12      (c) 6      (d) 9
20. The equation of two adjacent sides of rhombus are given by  $y = x$  and  $y = 7x$ . The diagonals of the rhombus intersect each other at the point  $(1, 2)$ . Then the area of the rhombus is :
- (a)  $\frac{10}{3}$       (b)  $\frac{20}{3}$       (c)  $\frac{40}{3}$       (d)  $\frac{50}{3}$
21. The point  $P(3, 3)$  is reflected across the line  $y = -x$ . Then it is translated horizontally 3 units to the left and vertically 3 units up. Finally, it is reflected across the line  $y = x$ . What are the coordinates of the point after these transformations ?
- (a)  $(0, -6)$       (b)  $(0, 0)$   
(c)  $(-6, 6)$       (d)  $(-6, 0)$
22. The equations  $x = t^3 + 9$  and  $y = \frac{3t^3}{4} + 6$  represents a straight line where  $t$  is a parameter. Then  $y$ -intercept of the line is :
- (a)  $-\frac{3}{4}$       (b) 9      (c) 6      (d) 1
23. The combined equation of two adjacent sides of a rhombus formed in first quadrant is  $7x^2 - 8xy + y^2 = 0$ ; then slope of its longer diagonal is :
- (a)  $-\frac{1}{2}$       (b)  $-2$       (c) 2      (d)  $\frac{1}{2}$
24. The number of integral points inside the triangle made by the line  $3x + 4y - 12 = 0$  with the coordinate axes which are equidistant from at least two sides is/are :  
(an integral point is a point both of whose coordinates are integers.)
- (a) 1      (b) 2      (c) 3      (d) 4

25. The area of triangle formed by the straight lines whose equations are  $y = 4x + 2$ ,  $2y = x + 3$  and  $x = 0$  is :
- (a)  $\frac{25}{7\sqrt{2}}$                       (b)  $\frac{\sqrt{2}}{28}$                       (c)  $\frac{1}{28}$                       (d)  $\frac{15}{7}$
26. In a triangle  $ABC$ , if  $A$  is  $(1, 2)$  and the equations of the medians through  $B$  and  $C$  are  $x + y = 5$  and  $x = 4$  respectively then  $B$  must be :
- (a)  $(1, 4)$                       (b)  $(7, -2)$                       (c)  $(4, 1)$                       (d)  $(-2, 7)$
27. The equation of image of pair of lines  $y = |x - 1|$  with respect to  $y$ -axis is :
- (a)  $x^2 - y^2 - 2x + 1 = 0$                       (b)  $x^2 - y^2 - 4x + 4 = 0$   
(c)  $4x^2 - 4x - y^2 + 1 = 0$                       (d)  $x^2 - y^2 + 2x + 1 = 0$
28. If  $P, Q$  and  $R$  are three points with coordinates  $(1, 4)$ ,  $(4, 5)$  and  $(m, m)$  respectively, then the value of  $m$  for which  $PR + RQ$  is minimum, is :
- (a) 4                      (b) 3                      (c)  $\frac{17}{8}$                       (d)  $\frac{7}{2}$
29. The vertices of triangle  $ABC$  are  $A(-1, -7)$ ,  $B(5, 1)$  and  $C(1, 4)$ . The equation of the bisector of the angle  $ABC$  of  $\triangle ABC$  is :
- (a)  $y + 2x - 11 = 0$                       (b)  $x - 7y + 2 = 0$   
(c)  $y - 2x + 9 = 0$                       (d)  $y + 7x - 36 = 0$
30. If one of the lines given by  $6x^2 - xy + 4cy^2 = 0$  is  $3x + 4y = 0$ , then  $c =$
- (a) -3                      (b) -1                      (c) 3                      (d) 1
31. The equations of  $L_1$  and  $L_2$  are  $y = mx$  and  $y = nx$ , respectively. Suppose  $L_1$  make twice as large of an angle with the horizontal (measured counterclockwise from the positive  $x$ -axis) as does  $L_2$  and that  $L_1$  has 4 times the slope of  $L_2$ . If  $L_1$  is not horizontal, then the value of the product  $(mn)$  equals:
- (a)  $\frac{\sqrt{2}}{2}$                       (b)  $-\frac{\sqrt{2}}{2}$   
(c) 2                      (d) -2
32. Given  $A(0, 0)$  and  $B(x, y)$  with  $x \in (0, 1)$  and  $y > 0$ . Let the slope of the line  $AB$  equals  $m_1$ . Point  $C$  lies on the line  $x = 1$  such that the slope of  $BC$  equals  $m_2$  where  $0 < m_2 < m_1$ . If the area of the triangle  $ABC$  can be expressed as  $(m_1 - m_2)f(x)$ , then the largest possible value of  $f(x)$  is:
- (a) 1                      (b)  $1/2$   
(c)  $1/4$                       (d)  $1/8$
33. If non-zero numbers  $a, b, c$  are in H.P., then the straight line  $\frac{x}{a} + \frac{y}{b} + \frac{1}{c} = 0$  always passes through a fixed point, co-ordinate of fixed point is :
- (a)  $(-1, 2)$                       (b)  $(-1, -2)$                       (c)  $(1, -2)$                       (d)  $\left(1, \frac{1}{2}\right)$

34. If  $\frac{x^2}{a} + \frac{y^2}{b} + \frac{2xy}{h} = 0$  represent pair of straight lines and slope of one line is twice the other, then  $ab : h^2$  is :
- (a) 9 : 8                      (b) 8 : 9                      (c) 1 : 2                      (d) 2 : 1
35. **Statement-1:** A variable line drawn through a fixed point cuts the coordinate axes at  $A$  and  $B$ . The locus of mid-point of  $AB$  is a circle.  
**because**
- Statement-2:** Through 3 non-collinear points in a plane, only one circle can be drawn.
- (a) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
(b) Statement-1 is true, statement-2 is true and statement-2 is not the correct explanation for statement-1.  
(c) Statement-1 is true, statement-2 is false.  
(d) Statement-1 is false, statement-2 is true.
36. A line passing through origin and is perpendicular to two parallel lines  $2x + y + 6 = 0$  and  $4x + 2y - 9 = 0$ , then the ratio in which the origin divides this line segment is :
- (a) 1 : 2                      (b) 1 : 1  
(c) 5 : 4                      (d) 3 : 4
37. If a vertex of a triangle is  $(1, 1)$  and the mid-points of two sides through this vertex are  $(-1, 2)$  and  $(3, 2)$ , then the centroid of the triangle is :
- (a)  $\left(-1, \frac{7}{3}\right)$               (b)  $\left(-\frac{1}{3}, \frac{7}{3}\right)$               (c)  $\left(1, \frac{7}{3}\right)$               (d)  $\left(\frac{1}{3}, \frac{7}{3}\right)$
38. The diagonals of parallelogram  $PQRS$  are along the lines  $x + 3y = 4$  and  $6x - 2y = 7$ . Then  $PQRS$  must be :
- (a) rectangle                      (b) square  
(c) rhombus                      (d) neither rhombus nor rectangle
39. The two points on the line  $x + y = 4$  that lie at a unit perpendicular distance from the line  $4x + 3y = 10$  are  $(a_1, b_1)$  and  $(a_2, b_2)$ , then  $a_1 + b_1 + a_2 + b_2 =$
- (a) 5                      (b) 6                      (c) 7                      (d) 8
40. The orthocentre of the triangle formed by the lines  $x + y = 1$ ,  $2x + 3y = 6$  and  $4x - y + 4 = 0$  lies in :
- (a) first quadrant                      (b) second quadrant  
(c) third quadrant                      (d) fourth quadrant
41. The equation of the line passing through the intersection of the lines  $3x + 4y = -5$ ,  $4x + 6y = 6$  and perpendicular to  $7x - 5y + 3 = 0$  is :
- (a)  $5x + 7y - 2 = 0$                       (b)  $5x - 7y + 2 = 0$   
(c)  $7x - 5y + 2 = 0$                       (d)  $5x + 7y + 2 = 0$

42. The points  $(2, 1)$ ,  $(8, 5)$  and  $(x, 7)$  lie on a straight line. Then the value of  $x$  is :  
 (a) 10 (b) 11 (c) 12 (d)  $\frac{35}{3}$
43. In a parallelogram  $PQRS$  (taken in order),  $P$  is the point  $(-1, -1)$ ,  $Q$  is  $(8, 0)$  and  $R$  is  $(7, 5)$ . Then  $S$  is the point :  
 (a)  $(-1, 4)$  (b)  $(-2, 2)$  (c)  $(-2, \frac{7}{2})$  (d)  $(-2, 4)$
44. The area of triangle whose vertices are  $(a, a)$ ,  $(a + 1, a + 1)$ ,  $(a + 2, a)$  is :  
 (a)  $a^3$  (b)  $2a$  (c) 1 (d) 2
45. The equation  $x^2 + y^2 - 2xy - 1 = 0$  represents :  
 (a) two parallel straight lines (b) two perpendicular straight lines  
 (c) a point (d) a circle
46. Let  $A \equiv (-2, 0)$  and  $B \equiv (2, 0)$ , then the number of integral values of  $a$ ,  $a \in [-10, 10]$  for which line segment  $AB$  subtends an acute angle at point  $C \equiv (a, a + 1)$  is :  
 (a) 15 (b) 17 (c) 19 (d) 21
47. The angle between sides of a rhombus whose  $\sqrt{2}$  times sides is mean of its two diagonal, is equal to :  
 (a)  $300^\circ$  (b)  $45^\circ$  (c)  $60^\circ$  (d)  $90^\circ$
48. A rod of  $AB$  of length 3 rests on a wall as follows :



$P$  is a point on  $AB$  such that  $AP : PB = 1 : 2$ . If the rod slides along the wall, then the locus of  $P$  lies on

- (a)  $2x + y + xy = 2$  (b)  $4x^2 + xy + xy + y^2 = 4$   
 (c)  $4x^2 + y^2 = 4$  (d)  $x^2 + y^2 - x - 2y = 0$
49. If  $\frac{x^2}{a} + \frac{y^2}{b} + \frac{2xy}{h} = 0$ , represents pair of straight lines and slope of one line is twice the other. Then  $ab : h^2$  is :  
 (a)  $8 : 9$  (b)  $1 : 2$  (c)  $2 : 1$  (d)  $9 : 8$

50. Locus of point of reflection of point  $(a, 0)$  w.r.t. the line  $yt = x + at^2$  is given by ( $t$  is parameter,  $t \in \mathbb{R}$ ):
- (a)  $x - a = 0$                       (b)  $y - a = 0$                       (c)  $x + a = 0$                       (d)  $y + a = 0$
51. A light ray emerging from the point source placed at  $P(1, 3)$  is reflected at a point  $Q$  in the  $x$ -axis. If the reflected ray passes through  $R(6, 7)$ , then abscissa of  $Q$  is :
- (a)  $\frac{5}{2}$                                       (b) 3                                      (c)  $\frac{7}{2}$                                       (d) 1
52. If the axes are rotated through  $60^\circ$  in the anticlockwise sense, find the transformed form of the equation  $x^2 - y^2 = a^2$  :
- (a)  $X^2 + Y^2 - 3\sqrt{3}XY = 2a^2$                                       (b)  $X^2 + Y^2 = a^2$   
(c)  $Y^2 - X^2 - 2\sqrt{3}XY = 2a^2$                                       (d)  $X^2 - Y^2 + 2\sqrt{3}XY = 2a^2$
53. The straight line  $3x + y - 4 = 0$ ,  $x + 3y - 4 = 0$  and  $x + y = 0$  form a triangle which is :
- (a) equilateral                                      (b) right-angled  
(c) acute-angled and isosceles                                      (d) obtuse-angled and isosceles
54. If  $m$  and  $b$  are real numbers and  $mb > 0$ , then the line whose equation is  $y = mx + b$  cannot contain the point:
- (a)  $(0, 2008)$                                       (b)  $(2008, 0)$   
(c)  $(0, -2008)$                                       (d)  $(20, -100)$
55. The number of possible straight lines, passing through  $(2, 3)$  and forming a triangle with coordinate axes, whose area is 12 sq. units, is:
- (a) one    (b) two  
(c) three    (d) four
56. If  $x_1, x_2, x_3$  and  $y_1, y_2, y_3$  are both in G.P with the same common ratio then the points  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$
- (a) lie on a straight line                                      (b) lie on a circle  
(c) are vertices of a triangle                                      (d) None of these
57. Locus of centroid of the triangle whose vertices are  $(a \cos t, a \sin t)$ ,  $(b \sin t, -b \cos t)$  and  $(1, 0)$ ; where  $t$  is a parameter is :
- (a)  $(3x - 1)^2 + (3y)^2 = a^2 - b^2$                                       (b)  $(3x - 1)^2 + (3y)^2 = a^2 + b^2$   
(c)  $(3x + 1)^2 + (3y)^2 = a^2 + b^2$                                       (d)  $(3x + 1)^2 + (3y)^2 = a^2 - b^2$
58. The equation of the straight line passing through  $(4, 3)$  and making intercepts on co-ordinate axes whose sum is  $-1$  is :
- (a)  $\frac{x}{2} + \frac{y}{3} = -1$  and  $\frac{x}{-2} + \frac{y}{1} = -1$                                       (b)  $\frac{x}{2} - \frac{y}{3} = -1$  and  $\frac{x}{-2} + \frac{y}{1} = -1$   
(c)  $\frac{x}{2} + \frac{y}{3} = 1$  and  $\frac{x}{2} + \frac{y}{1} = 1$                                       (d)  $\frac{x}{2} - \frac{y}{3} = 1$  and  $\frac{x}{-2} + \frac{y}{1} = 1$

59. Let  $A \equiv (3, 2)$  and  $B \equiv (5, 1)$ .  $ABP$  is an equilateral triangle is constructed one the side of  $AB$  remote from the origin then the orthocentre of triangle  $ABP$  is:
- (a)  $\left(4 - \frac{1}{2}\sqrt{3}, \frac{3}{2} - \sqrt{3}\right)$  (b)  $\left(4 + \frac{1}{2}\sqrt{3}, \frac{3}{2} + \sqrt{3}\right)$   
 (c)  $\left(4 - \frac{1}{6}\sqrt{3}, \frac{3}{2} - \frac{1}{3}\sqrt{3}\right)$  (d)  $\left(4 + \frac{1}{6}\sqrt{3}, \frac{3}{2} + \frac{1}{3}\sqrt{3}\right)$
60. Area of the triangle formed by the lines through point  $(6, 0)$  and at a perpendicular distance of 5 from point  $(1, 3)$  and line  $y = 16$  in square units is :
- (a) 160 (b) 200 (c) 240 (d) 130
61. The straight lines  $3x + y - 4 = 0$ ,  $x + 3y - 4 = 0$  and  $x + y = 0$  form a triangle which is :
- (a) equilateral (b) right-angled  
 (c) acute-angled and isosceles (d) obtuse-angled and isosceles
62. The orthocentre of the triangle with vertices  $(5, 0)$ ,  $(0, 0)$ ,  $\left(\frac{5}{2}, \frac{5\sqrt{3}}{2}\right)$  is :
- (a)  $(2, 3)$  (b)  $\left(\frac{5}{2}, \frac{5}{2\sqrt{3}}\right)$  (c)  $\left(\frac{5}{6}, \frac{5}{2\sqrt{3}}\right)$  (d)  $\left(\frac{5}{2}, \frac{5}{\sqrt{3}}\right)$
63. All chords of a curve  $3x^2 - y^2 - 2x + 4y = 0$  which subtends a right angle at the origin passes through a fixed point, which is :
- (a)  $(1, 2)$  (b)  $(1, -2)$  (c)  $(2, 1)$  (d)  $(-2, 1)$
64. Let  $P(-1, 0)$ ,  $Q(0, 0)$ ,  $R(3, 3\sqrt{3})$  be three points then the equation of the bisector of the angle  $\angle PQR$  is :
- (a)  $\frac{\sqrt{3}}{2}x + y = 0$  (b)  $x + \sqrt{3}y = 0$  (c)  $\sqrt{3}x + y = 0$  (d)  $x + \frac{\sqrt{3}}{2}y = 0$

## Answers

1.	(b)	2.	(c)	3.	(c)	4.	(c)	5.	(c)	6.	(c)	7.	(b)	8.	(b)	9.	(b)	10.	(a)
11.	(d)	12.	(d)	13.	(d)	14.	(c)	15.	(d)	16.	(b)	17.	(c)	18.	(b)	19.	(d)	20.	(a)
21.	(a)	22.	(a)	23.	(c)	24.	(a)	25.	(c)	26.	(b)	27.	(d)	28.	(a)	29.	(b)	30.	(a)
31.	(c)	32.	(d)	33.	(c)	34.	(a)	35.	(d)	36.	(d)	37.	(c)	38.	(c)	39.	(d)	40.	(a)
41.	(d)	42.	(b)	43.	(d)	44.	(c)	45.	(a)	46.	(c)	47.	(d)	48.	(c)	49.	(d)	50.	(c)
51.	(a)	52.	(c)	53.	(d)	54.	(b)	55.	(c)	56.	(a)	57.	(b)	58.	(d)	59.	(d)	60.	(c)

**Exercise-2 : One or More than One Answer is/are Correct**

- A line makes intercepts on co-ordinate axes whose sum is 9 and their product is 20 ; then its equation is/are :
 

(a) $4x + 5y - 20 = 0$	(b) $5x + 4y - 20 = 0$
(c) $4x - 5y - 20 = 0$	(d) $4x + 5y + 20 = 0$
- The equation(s) of the medians of the triangle formed by the points (4, 8), (3, 2) and (5, -6) is/are :
 

(a) $x = 4$	(b) $x = 5y - 3$
(c) $2x + 3y - 12 = 0$	(d) $22x + 3y - 92 = 0$
- The value(s) of  $t$  for which the lines  $2x + 3y = 5$ ,  $t^2x + ty - 6 = 0$  and  $3x - 2y - 1 = 0$  are concurrent, can be :
 

(a) $t = 2$	(b) $t = -3$
(c) $t = -2$	(d) $t = 3$
- If one of the lines given by the equation  $ax^2 + 6xy + by^2 = 0$  bisects the angle between the co-ordinate axes, then value of  $(a + b)$  can be :
 

(a) -6	(b) 3	(c) 6	(d) 12
--------	-------	-------	--------
- Suppose  $ABCD$  is a quadrilateral such that the coordinates of  $A$ ,  $B$  and  $C$  are (1, 3), (-2, 6) and (5, -8) respectively. For what choices of coordinates of  $D$  will make  $ABCD$  a trapezium ?
 

(a) (3, -6)	(b) (6, -9)	(c) (0, 5)	(d) (3, -1)
-------------	-------------	------------	-------------
- One diagonal of a square is the portion of the line  $\sqrt{3}x + y = 2\sqrt{3}$  intercepted by the axes. Then an extremity of the other diagonal is :
 

(a) $(1 + \sqrt{3}, \sqrt{3} - 1)$	(b) $(1 + \sqrt{3}, \sqrt{3} + 1)$
(c) $(1 - \sqrt{3}, \sqrt{3} - 1)$	(d) $(1 - \sqrt{3}, \sqrt{3} + 1)$
- Two sides of a rhombus  $ABCD$  are parallel to lines  $y = x + 2$  and  $y = 7x + 3$ . If the diagonals of the rhombus intersect at point (1, 2) and the vertex  $A$  is on the  $y$ -axis is, then the possible coordinates of  $A$  are:
 

(a) $(0, \frac{5}{2})$	(b) (0, 0)	(c) (0, 5)	(d) (0, 3)
------------------------	------------	------------	------------
- The equation of the sides of the triangle having (3, -1) as a vertex and  $x - 4y + 10 = 0$  and  $6x + 10y - 59 = 0$  as angle bisector and as median respectively drawn from different vertices, are :
 

(a) $6x + 7y - 13 = 0$	(b) $2x + 9y - 65 = 0$
(c) $18x + 13y - 41 = 0$	(d) $6x - 7y - 25 = 0$
- $A(1,3)$  and  $C(5, 1)$  are two opposite vertices of a rectangle  $ABCD$ . If the slope of  $BD$  is 2, then the coordinates of  $B$  can be :
 

(a) (4, 4)	(b) (5, 4)
(c) (2, 0)	(d) (1, 0)




10. All the points lying inside the triangle formed by the points (1, 3), (5, 6), and (-1, 2) satisfy:
- (a)  $3x + 2y \geq 0$  (b)  $2x + y + 1 \geq 0$   
 (c)  $-2x + 11 \geq 0$  (d)  $2x + 3y - 12 \geq 0$
11. The slope of a median, drawn from the vertex  $A$  of the triangle  $ABC$  is  $-2$ . The co-ordinates of vertices  $B$  and  $C$  are respectively  $(-1, 3)$  and  $(3, 5)$ . If the area of the triangle be 5 square units, then possible distance of vertex  $A$  from the origin is/are.
- (a) 6 (b) 4 (c)  $2\sqrt{2}$  (d)  $3\sqrt{2}$
12. The points  $A(0, 0)$ ,  $B(\cos \alpha, \sin \alpha)$  and  $C(\cos \beta, \sin \beta)$  are the vertices of a right angled triangle if:
- (a)  $\sin\left(\frac{\alpha - \beta}{2}\right) = \frac{1}{\sqrt{2}}$  (b)  $\cos\left(\frac{\alpha - \beta}{2}\right) = -\frac{1}{\sqrt{2}}$   
 (c)  $\cos\left(\frac{\alpha - \beta}{2}\right) = \frac{1}{\sqrt{2}}$  (d)  $\sin\left(\frac{\alpha - \beta}{2}\right) = -\frac{1}{\sqrt{2}}$

### Answers

1.	(a, b)	2.	(a, c, d)	3.	(a, b)	4.	(a, c)	5.	(b, d)	6.	(b, c)
7.	(a, b)	8.	(b, c, d)	9.	(a, c)	10.	(a, b, c, d)	11.	(a, c)	12.	(a, b, c)




**Exercise-4 : Matching Type Problems**

1.

Column-I		Column-II	
(A)	If $a, b, c$ are in A.P, then lines $ax + by + c = 0$ are concurrent at:	(P)	$(-4, -7)$
(B)	A point on the line $x + y = 4$ which lies at a unit distance from the line $4x + 3y = 10$ is :	(Q)	$(-7, 11)$
(C)	Orthocentre of triangle made by lines $x + y = 1$ , $x - y + 3 = 0$ , $2x + y = 7$ is	(R)	$(1, -2)$
(D)	Two vertex of a triangle are $(5, -1)$ and $(-2, 3)$ . If orthocentre is the origin then coordinates of the third vertex are	(S)	$(-1, 2)$
		(T)	$(0, 0)$

2.

Column-I		Column-II	
(A)	If $\sum_{r=1}^{n+1} \left( \sum_{k=1}^n {}^k C_{r-1} \right) = 30$ , then $n$ is equal to	(P)	1
(B)	The number of integral values of $g$ for which atmost one member of the family of lines given by $(1 + 2\lambda)x + (1 - \lambda)y + 2 + 4\lambda = 0$ ( $\lambda$ is real parameter) is tangent to the circle $x^2 + y^2 + 4gx + 18x + 17y + 4g^2 = 0$ can be	(Q)	4
(C)	Number of solutions of the equation $\sin 9x + \sin 5x + 2\sin^2 x = 1$ in interval $(0, \pi)$ is	(R)	7
(D)	If the roots of the equation $x^2 + ax + b = 0$ ( $a, b \in R$ ) are $\tan 65^\circ$ and $\tan 70^\circ$ , then $(a + b)$ equals.	(S)	10

3.

Column-I		Column-II	
(A)	Exact value of $\cos 40^\circ (1 - 2\sin 10^\circ) =$	(P)	$\frac{1}{4}$

<b>(B)</b>	Value of $\lambda$ for which lines are concurrent $x + y + 1 = 0$ , $3x + 2\lambda y + 4 = 0$ , $x + y - 3\lambda = 0$ can be	<b>(Q)</b>	$\frac{1}{2}$
<b>(C)</b>	Points $(k, 2 - 2k)$ , $(-k + 1, 2k)$ and $(-4 - k, 6 - 2k)$ are collinear then sum of all possible real values of 'k' is	<b>(R)</b>	$\frac{3}{2}$
<b>(D)</b>	Value of $\sum_{k=3}^{\infty} \sin^k\left(\frac{\pi}{6}\right) =$	<b>(S)</b>	$-\frac{1}{2}$

### Answers

1. A $\rightarrow$ R; B $\rightarrow$ Q; C $\rightarrow$ S; D $\rightarrow$ P
2. A $\rightarrow$ Q; B $\rightarrow$ R; C $\rightarrow$ S; D $\rightarrow$ P
3. A $\rightarrow$ Q; B $\rightarrow$ R; C $\rightarrow$ S; D $\rightarrow$ P

### Exercise-5 : Subjective Type Problems

1. If the area of the quadrilateral  $ABCD$  whose vertices are  $A(1, 1)$ ,  $B(7, -3)$ ,  $C(12, 2)$  and  $D(7, 21)$  is  $\Delta$ . Find the sum of the digits of  $\Delta$ .
2. The equation of a line through the mid-point of the sides  $AB$  and  $AD$  of rhombus  $ABCD$ , whose one diagonal is  $3x - 4y + 5 = 0$  and one vertex is  $A(3, 1)$  is  $ax + by + c = 0$ . Find the absolute value of  $(a + b + c)$  where  $a, b, c$  are integers expressed in lowest form.
3. If the point  $(\alpha, \alpha^4)$  lies on or inside the triangle formed by lines  $x^2y + xy^2 - 2xy = 0$ , then the largest value of  $\alpha$  is.
4. The minimum value of  $[(x_1 - x_2)^2 + (12 - \sqrt{1 - x_1^2} - \sqrt{4x_2})^2]^{1/2}$  for all permissible values of  $x_1$  and  $x_2$  is equal to  $a\sqrt{b} - c$  where  $a, b, c \in N$ , then find the value of  $a + b - c$ .
5. The number of lines that can be drawn passing through point  $(2, 3)$  so that its perpendicular distance from  $(-1, 6)$  is equal to 6 is :
6. The graph of  $x^4 = x^2y^2$  is a union of  $n$  different lines, then the value of  $n$  is.
7. The orthocentre of triangle formed by lines  $x + y - 1 = 0$ ,  $2x + y - 1 = 0$  and  $y = 0$  is  $(h, k)$ , then  $\frac{1}{k^2} =$
8. Find the integral value of  $a$  for which the point  $(-2, a)$  lies in the interior of the triangle formed by the lines  $y = x$ ,  $y = -x$  and  $2x + 3y = 6$ .
9. Let  $A = (-1, 0)$ ,  $B = (3, 0)$  and  $PQ$  be any line passing through  $(4, 1)$ . The range of the slope of  $PQ$  for which there are two points on  $PQ$  at which  $AB$  subtends a right angle is  $(\lambda_1, \lambda_2)$ , then  $5(\lambda_1 + \lambda_2)$  is equal to.
10. Given that the three points where the curve  $y = bx^2 - 2$  intersects the  $x$ -axis and  $y$ -axis form an equilateral triangle. Find the value of  $2b$ .

### Answers

1.	6	2.	1	3.	1	4.	8	5.	0	6.	3	7.	4
8.	3	9.	6	10.	5								



### Exercise-1 : Single Choice Problems

- The locus of mid-points of the chords of the circle  $x^2 - 2x + y^2 - 2y + 1 = 0$  which are of unit length is :
 

(a) $(x-1)^2 + (y-1)^2 = \frac{3}{4}$	(b) $(x-1)^2 + (y-1)^2 = 2$
(c) $(x-1)^2 + (y-1)^2 = \frac{1}{4}$	(d) $(x-1)^2 + (y-1)^2 = \frac{2}{3}$
- The length of a common internal tangent to two circles is 5 and a common external tangent is 15, then the product of the radii of the two circles is :
 

(a) 25	(b) 50	(c) 75	(d) 30
--------	--------	--------	--------
- A circle with center (2, 2) touches the coordinate axes and a straight line  $AB$  where  $A$  and  $B$  lie on positive direction of coordinate axes such that the circle lies between origin and the line  $AB$ . If  $O$  be the origin then the locus of circumcenter of  $\triangle OAB$  will be:
 

(a) $xy = x + y + \sqrt{x^2 + y^2}$	(b) $xy = x + y - \sqrt{x^2 + y^2}$
(c) $xy + x + y = \sqrt{x^2 + y^2}$	(d) $xy + x + y + \sqrt{x^2 + y^2} = 0$
- Length of chord of contact of point (4, 4) with respect to the circle  $x^2 + y^2 - 2x - 2y - 7 = 0$  is:
 

(a) $\frac{3}{\sqrt{2}}$	(b) $3\sqrt{2}$	(c) 3	(d) 6
--------------------------	-----------------	-------	-------
- Let  $P, Q, R, S$  be the feet of the perpendiculars drawn from a point (1, 1) upon the lines  $x + 4y = 12$ ;  $x - 4y + 4 = 0$  and their angle bisectors respectively; then equation of the circle which passes through  $Q, R, S$  is :
 

(a) $x^2 + y^2 - 5x + 3y - 6 = 0$	(b) $x^2 + y^2 - 5x - 3y + 6 = 0$
(c) $x^2 + y^2 - 5x - 3y - 6 = 0$	(d) None of these

6. From a point 'P' on the line  $2x + y + 4 = 0$ ; which is nearest to the circle  $x^2 + y^2 - 12y + 35 = 0$ , tangents are drawn to given circle. The area of quadrilateral  $PACB$  (where 'C' is the center of circle and  $PA$  &  $PB$  are the tangents.) is :
- (a) 8 (b)  $\sqrt{110}$  (c)  $\sqrt{19}$  (d) None of these
7. The line  $2x - y + 1 = 0$  is tangent to the circle at the point (2, 5) and the centre of the circles lies on  $x - 2y = 4$ . The radius of the circle is:
- (a)  $3\sqrt{5}$  (b)  $5\sqrt{3}$   
(c)  $2\sqrt{5}$  (d)  $5\sqrt{2}$
8. If  $A(\cos \alpha, \sin \alpha)$ ,  $B(\sin \alpha, -\cos \alpha)$ ,  $C(1, 2)$  are the vertices of a triangle, then as  $\alpha$  varies the locus of centroid of the  $\Delta ABC$  is a circle whose radius is :
- (a)  $\frac{2\sqrt{2}}{3}$  (b)  $\frac{\sqrt{4}}{3}$  (c)  $\frac{2}{3}$  (d)  $\frac{\sqrt{2}}{9}$
9. Tangents drawn to circle  $(x - 1)^2 + (y - 1)^2 = 5$  at point  $P$  meets the line  $2x + y + 6 = 0$  at  $Q$  on the  $x$ -axis. Length  $PQ$  is equal to :
- (a)  $\sqrt{12}$  (b)  $\sqrt{10}$  (c) 4 (d)  $\sqrt{15}$
10.  $ABCD$  is square in which  $A$  lies on positive  $y$ -axis and  $B$  lies on the positive  $x$ -axis. If  $D$  is the point (12, 17), then co-ordinate of  $C$  is :
- (a) (17, 12) (b) (17, 5) (c) (17, 16) (d) (15, 3)
11. **Statement-1:** The lines  $y = mx + 1 - m$  for all values of  $m$  is a normal to the circle  $x^2 + y^2 - 2x - 2y = 0$ .
- Statement-2:** The line  $L$  passes through the centre of the circle.
- (a) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
(b) Statement-1 is true, statement-2 is true and statement-2 is not the correct explanation for statement-1.  
(c) Statement-1 is true, statement-2 is false.  
(d) Statement-1 is false, statement-2 is true.
12.  $A(1, 0)$  and  $B(0, 1)$  are two fixed points on the circle  $x^2 + y^2 = 1$ .  $C$  is a variable point on this circle. As  $C$  moves, the locus of the orthocentre of the triangle  $ABC$  is :
- (a)  $x^2 + y^2 - 2x - 2y + 1 = 0$  (b)  $x^2 + y^2 - x - y = 0$   
(c)  $x^2 + y^2 = 4$  (d)  $x^2 + y^2 + 2x - 2y + 1 = 0$
13. Equation of a circle passing through (1, 2) and (2, 1) and for which line  $x + y = 2$  is a diameter ; is :
- (a)  $x^2 + y^2 + 2x + 2y - 11 = 0$  (b)  $x^2 + y^2 - 2x - 2y - 1 = 0$   
(c)  $x^2 + y^2 - 2x - 2y + 1 = 0$  (d) None of these

14. The area of an equilateral triangle inscribed in a circle of radius 4 cm, is :  
(a)  $12 \text{ cm}^2$  (b)  $9\sqrt{3} \text{ cm}^2$   
(c)  $8\sqrt{3} \text{ cm}^2$  (d)  $12\sqrt{3} \text{ cm}^2$
15. Let all the points on the curve  $x^2 + y^2 - 10x = 0$  are reflected about the line  $y = x + 3$ . The locus of the reflected points is in the form  $x^2 + y^2 + gx + fy + c = 0$ . The value of  $(g + f + c)$  is equal to :  
(a) 28 (b) -28 (c) 38 (d) -38
16. The shortest distance from the line  $3x + 4y = 25$  to the circle  $x^2 + y^2 = 6x - 8y$  is equal to:  
(a)  $7/5$  (b)  $9/5$  (c)  $11/5$  (d)  $32/5$
17. In the  $xy$ -plane, the length of the shortest path from  $(0, 0)$  to  $(12, 16)$  that does not go inside the circle  $(x - 6)^2 + (y - 8)^2 = 25$  is:  
(a)  $10\sqrt{3}$  (b)  $10\sqrt{5}$   
(c)  $10\sqrt{3} + \frac{5\pi}{3}$  (d)  $10 + 5\pi$
18. A circle is inscribed in an equilateral triangle with side lengths 6 unit. Another circle is drawn inside the triangle (but outside the first circle), tangent to the first circle and two of the sides of the triangle. The radius of the smaller circle is:  
(a)  $1/\sqrt{3}$  (b)  $2/3$   
(c)  $1/2$  (d) 1
19. The equation of the tangent to the circle  $x^2 + y^2 - 4x = 0$  which is perpendicular to the normal drawn through the origin can be :  
(a)  $x = 1$  (b)  $x = 2$  (c)  $x + y = 2$  (d)  $x = 4$
20. The equation of the line parallel to the line  $3x + 4y = 0$  and touching the circle  $x^2 + y^2 = 9$  in the first quadrant is :  
(a)  $3x + 4y = 15$  (b)  $3x + 4y = 45$   
(c)  $3x + 4y = 9$  (d)  $3x + 4y = 12$
21. The centres of the three circles  $x^2 + y^2 - 10x + 9 = 0$ ,  $x^2 + y^2 - 6x + 2y + 1 = 0$ ,  $x^2 + y^2 - 9x - 4y + 2 = 0$   
(a) lie on the straight line  $x - 2y = 5$  (b) lie on circle  $x^2 + y^2 = 25$   
(c) do not lie on straight line (d) lie on circle  $x^2 + y^2 + x + y - 17 = 0$
22. The equation of the diameter of the circle  $x^2 + y^2 + 2x - 4y = 4$  that is parallel to  $3x + 5y = 4$  is:  
(a)  $3x + 5y = -7$  (b)  $3x + 5y = 7$   
(c)  $3x + 5y = 9$  (d)  $3x + 5y = 1$



23. There are two circles passing through points  $A(-1, 2)$  and  $B(2, 3)$  having radius  $\sqrt{5}$ . Then the length of intercept on  $x$ -axis of the circle intersecting  $x$ -axis is :  
 (a) 2 (b) 3 (c) 4 (d) 5
24. A square  $OABC$  is formed by line pairs  $xy = 0$  and  $xy + 1 = x + y$  where ' $O$ ' is the origin. A circle with centre  $C_1$  inside the square is drawn to touch the line pair  $xy = 0$  and another circle with centre  $C_2$  and radius twice that of  $C_1$ , is drawn to touch the circle  $C_1$  and the other line pair. The radius of the circle with centre  $C_1$  is:  
 (a)  $\frac{\sqrt{2}}{\sqrt{3}(\sqrt{2}+1)}$  (b)  $\frac{2\sqrt{2}}{3(\sqrt{2}+1)}$   
 (c)  $\frac{\sqrt{2}}{3(\sqrt{2}+1)}$  (d)  $\frac{\sqrt{2}+1}{3\sqrt{2}}$
25. The equation of the circle circumscribing the triangle formed by the points  $(3, 4)$ ,  $(1, 4)$  and  $(3, 2)$  is :  
 (a)  $8x^2 + 8y^2 - 16x - 13y = 0$  (b)  $x^2 + y^2 - 4x - 8y + 19 = 0$   
 (c)  $x^2 + y^2 - 4x - 6y + 11 = 0$  (d)  $x^2 + y^2 - 6x - 6y + 17 = 0$
26. The equation of the tangent to circle  $x^2 + y^2 + 2gx + 2fy = 0$  at the origin is :  
 (a)  $fx + gy = 0$  (b)  $gx + fy = 0$  (c)  $x = 0$  (d)  $y = 0$
27. The line  $y = x$  is tangent at  $(0, 0)$  to a circle of radius 1. The centre of the circle is :  
 (a) either  $\left(-\frac{1}{2}, \frac{1}{2}\right)$  or  $\left(\frac{1}{2}, -\frac{1}{2}\right)$  (b) either  $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  or  $\left(-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$   
 (c) either  $\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$  or  $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  (d) either  $(1, 0)$  or  $(-1, 0)$
28. The circles  $x^2 + y^2 + 6x + 6y = 0$  and  $x^2 + y^2 - 12x - 12y = 0$  :  
 (a) cut orthogonally (b) touch each other internally  
 (c) intersect in two points (d) touch each other externally
29. In a right triangle  $ABC$ , right angled at  $A$ , on the leg  $AC$  as diameter, a semicircle is described. The chord joining  $A$  with the point of intersection  $D$  of the hypotenuse and the semicircle, then the length  $AC$  equals to:  
 (a)  $\frac{AB \cdot AD}{\sqrt{AB^2 + AD^2}}$  (b)  $\frac{AB \cdot AD}{AB + AD}$   
 (c)  $\sqrt{AB \cdot AD}$  (d)  $\frac{AB \cdot AD}{\sqrt{AB^2 - AD^2}}$
30. Radical centre of the circles drawn on the sides as a diameter of triangle formed by the lines  $3x - 4y + 6 = 0$ ,  $x - y + 2 = 0$  and  $4x + 3y - 17 = 0$  is :  
 (a)  $(3, 2)$  (b)  $(3, -2)$  (c)  $(2, -3)$  (d)  $(2, 3)$

- 31. Statement-1:** A circle can be inscribed in a quadrilateral whose sides are  $3x - 4y = 0$ ,  $3x - 4y = 5$ ,  $3x + 4y = 0$  and  $3x + 4y = 7$ .
- Statement-2:** A circle can be inscribed in a parallelogram if and only if it is a rhombus.
- (a) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
 (b) Statement-1 is true, statement-2 is true and statement-2 is not the correct explanation for statement-1.  
 (c) Statement-1 is true, statement-2 is false.  
 (d) Statement-1 is false, statement-2 is true.
- 32.** If  $x = 3$  is the chord of contact of the circle  $x^2 + y^2 = 81$ , then the equation of the corresponding pair of tangents, is:  
 (a)  $x^2 - 8y^2 + 54x + 729 = 0$ .  
 (b)  $x^2 - 8y^2 - 54x + 729 = 0$   
 (c)  $x^2 - 8y^2 - 54x - 729 = 0$   
 (d)  $x^2 - 8y^2 = 729$
- 33.** The shortest distance from the line  $3x + 4y = 25$  to the circle  $x^2 + y^2 = 6x - 8y$  is equal to :  
 (a)  $\frac{7}{3}$  (b)  $\frac{9}{5}$  (c)  $\frac{11}{5}$  (d)  $\frac{7}{5}$
- 34.** The circle with equation  $x^2 + y^2 = 1$  intersects the line  $y = 7x + 5$  at two distinct points  $A$  and  $B$ . Let  $C$  be the point at which the positive  $x$ -axis intersects the circle. The angle  $ACB$  is :  
 (a)  $\tan^{-1} \frac{4}{3}$  (b)  $\cot^{-1}(-1)$  (c)  $\tan^{-1} 1$  (d)  $\cot^{-1} \frac{4}{3}$
- 35.** The abscissae of two points  $A$  and  $B$  are the roots of the equation  $x^2 + 2ax - b^2 = 0$  and their ordinates are the roots of the equation  $x^2 + 2px - q^2 = 0$ . The radius of the circle with  $AB$  as diameter is ::  
 (a)  $\sqrt{a^2 + b^2 + p^2 + q^2}$  (b)  $\sqrt{a^2 + p^2}$   
 (c)  $\sqrt{b^2 + q^2}$  (d)  $\sqrt{a^2 + b^2 + p^2 + 1}$
- 36.** Let  $C$  be the circle of radius unity centred at the origin. If two positive numbers  $x_1$  and  $x_2$  are such that the line passing through  $(x_1, -1)$  and  $(x_2, 1)$  is tangent to  $C$  then:  
 (a)  $x_1 x_2 = 1$  (b)  $x_1 x_2 = -1$   
 (c)  $x_1 + x_2 = 1$  (d)  $4x_1 x_2 = 1$
- 37.** A circle bisects the circumference of the circle  $x^2 + y^2 + 2y - 3 = 0$  and touches the line  $x = y$  at the point  $(1, 1)$ . Its radius is :  
 (a)  $\frac{3}{\sqrt{2}}$  (b)  $\frac{9}{\sqrt{2}}$  (c)  $4\sqrt{2}$  (d)  $3\sqrt{2}$
- 38.** The distance between the chords of contact of tangents to the circle  $x^2 + y^2 + 2gx + 2fy + c = 0$  from the origin and the point  $(g, f)$  is:

$$(a) \sqrt{g^2 + f^2} \qquad (b) \frac{\sqrt{g^2 + f^2 - c}}{2}$$

$$(c) \frac{g^2 + f^2 - c}{2\sqrt{g^2 + f^2}} \qquad (d) \frac{\sqrt{g^2 + f^2 + c}}{2\sqrt{g^2 + f^2}}$$

39. If the tangents  $AP$  and  $AQ$  are drawn from the point  $A(3, -1)$  to the circle  $x^2 + y^2 - 3x + 2y - 7 = 0$  and  $C$  is the centre of circle, then the area of quadrilateral  $APCQ$  is :  
 (a) 9 (b) 4 (c) 2 (d) non-existent
40. Number of integral value(s) of  $k$  for which no tangent can be drawn from the point  $(k, k + 2)$  to the circle  $x^2 + y^2 = 4$  is :  
 (a) 0 (b) 1 (c) 2 (d) 3
41. If the length of the normal for each point on a curve is equal to the radius vector, then the curve :  
 (a) is a circle passing through origin  
 (b) is a circle having centre at origin and radius  $> 0$   
 (c) is a circle having centre on  $x$ -axis and touching  $y$ -axis  
 (d) is a circle having centre on  $y$ -axis and touching  $x$ -axis
42. A circle of radius unity is centred at origin. Two particles start moving at the same time from the point  $(1, 0)$  and move around the circle in opposite direction. One of the particle moves counter clockwise with constant speed  $v$  and the other moves clockwise with constant speed  $3v$ . After leaving  $(1, 0)$ , the two particles meet first at a point  $P$ , and continue until they meet next at point  $Q$ . The coordinates of the point  $Q$  are:  
 (a)  $(1, 0)$  (b)  $(0, 1)$   
 (c)  $(0, -1)$  (d)  $(-1, 0)$
43. A variable circle is drawn to touch the  $x$ -axis at the origin. The locus of the pole of the straight line  $lx + my + n = 0$  w.r.t the variable circle has the equation:  
 (a)  $x(my - n) - ly^2 = 0$  (b)  $x(my + n) - ly^2 = 0$   
 (c)  $x(my - n) + ly^2 = 0$  (d) none of these
44. The minimum length of the chord of the circle  $x^2 + y^2 + 2x + 2y - 7 = 0$  which is passing through  $(1, 0)$  is :  
 (a) 2 (b) 4 (c)  $2\sqrt{2}$  (d)  $\sqrt{5}$
45. Three concentric circles of which the biggest is  $x^2 + y^2 = 1$ , have their radii in A.P. If the line  $y = x + 1$  cuts all the circles in real and distinct points. The interval in which the common difference of the A.P. will lie is:  
 (a)  $\left(0, \frac{1}{4}\right)$  (b)  $\left(0, \frac{1}{2\sqrt{2}}\right)$  (c)  $\left(0, \frac{2 - \sqrt{2}}{4}\right)$  (d) none

46. The locus of the point of intersection of the tangent to the circle  $x^2 + y^2 = a^2$ , which include an angle of  $45^\circ$  is the curve  $(x^2 + y^2)^2 = \lambda a^2(x^2 + y^2 - a^2)$ . The value of  $\lambda$  is:
- (a) 2 (b) 4  
(c) 8 (d) 16
47. A circle touches the line  $y = x$  at point  $(4, 4)$  on it. The length of the chord on the line  $x + y = 0$  is  $6\sqrt{2}$ . Then one of the possible equation of the circle is :
- (a)  $x^2 + y^2 + x - y + 30 = 0$  (b)  $x^2 + y^2 + 2x - 18y + 32 = 0$   
(c)  $x^2 + y^2 + 2x + 18y + 32 = 0$  (d)  $x^2 + y^2 - 2x - 22y + 32 = 0$
48. Point on the circle  $x^2 + y^2 - 2x + 4y - 4 = 0$  which is nearest to the line  $y = 2x + 11$  is :
- (a)  $\left(1 - \frac{6}{\sqrt{5}}, -2 + \frac{3}{\sqrt{5}}\right)$  (b)  $\left(1 + \frac{6}{\sqrt{5}}, -2 - \frac{3}{\sqrt{5}}\right)$   
(c)  $\left(1 - \frac{6}{\sqrt{5}}, -2 - \frac{3}{\sqrt{5}}\right)$  (d) None of these
49. A foot of the normal from the point  $(4, 3)$  to a circle is  $(2, 1)$  and a diameter of the circle has the equation  $2x - y - 2 = 0$ . Then the equation of the circle is:
- (a)  $x^2 + y^2 - 4y + 2 = 0$  (b)  $x^2 + y^2 - 4y + 1 = 0$   
(c)  $x^2 + y^2 - 2x - 1 = 0$  (d)  $x^2 + y^2 - 2x + 1 = 0$
50. If  $\left(a, \frac{1}{a}\right)$ ,  $\left(b, \frac{1}{b}\right)$ ,  $\left(c, \frac{1}{c}\right)$  and  $\left(d, \frac{1}{d}\right)$  are four distinct points on a circle of radius 4 units then,  $abcd$  is equal to:
- (a) 4 (b)  $\frac{1}{4}$  (c) 1 (d) 16

### Answers

1.	(a)	2.	(b)	3.	(a)	4.	(b)	5.	(b)	6.	(c)	7.	(a)	8.	(d)	9.	(a)	10.	(b)
11.	(a)	12.	(a)	13.	(c)	14.	(d)	15.	(c)	16.	(a)	17.	(c)	18.	(a)	19.	(d)	20.	(a)
21.	(c)	22.	(b)	23.	(c)	24.	(c)	25.	(c)	26.	(b)	27.	(c)	28.	(d)	29.	(d)	30.	(d)
31.	(d)	32.	(b)	33.	(d)	34.	(c)	35.	(a)	36.	(a)	37.	(b)	38.	(c)	39.	(d)	40.	(b)
41.	(b)	42.	(d)	43.	(a)	44.	(b)	45.	(c)	46.	(c)	47.	(b)	48.	(a)	49.	(c)	50.	(c)

### Exercise-2 : One or More than One Answer is/are Correct

- Number of circle touching both the axes and the line  $x + y = 4$  is greater than or equal to :
  - 1
  - 2
  - 3
  - 4
- Which of the following is/are true ?  
The circles  $x^2 + y^2 - 6x - 6y + 9 = 0$  and  $x^2 + y^2 + 6x + 6y + 9 = 0$  are such that :
  - They do not intersect
  - They touch each other
  - Their exterior common tangents are parallel
  - Their interior common tangents are perpendicular
- Let ' $\alpha$ ' be a variable parameter, then the length of the chord of the curve :  

$$(x - \sin^{-1} \alpha)(x - \cos^{-1} \alpha) + (y - \sin^{-1} \alpha)(y + \cos^{-1} \alpha) = 0$$
 along the line  $x = \frac{\pi}{4}$  can not be equal to :
  - $\frac{\pi}{3}$
  - $\frac{\pi}{6}$
  - $\frac{\pi}{4}$
  - $\frac{\pi}{2}$
- If the point  $(1, 4)$  lies inside the circle  $x^2 + y^2 - 6x - 10y + p = 0$  and the circle does not touch or intersect the coordinate axes, then which of the following must be correct :
  - $p < 29$
  - $p > 25$
  - $p > 27$
  - $p < 27$
- The equation of a circle  $S_1 = 0$  is  $x^2 + y^2 = 4$ , locus of the intersection of orthogonal tangents to the circle is the curve  $C_1$  and the locus of the intersection of perpendicular tangents to the curve  $C_1$  is the curve  $C_2$ , then :
  - $C_2$  is a circle
  - $C_1, C_2$  are circles having different centres
  - $C_1, C_2$  are circles having same centres
  - area enclosed between  $C_1$  and  $C_2$  is  $8\pi$
- If two distinct chords drawn from the point  $(p, q)$  on the circle  $x^2 + y^2 = px + qy$  (where  $pq \neq 0$ ) are bisected by the  $x$ -axis, then :
  - $p^2 = q^2$
  - $p^2 > q^2$
  - $p^2 < 8q^2$
  - $p^2 > 8q^2$
- If  $a = \max\{(x+2)^2 + (y-3)^2\}$  and  $b = \min\{(x+2)^2 + (y-3)^2\}$  where  $x, y$  satisfying  $x^2 + y^2 + 8x - 10y - 40 = 0$ , then :
  - $a + b = 18$
  - $a + b = 178$
  - $a - b = 4\sqrt{2}$
  - $a - b = 72\sqrt{2}$

8. The locus of points of intersection of the tangents to  $x^2 + y^2 = a^2$  at the extremities of a chord of circle  $x^2 + y^2 = a^2$  which touches the circle  $x^2 + y^2 - 2ax = 0$  is/are :
- (a)  $y^2 = a(a - 2x)$  (b)  $x^2 = a(a - 2y)$   
(c)  $x^2 + y^2 = (x - a)^2$  (d)  $x^2 + y^2 = (y - a)^2$
9. A circle passes through the points  $(-1, 1)$ ,  $(0, 6)$  and  $(5, 5)$ . The point(s) on this circle, the tangent(s) at which is/are parallel to the straight line joining the origin to its centre is/are  
(a)  $(1, -5)$  (b)  $(5, 1)$  (c)  $(-5, -1)$  (d)  $(-1, 5)$
10. A square is inscribed in the circle  $x^2 + y^2 - 2x + 4y - 93 = 0$  with the sides parallel to the co-ordinate axes. The co-ordinate of the vertices are :  
(a)  $(8, 5)$  (b)  $(8, 9)$  (c)  $(-6, 5)$  (d)  $(-6, -9)$

## Answers

1.	(a, b, c, d)	2.	(a, c, d)	3.	(a, b, c)	4.	(a, b)	5.	(a, c, d)	6.	(b, d)
7.	(b, d)	8.	(a, c)	9.	(b, d)	10.	(a, c)				

### Exercise-3 : Comprehension Type Problems

#### Paragraph for Question Nos. 1 to 3

Let each of the circles,

$$S_1 \equiv x^2 + y^2 + 4y - 1 = 0,$$

$$S_2 \equiv x^2 + y^2 + 6x + y + 8 = 0,$$

$$S_3 \equiv x^2 + y^2 - 4x - 4y - 37 = 0$$

touches the other two. Let  $P_1, P_2, P_3$  be the points of contact of  $S_1$  and  $S_2$ ,  $S_2$  and  $S_3$ ,  $S_3$  and  $S_1$  respectively and  $C_1, C_2, C_3$  be the centres of  $S_1, S_2, S_3$  respectively.

- The co-ordinates of  $P_1$  are :  
 (a)  $(2, -1)$       (b)  $(2, 1)$       (c)  $(-2, 1)$       (d)  $(-2, -1)$
- The ratio  $\frac{\text{area}(\Delta P_1 P_2 P_3)}{\text{area}(\Delta C_1 C_2 C_3)}$  is equal to :  
 (a)  $3 : 2$       (b)  $2 : 5$       (c)  $5 : 3$       (d)  $2 : 3$
- $P_2$  and  $P_3$  are image of each other with respect to line :  
 (a)  $y = x + 1$       (b)  $y = -x$       (c)  $y = x$       (d)  $y = -x + 2$

#### Paragraph for Question Nos. 4 to 6

Let  $A(3, 7)$  and  $B(6, 5)$  are two points.  $C : x^2 + y^2 - 4x - 6y - 3 = 0$  is a circle.

- The chords in which the circle  $C$  cuts the members of the family  $S$  of circle passing through  $A$  and  $B$  are concurrent at :  
 (a)  $(2, 3)$       (b)  $\left(2, \frac{23}{3}\right)$       (c)  $\left(3, \frac{23}{2}\right)$       (d)  $(3, 2)$
- Equation of the member of the family of circles  $S$  that bisects the circumference of  $C$  is :  
 (a)  $x^2 + y^2 - 5x - 1 = 0$       (b)  $x^2 + y^2 - 5x + 6y - 1 = 0$   
 (c)  $x^2 + y^2 - 5x - 6y - 1 = 0$       (d)  $x^2 + y^2 + 5x - 6y - 1 = 0$
- If  $O$  is the origin and  $P$  is the center of  $C$ , then absolute value of difference of the squares of the lengths of the tangents from  $A$  and  $B$  to the circle  $C$  is equal to :  
 (a)  $(AB)^2$       (b)  $(OP)^2$       (c)  $|(AP)^2 - (BP)^2|$       (d)  $(AP)^2 + (BP)^2$

#### Paragraph for Question Nos. 7 to 8

Let the diameter of a subset  $S$  of the plane be defined as the maximum of the distance between arbitrary pairs of points of  $S$ .

- Let  $S = \{(x, y) : (y - x) \leq 0, x + y \geq 0, x^2 + y^2 \leq 2\}$ , then the diameter of  $S$  is :  
 (a) 2      (b) 4      (c)  $\sqrt{2}$       (d)  $2\sqrt{2}$

8. Let  $S = \{(x, y) : (\sqrt{5} - 1)x - \sqrt{10 + 2\sqrt{5}} y \geq 0, (\sqrt{5} - 1)x + \sqrt{10 + 12\sqrt{5}} y \geq 0, x^2 + y^2 \leq 9\}$  then the diameter of  $S$  is :

- (a)  $\frac{3}{2}(\sqrt{5} - 1)$       (b)  $3(\sqrt{5} - 1)$       (c)  $3\sqrt{2}$       (d) 3

**Paragraph for Question Nos. 9 to 10**

Let  $L_1, L_2$  and  $L_3$  be the lengths of tangents drawn from a point  $P$  to the circles  $x^2 + y^2 = 4$ ,  $x^2 + y^2 - 4x = 0$  and  $x^2 + y^2 - 4y = 0$  respectively. If  $L_1^4 = L_2^2 L_3^2 + 16$  then the locus of  $P$  are the curves,  $C_1$  (a straight line) and  $C_2$  (a circle).

9. Circum centre of the triangle formed by  $C_1$  and two other lines which are at angle of  $45^\circ$  with  $C_1$  and tangent to  $C_2$  is :

- (a) (1, 1)      (b) (0, 0)      (c) (-1, -1)      (d) (2, 2)

10. If  $S_1, S_2$  and  $S_3$  are three circles congruent to  $C_2$  and touch both  $C_1$  and  $C_2$ ; then the area of triangle formed by joining centres of the circles  $S_1, S_2$  and  $S_3$  is (in square units)

- (a) 2      (b) 4      (c) 8      (d) 16

**Answers**

1.	(d)	2.	(b)	3.	(c)	4.	(b)	5.	(c)	6.	(c)	7.	(a)	8.	(d)	9.	(b)	10.	(c)
----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	-----	-----




**Exercise-4 : Matching Type Problems**

1.

Column-I		Column-II	
(A)	The triangle $PQR$ is inscribed in the circle $x^2 + y^2 = 169$ . If $Q(5, 12)$ and $R(-12, 5)$ then $\angle QPR$ is	(P)	$\pi/6$
(B)	The angle between the lines joining the origin to the points of intersection of the line $4x + 3y = 24$ with circle $(x - 3)^2 + (y - 4)^2 = 25$	(Q)	$\pi/4$
(C)	Two parallel tangents drawn to given circle are cut by a third tangent. The angle subtended by the portion of third tangent between the given tangents at the centre is	(R)	$\pi/3$
(D)	A chord is drawn joining the point of contact of tangents drawn from a point $P$ to the circle. If the chord subtends an angle $\pi/2$ at the centre then the angle included between the tangents at $P$ is	(S)	$\pi/2$
		(T)	$\pi$

2.

Column-I		Column-II	
(A)	A ray of light coming from the point $(1, 2)$ is reflected at a point $A$ on the $x$ -axis then passes through the point $(5, 3)$ . The coordinates of the point $A$ are :	(P)	$\left(\frac{13}{5}, 0\right)$
(B)	The equation of three sides of triangle $ABC$ are $x + y = 3$ , $x - y = 5$ and $3x + y = 4$ . Considering the sides as diameter, three circles $S_1, S_2, S_3$ are drawn whose radical centre is at :	(Q)	$(4, -1)$
(C)	If the straight line $x - 2y + 1 = 0$ intersects the circle $x^2 + y^2 = 25$ at the points $P$ and $Q$ , then the coordinate of the point of intersection of tangents drawn at $P$ and $Q$ to the circle is	(R)	$(-25, 50)$
(D)	The equation of three sides of a triangle are $4x + 3y + 9 = 0$ , $2x + 3 = 0$ and $3y - 4 = 0$ . The circumcentre of the triangle is :	(S)	$\left(\frac{-19}{8}, \frac{1}{6}\right)$
		(T)	$(-1, 2)$

**Answers**
1. A  $\rightarrow$  Q; B  $\rightarrow$  S; C  $\rightarrow$  S; D  $\rightarrow$  S2. A  $\rightarrow$  P; B  $\rightarrow$  Q; C  $\rightarrow$  R; D  $\rightarrow$  S

### Exercise-5 : Subjective Type Problems


1. Tangents are drawn to circle  $x^2 + y^2 = 1$  at its intersection points (distinct) with the circle  $x^2 + y^2 + (\lambda - 3)x + (2\lambda + 2)y + 2 = 0$ . The locus of intersection of tangents is a straight line, then the slope of that straight line is.
2. The radical centre of the three circles is at the origin. The equations of the two of the circles are  $x^2 + y^2 = 1$  and  $x^2 + y^2 + 4x + 4y - 1 = 0$ . If the third circle passes through the points (1, 1) and (-2, 1); and its radius can be expressed in the form of  $\frac{p}{q}$ , where  $p$  and  $q$  are relatively prime positive integers. Find the value of  $(p + q)$ .
3. Let  $S = \{(x, y) \mid x, y \in \mathbb{R}, x^2 + y^2 - 10x + 16 = 0\}$ . The largest value of  $\frac{y}{x}$  can be put in the form  $\frac{m}{n}$  where  $m, n$  are relatively prime natural numbers, then  $m^2 + n^2 =$
4. In the above problem, the complete range of the expression  $x^2 + y^2 - 26x + 12y + 210$  is  $[a, b]$ , then  $b - 2a =$
5. If the line  $y = 2 - x$  is tangent to the circle  $S$  at the point  $P(1, 1)$  and circle  $S$  is orthogonal to the circle  $x^2 + y^2 + 2x + 2y - 2 = 0$ , then find the length of tangent drawn from the point (2, 2) to circle  $S$ .
6. Two circles having radii  $r_1$  and  $r_2$  passing through vertex  $A$  of a triangle  $ABC$ . One of the circle touches the side  $BC$  at  $B$  and other circle touches the side  $BC$  at  $C$ . If  $a = 5$  and  $A = 30^\circ$ , find  $\sqrt{r_1 r_2}$ .
7. A circle  $S$  of radius 'a' is the director circle of another circle  $S_1$ .  $S_1$  is the director circle of  $S_2$  and so on. If the sum of radius of  $S, S_1, S_2, S_3, \dots$  circles is '2' and  $a = (k - \sqrt{k})$ , then the value of  $k$  is .....
8. If  $r_1$  and  $r_2$  be the maximum and minimum radius of the circle which pass through the point (4, 3) and touch the circle  $x^2 + y^2 = 49$ , then  $\frac{r_1}{r_2}$  is .....
9. Let  $C$  be the circle  $x^2 + y^2 - 4x - 4y - 1 = 0$ . The number of points common to  $C$  and the sides of the rectangle determined by the lines  $x = 2, x = 5, y = -1$  and  $y = 5$  is  $P$  then find  $P$ .
10. Two congruent circles with centres at (2, 3) and (5, 6) intersects at right angle; find the radius of the circle.
11. The sum of abscissa and ordinate of a point on the circle  $x^2 + y^2 - 4x + 2y - 20 = 0$  which is nearest to  $\left(2, \frac{3}{2}\right)$  is :
12.  $AB$  is any chord of the circle  $x^2 + y^2 - 6x - 8y - 11 = 0$  which subtends an angle  $\frac{\pi}{2}$  at (1, 2). If locus of midpoint of  $AB$  is a circle  $x^2 + y^2 - 2ax - 2by - c = 0$ ; then find the value of  $(a + b + c)$ .

13. If circles  $x^2 + y^2 = c$  with radius  $\sqrt{3}$  and  $x^2 + y^2 + ax + by + c = 0$  with radius  $\sqrt{6}$  intersect at two points  $A$  and  $B$ . If length of  $AB = \sqrt{l}$ . Find  $l$ .

## Answers

1.	2	2.	5	3.	25	4.	66	5.	2	6.	5	7.	2
8.	6	9.	3	10.	3	11.	6	12.	8	13.	8		




**Exercise-1 : Single Choice Problems**

- Let  $PQ$  be the latus rectum of the parabola  $y^2 = 4x$  with vertex  $A$ . Minimum length of the projection of  $PQ$  on a tangent drawn in portion of parabola  $PAQ$  is :
  - 2
  - $2\sqrt{3}$
  - 4
  - $2\sqrt{2}$
- A normal is drawn to the parabola  $y^2 = 9x$  at the point  $P(4, 6)$ . A circle is described on  $SP$  as diameter; where  $S$  is the focus. The length of the intercept made by the circle on the normal at point  $P$  is :
  - $\frac{17}{4}$
  - $\frac{15}{4}$
  - 4
  - 5
- A trapezium is inscribed in the parabola  $y^2 = 4x$ , such that its diagonal pass through the point  $(1, 0)$  and each has length  $\frac{25}{4}$ . If the area of the trapezium be  $P$ , then  $4P$  is equal to :
  - 70
  - 71
  - 80
  - 75
- The length of normal chord of parabola  $y^2 = 4x$ , which subtends an angle of  $90^\circ$  at the vertex is :
  - $6\sqrt{3}$
  - $7\sqrt{2}$
  - $8\sqrt{2}$
  - $9\sqrt{2}$
- If  $b$  and  $c$  are the lengths of the segments of any focal chord of a parabola  $y^2 = 4ax$ . Then the length of semi-latus rectum is :
  - $\frac{bc}{b+c}$
  - $\frac{2bc}{b+c}$
  - $\frac{b+c}{2}$
  - $\sqrt{bc}$
- The length of the shortest path that begins at the point  $(-1, 1)$ , touches the  $x$ -axis and then ends at a point on the parabola  $(x-y)^2 = 2(x+y-4)$ , is :
  - $3\sqrt{2}$
  - 5
  - $4\sqrt{10}$
  - 13

7. If the normals at three points  $P, Q, R$  of the parabola  $y^2 = 4ax$  meet in a point  $O'$  and  $S$  be its focus, then  $|SP| \cdot |SQ| \cdot |SR|$  is equal to :
- (a)  $a^3$  (b)  $a^2(SO')$   
 (c)  $a(SO')^2$  (d) None of these
8. Let  $P$  and  $Q$  are points on the parabola  $y^2 = 4ax$  with vertex  $O$ , such that  $OP$  is perpendicular to  $OQ$  and have lengths  $r_1$  and  $r_2$  respectively, then the value of  $\frac{r_1^{4/3} r_2^{4/3}}{r_1^{2/3} + r_2^{2/3}}$  is :
- (a)  $16a^2$  (b)  $a^2$  (c)  $4a$  (d) None of these
9. Length of the shortest chord of the parabola  $y^2 = 4x + 8$ , which belongs to the family of lines  $(1 + \lambda)y + (\lambda - 1)x + 2(1 - \lambda) = 0$ , is :
- (a) 6 (b) 5 (c) 8 (d) 2
10. If locus of mid-point of any normal chord of the parabola :  
 $y^2 = 4x$  is  $x - a = \frac{b}{y^2} + \frac{y^2}{c}$ ;
- where  $a, b, c \in N$ , then  $(a + b + c)$  equals to :
- (a) 5 (b) 8 (c) 10 (d) None of these
11. Let tangents at  $P$  and  $Q$  to curve  $y^2 - 4x - 2y + 5 = 0$  intersect at  $T$ . If  $S(2, 1)$  is a point such that  $(SP)(SQ) = 16$ , then the length  $ST$  is equal to :
- (a) 3 (b) 4 (c) 5 (d) None of these
12. Abscissa of two points  $P$  and  $Q$  on parabola  $y^2 = 8x$  are roots of equation  $x^2 - 17x + 11 = 0$ . Let Tangents at  $P$  and  $Q$  meet at point  $T$ , then distance of  $T$  from the focus of parabola is :
- (a) 7 (b) 6 (c) 5 (d) 4
13. If  $Ax + By = 1$  is a normal to the curve  $ay = x^2$ , then :
- (a)  $4A^2(1 - aB) = aB^3$  (b)  $4A^2(2 + aB) = aB^3$   
 (c)  $4A^2(1 + aB) + aB^3 = 0$  (d)  $2A^2(2 - aB) = aB^3$
14. The equation of a curve which passes through the point  $(3, 1)$ , such that the segment of any tangent between the point of tangency and the  $x$ -axis is bisected at its point of intersection with  $y$ -axis, is :
- (a)  $x = 3y^2$  (b)  $x^2 = 9y$  (c)  $x = y^2 + 2$  (d)  $2x = 3y^2 + 3$
15. The parabola  $y = 4 - x^2$  has vertex  $P$ . It intersects  $x$ -axis at  $A$  and  $B$ . If the parabola is translated from its initial position to a new position by moving its vertex along the line  $y = x + 4$ , so that it intersects  $x$ -axis at  $B$  and  $C$ , then abscissa of  $C$  will be :
- (a) 3 (b) 4 (c) 6 (d) 8

16. A focal chord for parabola  $y^2 = 8(x + 2)$  is inclined at an angle of  $60^\circ$  with positive  $x$ -axis and intersects the parabola at  $P$  and  $Q$ . Let perpendicular bisector of the chord  $PQ$  intersects the  $x$ -axis at  $R$ ; then the distance of  $R$  from focus is :
- (a)  $\frac{8}{3}$                       (b)  $\frac{16\sqrt{3}}{3}$                       (c)  $\frac{16}{3}$                       (d)  $8\sqrt{3}$
17. The Director circle of the parabola  $(y - 2)^2 = 16(x + 7)$  touches the circle  $(x - 1)^2 + (y + 1)^2 = r^2$ , then  $r$  is equal to :
- (a) 10                      (b) 11                      (c) 12                      (d) None of these
18. The chord of contact of a point  $A(x_A, y_A)$  of  $y^2 = 4x$  passes through  $(3, 1)$  and point  $A$  lies on  $x^2 + y^2 = 5^2$ . Then :
- (a)  $5x_A^2 + 24x_A + 11 = 0$                       (b)  $13x_A^2 + 8x_A - 21 = 0$   
(c)  $5x_A^2 + 24x_A + 61 = 0$                       (d)  $13x_A^2 + 21x_A - 31 = 0$

### Answers

1. (d)	2. (b)	3. (d)	4. (a)	5. (b)	6. (a)	7. (c)	8. (a)	9. (c)	10. (b)
11. (b)	12. (a)	13. (d)	14. (a)	15. (d)	16. (c)	17. (c)	18. (a)		

**Exercise-2 : One or More than One Answer is/are Correct**


1.  $PQ$  is a double ordinate of the parabola  $y^2 = 4ax$ . If the normal at  $P$  intersect the line passing through  $Q$  and parallel to  $x$ -axis at  $G$ ; then locus of  $G$  is a parabola with :
- (a) vertex at  $(4a, 0)$                       (b) focus at  $(5a, 0)$   
(c) directrix as the line  $x - 3a = 0$       (d) length of latus rectum equal to  $4a$

**Answers**

1. (a, b, c, d)





 **Exercise-4 : Matching Type Problems**

1.

Column-I		Column-II	
(A)	The equation of tangent to the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ which cuts off equal intercepts on axes is $x - y = a$ where $ a $ equal to	(P)	$\sqrt{2}$
(B)	The normal $y = mx - 2am - am^2$ to the parabola $y^2 = 4ax$ subtends a right angle at the vertex if $ m $ equal to	(Q)	$\sqrt{3}$
(C)	The equation of the common tangent to parabola $y^2 = 4x$ and $x^2 = 4y$ is $x + y + \frac{k}{\sqrt{3}} = 0$ , then $k$ is equal to	(R)	$\sqrt{8}$
(D)	An equation of common tangent to parabola $y^2 = 8x$ and the hyperbola $3x^2 - y^2 = 3$ is $4x - 2y + \frac{k}{\sqrt{2}} = 0$ , then $k$ is equal to	(S)	$\sqrt{41}$
		(T)	2

2.

Column-I		Column-II	
(A)	Area of $\Delta PQR$ is equal to	(P)	2
(B)	Radius of circumcircle of $\Delta PQR$ is equal to	(Q)	$\frac{5}{2}$
(C)	Distance of the vertex from the centroid of $\Delta PQR$ is equal to	(R)	$\frac{3}{2}$
(D)	Distance of the centroid from the circumcentre of $\Delta PQR$ is equal to	(S)	$\frac{2}{3}$
		(T)	$\frac{11}{6}$

**Answers**
1. A  $\rightarrow$  S; B  $\rightarrow$  P; C  $\rightarrow$  Q; D  $\rightarrow$  R2. A  $\rightarrow$  P; B  $\rightarrow$  Q; C  $\rightarrow$  S; D  $\rightarrow$  T

### Exercise-5 : Subjective Type Problems

- Points  $A$  and  $B$  lie on the parabola  $y = 2x^2 + 4x - 2$ , such that origin is the mid-point of the line segment  $AB$ . If ' $l$ ' be the length of the line segment  $AB$ , then find the unit digit of  $l^2$ .
- For the parabola  $y = -x^2$ , let  $a < 0$  and  $b > 0$ ;  $P(a, -a^2)$  and  $Q(b, -b^2)$ . Let  $M$  be the mid-point of  $PQ$  and  $R$  be the point of intersection of the vertical line through  $M$ , with the parabola. If the ratio of the area of the region bounded by the parabola and the line segment  $PQ$  to the area of the triangle  $PQR$  be  $\frac{\lambda}{\mu}$ ; where  $\lambda$  and  $\mu$  are relatively prime positive integers, then find the value of  $(\lambda + \mu)$  :
- The chord  $AC$  of the parabola  $y^2 = 4ax$  subtends an angle of  $90^\circ$  at points  $B$  and  $D$  on the parabola. If points  $A, B, C$  and  $D$  are represented by  $(at_i^2, 2at_i)$ ,  $i = 1, 2, 3, 4$  respectively, then find the value of  $\left| \frac{t_2 + t_4}{t_1 + t_3} \right|$ .

### Answers

1.	8	2.	7	3.	1														
----	---	----	---	----	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--



### Exercise-1 : Single Choice Problems

1. If  $CF$  be the perpendicular from the centre  $C$  of the ellipse  $\frac{x^2}{12} + \frac{y^2}{8} = 1$ , on the tangent at any point  $P$  and  $G$  is the point where the normal at  $P$  meets the major axis, then the value of  $(CF \cdot PG)$  equals to :  
 (a) 5 (b) 6 (c) 8 (d) None of these
2. The minimum length of intercept on any tangent to the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$  cut by the circle  $x^2 + y^2 = 25$  is :  
 (a) 8 (b) 9 (c) 2 (d) 11
3. The point on the ellipse  $x^2 + 2y^2 = 6$ , whose distance from the line  $x + y = 7$  is minimum is :  
 (a) (2, 3) (b) (2, 1) (c) (1, 0) (d) None of these
4. If lines  $2x + 3y = 10$  and  $2x - 3y = 10$  are tangents at the extremities of a latus rectum of an ellipse; whose centre is origin, then the length of the latus rectum is :  
 (a)  $\frac{110}{27}$  (b)  $\frac{98}{27}$  (c)  $\frac{100}{27}$  (d)  $\frac{120}{27}$
5. The area bounded by the circle  $x^2 + y^2 = a^2$  and the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is equal to the area of another ellipse having semi-axes :  
 (a)  $a + b$  and  $b$  (b)  $a - b$  and  $a$  (c)  $a$  and  $b$  (d) None of these
6. If  $F_1$  and  $F_2$  are the feet of the perpendiculars from foci  $S_1$  and  $S_2$  of the ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1$  on the tangent at any point  $P$  of the ellipse, then :  
 (a)  $S_1F_1 + S_2F_2 \geq 2$  (b)  $S_1F_1 + S_2F_2 \geq 3$  (c)  $S_1F_1 + S_2F_2 \geq 6$  (d)  $S_1F_1 + S_2F_2 \geq 8$

7. Consider the ellipse  $\frac{x^2}{f(k^2 + 2k + 5)} + \frac{y^2}{f(k + 11)} = 1$ , where  $f(x)$  is a positive decreasing function, then the value of  $k$  for which major axis coincides with  $x$ -axis is :
- (a)  $k \in (-7, -5)$       (b)  $k \in (-5, -3)$       (c)  $k \in (-3, 2)$       (d) None of these
8. If area of the ellipse  $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$  inscribed in a square of side length  $5\sqrt{2}$  is  $A$ , then  $\frac{A}{\pi}$  equals to :
- (a) 12      (b) 10      (c) 8      (d) 11
9. Any chord of the conic  $x^2 + y^2 + xy = 1$  passing through origin is bisected at a point  $(p, q)$ , then  $(p + q + 12)$  equals to :
- (a) 13      (b) 14      (c) 11      (d) 12
10. Tangents are drawn from the point  $(4, 2)$  to the curve  $x^2 + 9y^2 = 9$ , the tangent of angle between the tangents :
- (a)  $\frac{3\sqrt{3}}{5\sqrt{17}}$       (b)  $\frac{\sqrt{43}}{10}$       (c)  $\frac{\sqrt{43}}{5}$       (d)  $\sqrt{\frac{3}{17}}$

### Answers

1.	(c)	2.	(a)	3.	(b)	4.	(c)	5.	(b)	6.	(d)	7.	(c)	8.	(a)	9.	(d)	10.	(c)
----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----	-----	-----	-----




**Exercise-3 : Matching Type Problems**

1.

Column-I		Column-II	
(A)	If the tangent to the ellipse $x^2 + 4y^2 = 16$ at the point $P(4 \cos \phi, 2 \sin \phi)$ is a normal to the circle $x^2 + y^2 - 8x - 4y = 0$ then $\frac{\phi}{2}$ may be	(P)	0
(B)	The eccentric angle(s) of a point on the ellipse $x^2 + 3y^2 = 6$ at a distance 2 units from the centre of the ellipse is/are	(Q)	$\cos^{-1}\left(-\frac{2}{3}\right)$
(C)	The eccentric angle of point of intersection of the ellipse $x^2 + 4y^2 = 4$ and the parabola $x^2 + 1 = y$ is	(R)	$\frac{\pi}{4}$
(D)	If the normal at the point $P(\sqrt{14} \cos \theta, \sqrt{5} \sin \theta)$ to the ellipse $\frac{x^2}{14} + \frac{y^2}{5} = 1$ intersect it again at the point $Q(\sqrt{14} \cos 2\theta, \sqrt{5} \sin 2\theta)$ , then $\theta$ is	(S)	$\frac{5\pi}{4}$
		(T)	$\frac{\pi}{2}$

**Answers**
**1.** A  $\rightarrow$  P, R; B  $\rightarrow$  R, S; C  $\rightarrow$  P; D  $\rightarrow$  Q


**Exercise-4 : Subjective Type Problems**

1. For the ellipse  $\frac{x^2}{9} + \frac{y^2}{4} = 1$ . Let  $O$  be the centre and  $S$  and  $S'$  be the foci. For any point  $P$  on the ellipse the value of  $PS \cdot PS'd^2$  (where  $d$  is the distance of  $O$  from the tangent at  $P$ ) is equal to
2. Number of perpendicular tangents that can be drawn on the ellipse  $\frac{x^2}{16} + \frac{y^2}{25} = 1$  from point  $(6, 7)$  is

**Answers**

1.	4	2.	0									
----	---	----	---	--	--	--	--	--	--	--	--	--



 **Exercise-1 : Single Choice Problems**

- The normal to curve  $xy = 4$  at the point  $(1, 4)$  meets the curve again at :  
 (a)  $(-4, -1)$       (b)  $(-8, -\frac{1}{2})$       (c)  $(-16, -\frac{1}{4})$       (d)  $(-1, -4)$
- Let  $PQ : 2x + y + 6 = 0$  is a chord of the curve  $x^2 - 4y^2 = 4$ . Coordinates of the point  $R(\alpha, \beta)$  that satisfy  $\alpha^2 + \beta^2 - 1 \leq 0$ ; such that area of triangle  $PQR$  is minimum; are given by :  
 (a)  $(\frac{-2}{\sqrt{5}}, \frac{1}{\sqrt{5}})$       (b)  $(\frac{-2}{\sqrt{5}}, \frac{-1}{\sqrt{5}})$   
 (c)  $(\frac{2}{\sqrt{5}}, \frac{1}{\sqrt{5}})$       (d)  $(\frac{2}{\sqrt{5}}, \frac{-1}{\sqrt{5}})$
- If  $y = mx + c$  be a tangent to hyperbola  $\frac{x^2}{\lambda^2} - \frac{y^2}{(\lambda^3 + \lambda^2 + \lambda)^2} = 1$ , then least value of  $16m^2$  equals to :  
 (a) 0      (b) 1      (c) 4      (d) 9
- Let the double ordinate  $PP'$  of the hyperbola  $\frac{x^2}{4} - \frac{y^2}{3} = 1$  is produced both sides to meet asymptotes of hyperbola in  $Q$  and  $Q'$ . The product  $(PQ)(PQ')$  is equal to :  
 (a) 3      (b) 4      (c) 1      (d) 5
- If eccentricity of conjugate hyperbola of the given hyperbola :  

$$|\sqrt{(x-1)^2 + (y-2)^2} - \sqrt{(x-5)^2 + (y-5)^2}| = 3$$
 is  $e'$ , then value of  $8e'$  is :  
 (a) 12      (b) 14      (c) 17      (d) 10



6. A normal to the hyperbola  $\frac{x^2}{4} - \frac{y^2}{1} = 1$  has equal intercepts on positive  $x$  and positive  $y$ -axes. If this normal touches the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , then  $3(a^2 + b^2)$  is equal to :
- (a) 5                      (b) 25                      (c) 16                      (d) None of these
7. Locus of a point, whose chord of contact with respect to the circle  $x^2 + y^2 = 4$  is a tangent to the hyperbola  $xy = 1$  is a/an :
- (a) ellipse                      (b) circle  
(c) hyperbola                      (d) parabola
8. Let the chord  $x \cos \alpha + y \sin \alpha = p$  of the hyperbola  $\frac{x^2}{16} - \frac{y^2}{18} = 1$  subtends a right angle at the centre. Let diameter of the circle, concentric with the hyperbola, to which the given chord is a tangent is  $d$ , then  $\frac{d}{4}$  is equal to :
- (a) 4                      (b) 5                      (c) 6                      (d) 7
9. If the tangent and normal at a point on rectangular hyperbola cut-off intercept  $a_1, a_2$  on  $x$ -axis and  $b_1, b_2$  on the  $y$ -axis, then  $a_1 a_2 + b_1 b_2$  is equal to :
- (a) 2                      (b)  $\frac{1}{2}$                       (c) 0                      (d) -1

### Answers

1. (c)	2. (b)	3. (d)	4. (a)	5. (d)	6. (b)	7. (c)	8. (c)	9. (c)	
--------	--------	--------	--------	--------	--------	--------	--------	--------	--

**Exercise-2 : One or More than One Answer is/are Correct**

1. A common tangent to the hyperbola  $9x^2 - 16y^2 = 144$  and the circle  $x^2 + y^2 = 9$  is/are :
- (a)  $y = \frac{3}{\sqrt{7}}x + \frac{15}{\sqrt{7}}$  (b)  $y = 3\sqrt{\frac{2}{7}}x + \frac{25}{\sqrt{7}}$   
 (c)  $y = 2\sqrt{\frac{3}{7}}x + 15\sqrt{7}$  (d)  $y = -3\sqrt{\frac{2}{7}}x + \frac{25}{\sqrt{7}}$
2. Tangents are drawn to the hyperbola  $x^2 - y^2 = 3$  which are parallel to the line  $2x + y + 8 = 0$ . Then their points of contact is/are :
- (a) (2, 1) (b) (2, -1)  
 (c) (-2, -1) (d) (-2, 1)
3. If the line  $ax + by + c = 0$  is normal to the curve  $xy = 1$ , then:
- (a)  $a > 0, b > 0$  (b)  $a > 0, b < 0$   
 (c)  $b < 0, a < 0$  (d)  $a < 0, b > 0$
4. A circle cuts rectangular hyperbola  $xy = 1$  in the points  $(x_r, y_r)$ ,  $r = 1, 2, 3, 4$  then :
- (a)  $y_1y_2y_3y_4 = 1$  (b)  $x_1x_2x_3x_4 = 1$   
 (c)  $x_1x_2x_3x_4 = y_1y_2y_3y_4 = -1$  (d)  $y_1y_2y_3y_4 = 0$

**Answers**

1.	(b, d)	2.	(b, d)	3.	(b, d)	4.	(a, b)		
----	--------	----	--------	----	--------	----	--------	--	--



**Exercise-4 : Subjective Type Problems**

1. Let  $y = mx + c$  be a common tangent to  $\frac{x^2}{16} - \frac{y^2}{9} = 1$  and  $\frac{x^2}{4} + \frac{y^2}{3} = 1$ , then find the value of  $m^2 + c^2$ .
2. The maximum number of normals that can be drawn to an ellipse/hyperbola passing through a given point is :
3. Tangent at  $P$  to rectangular hyperbola  $xy = 2$  meets coordinate axes at  $A$  and  $B$ , then area of triangle  $OAB$  (where  $O$  is origin) is :

**Answers**

1.	8	2.	4	3.	4								
----	---	----	---	----	---	--	--	--	--	--	--	--	--

# Trigonometry

---


- 22.** Compound Angles
- 23.** Trigonometric Equations
- 24.** Solution of Triangles
- 25.** Inverse Trigonometric Functions

Chapter 22 – Compound Angles

**Exercise-1 : Single Choice Problems**

- $\left(\cos^4 \frac{\pi}{24} - \sin^4 \frac{\pi}{24}\right)$  equals :
  - $\frac{1}{\sqrt{2}}$
  - $\frac{\sqrt{6}-\sqrt{2}}{4}$
  - $\frac{\sqrt{6}+\sqrt{2}}{4}$
  - $\frac{\sqrt{3}+1}{2}$
- If a  $\sin x + b \cos(c+x) + b \cos(c-x) = \alpha$ ,  $\alpha > a$ , then the minimum value of  $|\cos c|$  is :
  - $\sqrt{\frac{\alpha^2 - a^2}{b^2}}$
  - $\sqrt{\frac{\alpha^2 - a^2}{2b^2}}$
  - $\sqrt{\frac{\alpha^2 - a^2}{3b^2}}$
  - $\sqrt{\frac{\alpha^2 - a^2}{4b^2}}$
- If all values of  $x \in (a, b)$  satisfy the inequality  $\tan x \tan 3x < -1$ ,  $x \in \left(0, \frac{\pi}{2}\right)$ , then the maximum value  $(b-a)$  is :
  - $\frac{\pi}{12}$
  - $\frac{\pi}{3}$
  - $\frac{\pi}{6}$
  - $\frac{\pi}{4}$
- $\sum_{r=1}^8 \tan(rA) \tan((r+1)A)$  where  $A = 36^\circ$  is :
  - $-10 - \tan A$
  - $-10 + \tan A$
  - $-10$
  - $-9$
- Let  $f(x) = 2 \operatorname{cosec} 2x + \sec x + \operatorname{cosec} x$ , then minimum value of  $f(x)$  for  $x \in \left(0, \frac{\pi}{2}\right)$  is :
  - $\frac{1}{\sqrt{2}-1}$
  - $\frac{2}{\sqrt{2}-1}$
  - $\frac{1}{\sqrt{2}+1}$
  - $\frac{2}{\sqrt{2}+1}$
- The exact value of  $\operatorname{cosec} 10^\circ + \operatorname{cosec} 50^\circ - \operatorname{cosec} 70^\circ$  is :
  - 4
  - 5
  - 6
  - 8
- If  $u = \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} + \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$ , then the difference between the maximum and minimum values of  $u^2$  is given by :
  - $2(a^2 + b^2)$
  - $2\sqrt{a^2 + b^2}$
  - $(a+b)^2$
  - $(a-b)^2$

8. If  $u_n = \sin(n\theta) \sec^n \theta$ ,  $v_n = \cos(n\theta) \sec^n \theta$ ,  $n \in N$ ,  $n \neq 1$ , then  $\frac{v_n - v_{n-1}}{u_{n-1}} + \frac{1}{n} \frac{u_n}{v_n} =$
- (a)  $-\cot \theta + \frac{1}{n} \tan(n\theta)$  (b)  $\cot \theta + \frac{1}{n} \tan(n\theta)$   
 (c)  $\tan \theta + \frac{1}{n} \tan(n\theta)$  (d)  $-\tan \theta + \frac{\tan(n\theta)}{n}$
9. If  $a \cos^2 3\alpha + b \cos^4 \alpha = 16 \cos^6 \alpha + 9 \cos^2 \alpha$  is an identity, then  
 (a)  $a = 1, b = 24$  (b)  $a = 3, b = 24$  (c)  $a = 4, b = 2$  (d)  $a = 7, b = 18$
10. Maximum value of  $\cos x (\sin x + \cos x)$  is equal to :  
 (a)  $\sqrt{2}$  (b) 2 (c)  $\frac{\sqrt{2}+1}{2}$  (d)  $\sqrt{2}+1$
11. If  $\frac{\sin A}{\sin B} = \frac{\sqrt{3}}{2}$  and  $\frac{\cos A}{\cos B} = \frac{\sqrt{5}}{2}$ ,  $0 < A, B < \frac{\pi}{2}$  then  $\tan A + \tan B$  is equal to :  
 (a)  $\sqrt{\frac{3}{5}}$  (b)  $\sqrt{\frac{5}{3}}$  (c)  $\frac{\sqrt{3}+\sqrt{5}}{\sqrt{5}}$  (d)  $\frac{\sqrt{3}+\sqrt{5}}{\sqrt{3}}$
12. Let  $0 \leq \alpha, \beta, \gamma, \delta \leq \pi$  where  $\beta$  and  $\gamma$  are not complementary such that  
 $2 \cos \alpha + 6 \cos \beta + 7 \cos \gamma + 9 \cos \delta = 0$   
 and  $2 \sin \alpha - 6 \sin \beta + 7 \sin \gamma - 9 \sin \delta = 0$   
 If  $\frac{\cos(\alpha + \delta)}{\cos(\beta + \gamma)} = \frac{m}{n}$  where  $m$  and  $n$  are relatively prime positive numbers, then the value of  $(m + n)$  is equal to :  
 (a) 11 (b) 10 (c) 9 (d) 7
13. If  $-\pi < \theta < -\frac{\pi}{2}$ , then  $\left| \sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} + \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}} \right|$  is equal to :  
 (a)  $2 \sec \theta$  (b)  $-2 \sec \theta$  (c)  $2 \sec \frac{\theta}{2}$  (d)  $-\sec \frac{\theta}{2}$
14. If  $A = \sum_{r=1}^3 \cos \frac{2r\pi}{7}$  and  $B = \sum_{r=1}^3 \cos \frac{2^r \pi}{7}$ , then :  
 (a)  $A + B = 0$  (b)  $2A + B = 0$  (c)  $A + 2B = 0$  (d)  $A = B$
15. In a  $\Delta PQR$  (as shown in figure) if  $x : y : z = 2 : 3 : 6$ , then the value of  $\angle QPR$  is :  
 (a)  $\frac{\pi}{6}$  (b)  $\frac{\pi}{4}$   
 (c)  $\frac{\pi}{3}$  (d)  $\frac{\pi}{2}$



16. If  $A = \sum_{r=1}^3 \cos \frac{2r\pi}{7}$  and  $B = \sum_{r=1}^3 \cos \frac{2^r \pi}{7}$ , then :
- (a)  $A + B = 0$       (b)  $2A + B = 0$       (c)  $A + 2B = 0$       (d)  $A - B = 0$
17. Let  $f(x) = \sin x + 2 \cos^2 x$ ;  $\frac{\pi}{6} \leq x \leq \frac{2\pi}{3}$ , then maximum value of  $f(x)$  is :
- (a) 1      (b)  $\frac{3}{2}$       (c) 2      (d)  $\frac{5}{2}$
18. In  $\triangle ABC$ ,  $\angle C = \frac{2\pi}{3}$  then the value of  $\cos^2 A + \cos^2 B - \cos A \cdot \cos B$  is equal to :
- (a)  $\frac{3}{4}$       (b)  $\frac{3}{2}$       (c)  $\frac{1}{2}$       (d)  $\frac{1}{4}$
19. The number of solutions of the equation  $4 \sin^2 x + \tan^2 x + \cot^2 x + \operatorname{cosec}^2 x = 6$  in  $[0, 2\pi]$  :
- (a) 1      (b) 2      (c) 3      (d) 4
20. If  $\sin A$ ,  $\cos A$  and  $\tan A$  are in G.P., then  $\cos^3 A + \cos^2 A$  is equal to :
- (a) 1      (b) 2      (c) 4      (d) none
21. Range of function  $f(x) = \sin\left(x + \frac{\pi}{6}\right) + \cos\left(x - \frac{\pi}{6}\right)$  is :
- (a)  $[-\sqrt{2}, \sqrt{2}]$       (b)  $[-\sqrt{2}(\sqrt{3} + 1), \sqrt{2}(\sqrt{3} + 1)]$   
 (c)  $\left[-\frac{\sqrt{3} + 1}{\sqrt{2}}, \frac{\sqrt{3} + 1}{\sqrt{2}}\right]$       (d)  $\left[-\frac{\sqrt{3} - 1}{\sqrt{2}}, \frac{\sqrt{3} - 1}{\sqrt{2}}\right]$
22. The value of  $\tan(\log_2 6) \cdot \tan(\log_2 3) \cdot \tan 1$  is always equal to :
- (a)  $\tan(\log_2 6) + \tan(\log_2 3) + \tan 1$       (b)  $\tan(\log_2 6) - \tan(\log_2 3) - \tan 1$   
 (c)  $\tan(\log_2 6) - \tan(\log_2 3) + \tan 1$       (d)  $\tan(\log_2 6) + \tan(\log_2 3) - \tan 1$
23. In a triangle  $ABC$ , side  $BC = 3$ ,  $AC = 4$  and  $AB = 5$ . The value of  $\sin A + \sin 2B + \sin 3C$  is equal to:
- (a)  $\frac{24}{25}$       (b)  $\frac{14}{25}$       (c)  $\frac{64}{25}$       (d) none
24. If  $A + B + C = 180^\circ$ , then  $\frac{\cos A \cos C + \cos(A + B) \cos(B + C)}{\cos A \sin C - \sin(A + B) \cos(B + C)}$  simplifies to :
- (a)  $-\cot C$       (b) 0      (c)  $\tan C$       (d)  $\cot C$
25. If  $\alpha + \gamma = 2\beta$  then the expression  $\frac{\sin \alpha - \sin \gamma}{\cos \gamma - \cos \alpha}$  simplifies to :
- (a)  $\tan \beta$       (b)  $-\tan \beta$       (c)  $\cot \beta$       (d)  $-\cot \beta$



26. The product  $\left(\cos \frac{x}{2}\right) \cdot \left(\cos \frac{x}{4}\right) \cdot \left(\cos \frac{x}{8}\right) \cdots \left(\cos \frac{x}{256}\right)$  is equal to :
- (a)  $\frac{\sin x}{128 \sin \frac{x}{256}}$  (b)  $\frac{\sin x}{256 \sin \frac{x}{256}}$  (c)  $\frac{\sin x}{128 \sin \frac{x}{128}}$  (d)  $\frac{\sin x}{512 \sin \frac{x}{512}}$
27. The value of the expression  $\frac{\sin 7\alpha + 6 \sin 5\alpha + 17 \sin 3\alpha + 12 \sin \alpha}{\sin 6\alpha + 5 \sin 4\alpha + 12 \sin 2\alpha}$ , where  $\alpha = \frac{\pi}{5}$  is equal to :
- (a)  $\frac{\sqrt{5}-1}{4}$  (b)  $\frac{\sqrt{5}+1}{4}$  (c)  $\frac{\sqrt{5}+1}{2}$  (d)  $\frac{\sqrt{5}-1}{2}$
28. In a triangle  $ABC$  if  $\sum \tan^2 A = \sum \tan A \tan B$ , then largest angle of the triangle in radian will be :
- (a)  $\frac{2\pi}{3}$  (b)  $\frac{\pi}{3}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{3\pi}{4}$
29. Which one of the following values is not the solution of the equation  $\log_{|\sin x|}(|\cos x|) + \log_{|\cos x|}(|\sin x|) = 2$
- (a)  $\frac{7\pi}{4}$  (b)  $\frac{11\pi}{4}$  (c)  $\frac{3\pi}{4}$  (d)  $\frac{3\pi}{8}$
30. Range of  $f(x) = \sin^6 x + \cos^6 x$  is :
- (a)  $\left[\frac{1}{4}, 1\right]$  (b)  $\left[\frac{1}{4}, \frac{3}{4}\right]$  (c)  $\left[\frac{3}{4}, 1\right]$  (d)  $[1, 2]$
31. If  $y = \frac{2 \sin \alpha}{1 + \cos \alpha + \sin \alpha}$ , then  $\frac{1 - \cos \alpha + \sin \alpha}{1 + \sin \alpha}$  is equal to :
- (a)  $\frac{1}{y}$  (b)  $y$  (c)  $1 - y$  (d)  $1 + y$
32. If  $\frac{\tan^3 A}{1 + \tan^2 A} + \frac{\cot^3 A}{1 + \cot^2 A} = p \sec A \operatorname{cosec} A + q \sin A \cos A$ , then :
- (a)  $p = 2, q = 1$  (b)  $p = 1, q = 2$  (c)  $p = 1, q = -2$  (d)  $p = 2, q = -1$
33. If  $\theta$  lies in the second quadrant. Then the value of  $\sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} + \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}}$  is equal to :
- (a)  $2 \sec \theta$  (b)  $-2 \sec \theta$  (c)  $2 \operatorname{cosec} \theta$  (d) 2
34. If  $y = (\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2$ , then minimum value of  $y$  is :
- (a) 7 (b) 8 (c) 9 (d) none of these
35. If  $\log_3 \sin x - \log_3 \cos x - \log_3 (1 - \tan x) - \log_3 (1 + \tan x) = -1$ , then  $\tan 2x$  is equal to (wherever defined)
- (a) -2 (b)  $\frac{3}{2}$  (c)  $\frac{2}{3}$  (d) 6

36. If  $\sin \theta + \operatorname{cosec} \theta = 2$ , then the value of  $\sin^8 \theta + \operatorname{cosec}^8 \theta$  is equal to :  
 (a) 2 (b)  $2^4$  (c)  $2^8$  (d) more than  $2^8$
37. If  $\tan^3 \theta + \cot^3 \theta = 52$ , then the value of  $\tan^2 \theta + \cot^2 \theta$  is equal to :  
 (a) 14 (b) 15  
 (c) 16 (d) 17
38. The maximum value of  $\log_{20}(3 \sin x - 4 \cos x + 15)$  is equal to :  
 (a) 1 (b) 2 (c) 3 (d) 4
39. If  $x^2 + y^2 = 9$  and  $4a^2 + 9b^2 = 16$ , then maximum value of  $4a^2x^2 + 9b^2y^2 - 12abxy$  is :  
 (a) 81 (b) 100 (c) 121 (d) 144
40. If  $A = \sqrt{\sin 2 - \sin \sqrt{3}}$ ,  $B = \sqrt{\cos 2 - \cos \sqrt{3}}$ , then which of the following statement is true ?  
 (a)  $A$  and  $B$  both are real numbers and  $A > B$   
 (b)  $A$  and  $B$  both are real numbers and  $A < B$   
 (c) Exactly one of  $A$  and  $B$  is not real number  
 (d) Both  $A$  and  $B$  are not real numbers
41. The number of real values of  $x$  such that  
 $(2^x + 2^{-x} - 2 \cos x)(3^{x+\pi} + 3^{-x-\pi} + 2 \cos x)(5^{\pi-x} + 5^{x-\pi} - 2 \cos x) = 0$  is :  
 (a) 1 (b) 2 (c) 3 (d) infinite
42. The equation  $e^{\sin x} - e^{-\sin x} - 4 = 0$  has :  
 (a) infinite number of real roots (b) no real roots  
 (c) exactly one real root (d) exactly four real roots
43. If  $\pi < \alpha < \frac{3\pi}{2}$ , then the expression  $\sqrt{4 \sin^4 \alpha + \sin^2 2\alpha} + 4 \cos^2 \left( \frac{\pi - \alpha}{4} \right)$  is equal to :  
 (a)  $2 + 4 \sin \alpha$  (b)  $2 - 4 \cos \alpha$  (c) 2 (d)  $2 - 4 \sin \alpha$
44.  $\left( \cos \frac{\pi}{12} - \sin \frac{\pi}{12} \right) \left( \tan \frac{\pi}{12} + \cot \frac{\pi}{12} \right) =$   
 (a)  $\frac{1}{\sqrt{2}}$  (b)  $4\sqrt{2}$  (c)  $\sqrt{2}$  (d)  $2\sqrt{2}$
45.  $\tan(100^\circ) + \tan(125^\circ) + \tan(100^\circ) \tan(125^\circ) =$   
 (a) 0 (b)  $\frac{1}{2}$  (c) -1 (d) 1
46. If  $\sin x + \sin^2 x = 1$ , then  $\cos^8 x + 2 \cos^6 x + \cos^4 x =$   
 (a) 2 (b) 1 (c) 3 (d)  $\frac{1}{2}$
47. The maximum value of  $\log_5(3x + 4y)$ , if  $x^2 + y^2 = 25$  is :  
 (a) 1 (b) 2 (c) 3 (d) 4

48. The number of values of  $\theta$  between  $-\pi$  and  $\frac{3\pi}{2}$  that satisfies the equation  $5 \cos 2\theta + 2 \cos^2 \frac{\theta}{2} + 1 = 0$  is :
- (a) 3 (b) 4 (c) 5 (d) 6
49. Given that  $\sin \beta = \frac{4}{5}$ ,  $0 < \beta < \pi$  and  $\tan \beta > 0$ , then  $((3 \sin(\alpha + \beta) - 4 \cos(\alpha + \beta)) \operatorname{cosec} \alpha)$  is equal to:
- (a) 2 (b) 3 (c) 4 (d) 5
50. The maximum value of  $\sin\left(x + \frac{\pi}{6}\right) + \cos\left(x + \frac{\pi}{6}\right)$  for  $x \in \left[0, \frac{\pi}{2}\right]$  is attained at  $x =$
- (a)  $\frac{\pi}{12}$  (b)  $\frac{\pi}{6}$  (c)  $\frac{\pi}{3}$  (d)  $\frac{\pi}{2}$
51. The values of 'a' for which the equation  $\sin x (\sin x + \cos x) = a$  has a real solution are
- (a)  $1 - \sqrt{2} \leq a \leq 1 + \sqrt{2}$  (b)  $2 - \sqrt{3} \leq a \leq 2 + \sqrt{3}$   
(c)  $0 \leq a \leq 2 + \sqrt{3}$  (d)  $\frac{1 - \sqrt{2}}{2} \leq a \leq \frac{1 + \sqrt{2}}{2}$
52. The value of  $\cos 12^\circ \cos 24^\circ \cos 36^\circ \cos 48^\circ \cos 60^\circ \cos 72^\circ \cos 84^\circ$  is :
- (a)  $\frac{1}{64}$  (b)  $\frac{1}{128}$  (c)  $\frac{1}{256}$  (d)  $\frac{1}{512}$
53. The ratio of the maximum value to minimum value of  $2 \cos^2 \theta + \cos \theta + 1$  is :
- (a) 32 : 7 (b) 32 : 9 (c) 4 : 1 (d) 2 : 1
54. If all values of  $x \in (a, b)$  satisfy the inequality  $\tan x \tan 3x < -1$ ,  $x \in \left(0, \frac{\pi}{2}\right)$ , then the maximum value  $(b - a)$  is :
- (a)  $\frac{\pi}{12}$  (b)  $\frac{\pi}{3}$  (c)  $\frac{\pi}{6}$  (d)  $\frac{\pi}{4}$
55. If a regular polygon of 'n' sides has circum radius = R and inradius = r; then each side of polygon is :
- (a)  $(R + r) \tan\left(\frac{\pi}{2n}\right)$  (b)  $2(R + r) \tan\left(\frac{\pi}{2n}\right)$   
(c)  $(R + r) \sin\left(\frac{\pi}{2n}\right)$  (d)  $2(R + r) \cot\left(\frac{\pi}{2n}\right)$
56. The value of  $\cos 12^\circ + \cos 84^\circ + \cos 156^\circ + \cos 132^\circ$  is:
- (a)  $\frac{1}{8}$  (b)  $-\frac{1}{2}$  (c) 1 (d)  $\frac{1}{2}$
57.  $\frac{\sin \theta}{\cos(3\theta)} + \frac{\sin(3\theta)}{\cos(9\theta)} + \frac{\sin(9\theta)}{\cos(27\theta)} + \frac{\sin(27\theta)}{\cos(81\theta)} =$

- (a)  $\frac{\sin(81\theta)}{2\cos(80\theta)\cos\theta}$  (b)  $\frac{\sin(80\theta)}{2\cos(81\theta)\cos\theta}$   
 (c)  $\frac{\sin(81\theta)}{\cos(80\theta)\cos\theta}$  (d)  $\frac{\sin(80\theta)}{\cos(81\theta)\cos\theta}$
58. The value of  $\left(\sin\frac{\pi}{9}\right)\left(4 + \sec\frac{\pi}{9}\right)$  is :  
 (a)  $\frac{1}{2}$  (b)  $\sqrt{2}$  (c) 1 (d)  $\sqrt{3}$
59. If  $\frac{dy}{dx} = \sin\left(\frac{x\pi}{2}\right)\cos(x\pi)$ , then  $y$  is strictly increasing in :  
 (a) (3, 4) (b)  $\left(\frac{5}{2}, \frac{7}{2}\right)$  (c) (2, 3) (d)  $\left(\frac{1}{2}, \frac{3}{2}\right)$
60. Smallest positive value of  $\theta$  satisfying the equation  $8\sin\theta\cos2\theta\sin3\theta\cos4\theta = \cos6\theta$ ; is :  
 (a)  $\frac{\pi}{18}$  (b)  $\frac{\pi}{22}$  (c)  $\frac{\pi}{24}$  (d) None of these
61. If an angle  $A$  of a triangle  $ABC$  is given by  $3\tan A + 1 = 0$ , then  $\sin A$  and  $\cos A$  are the roots of the equation  
 (a)  $10x^2 - 2\sqrt{10}x + 3 = 0$  (b)  $10x^2 - 2\sqrt{10}x - 3 = 0$   
 (c)  $10x^2 + 2\sqrt{10}x + 3 = 0$  (d)  $10x^2 + 2\sqrt{10}x - 3 = 0$
62. If  $\theta$  is an acute angle and  $\tan\theta = \frac{1}{\sqrt{7}}$ , then the value of  $\frac{\operatorname{cosec}^2\theta - \sec^2\theta}{\operatorname{cosec}^2\theta + \sec^2\theta}$  is :  
 (a)  $3/4$  (b)  $1/2$  (c) 2 (d)  $5/4$
63. If  $2\cos\theta + \sin\theta = 1$ , then  $7\cos\theta + 6\sin\theta$  equals  
 (a) 1 or 2 (b) 2 or 3 (c) 2 or 4 (d) 2 or 6
64. If  $\sin\theta + \operatorname{cosec}\theta = 2$ , then the value of  $\sin^8\theta + \operatorname{cosec}^8\theta$  is equal to :  
 (a) 2 (b)  $2^4$  (c)  $2^8$  (d) more than  $2^8$
65. If  $\tan^3\theta + \cot^3\theta = 52$ , then the value of  $\tan^2\theta + \cot^2\theta$  is equal to :  
 (a) 14 (b) 15 (c) 16 (d) 17
66. If  $ABCD$  is a cyclic quadrilateral such that  $12\tan A - 5 = 0$  and  $5\cos B + 3 = 0$  then  $\tan C + \tan D$  is equal to :  
 (a)  $\frac{21}{12}$  (b)  $\frac{11}{12}$  (c)  $-\frac{11}{12}$  (d)  $-\frac{21}{12}$
67. If  $\frac{\pi}{2} < \theta < \frac{3\pi}{2}$  then  $\sqrt{\tan^2\theta - \sin^2\theta}$  is equal to :  
 (a)  $\tan\theta\sin\theta$  (b)  $-\tan\theta\sin\theta$  (c)  $\tan\theta - \sin\theta$  (d)  $\sin\theta - \tan\theta$

68. The value of  $\frac{\sin 10^\circ + \sin 20^\circ}{\cos 10^\circ + \cos 20^\circ}$  equals  
 (a)  $2 + \sqrt{3}$  (b)  $\sqrt{2} - 1$  (c)  $2 - \sqrt{3}$  (d)  $\sqrt{2} + 1$
69. The expression  $\cos^6 \theta + \sin^6 \theta + 3 \sin^2 \theta \cos^2 \theta$  simplifies to :  
 (a) 0 (b) 1 (c) 2 (d) 3
70.  $\frac{\sin x + \cos x}{\sin x - \cos x} - \frac{\sec^2 x + 2}{\tan^2 x - 1} =$ , where  $x \in \left(0, \frac{\pi}{2}\right)$   
 (a)  $\frac{1}{\tan x + 1}$  (b)  $\frac{2}{1 + \tan x}$  (c)  $\frac{2}{1 + \cot x}$  (d)  $\frac{2}{1 - \tan x}$
71. If  $\frac{\cot \alpha + \cot(270^\circ + \alpha)}{\cot \alpha - \cot(270^\circ + \alpha)} - 2 \cos(135^\circ + \alpha) \cos(315^\circ - \alpha) = \lambda \cos 2\alpha$ , where  $\alpha \in \left(0, \frac{\pi}{2}\right)$ , then  $\lambda =$   
 (a) 0 (b) 1 (c) 2 (d) 4
72. The expression  $\frac{\sin \alpha + \cos \alpha}{\cos \alpha - \sin \alpha} \tan\left(\frac{\pi}{4} + \alpha\right) + 1$ ,  $\alpha \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$  simplifies to :  
 (a)  $\operatorname{cosec}^2\left(\frac{\pi}{4} - \alpha\right)$  (b)  $\sec^2\left(\frac{\pi}{4} - \alpha\right)$  (c)  $\tan^2\left(\frac{\pi}{4} - \alpha\right)$  (d)  $\cot^2\left(\frac{\pi}{4} - \alpha\right)$
73. The value of expression  $\frac{\tan \alpha + \sin \alpha}{2 \cos^2 \frac{\alpha}{2}}$  for  $\alpha = \frac{\pi}{4}$  is :  
 (a) 4 (b) 3 (c) 2 (d) 1
74.  $\cos 2\alpha - \cos 3\alpha - \cos 4\alpha + \cos 5\alpha$  simplifies to :  
 (a)  $-4 \sin \frac{\alpha}{2} \sin \alpha \cos \frac{7\alpha}{2}$  (b)  $4 \sin \frac{\alpha}{2} \sin \alpha \cos \frac{7\alpha}{2}$   
 (c)  $-4 \sin \frac{\alpha}{2} \sin \frac{7\alpha}{2} \cos \alpha$  (d)  $-4 \sin \alpha \cos \frac{\alpha}{2} \sin \frac{7\alpha}{2}$
75. If  $\tan \gamma = \sec \alpha \sec \beta + \tan \alpha \tan \beta$ , then the least value of  $\cos 2\gamma$  is :  
 (a) -1 (b)  $\frac{1}{2}$  (c)  $-\frac{1}{2}$  (d) 0
76. If  $\operatorname{cosec} x = \frac{2}{\sqrt{3}}$ ,  $\cot x = -\frac{1}{\sqrt{3}}$ ,  $x \in [0, 2\pi]$ , then  $\cos x + \cos 2x + \cos 3x + \dots + \cos 100x =$   
 (a)  $\frac{1}{2}$  (b)  $-\frac{1}{2}$  (c)  $-\frac{\sqrt{3}}{2}$  (d)  $\frac{\sqrt{3}}{2}$
77. The value of  $\sum_{r=0}^{10} \cos^3\left(\frac{\pi r}{3}\right)$  is equal to :  
 (a)  $-\frac{7}{8}$  (b)  $-\frac{9}{8}$  (c)  $-\frac{3}{8}$  (d)  $-\frac{1}{8}$

78. The value of the expression  $\frac{1 - 4 \sin 10^\circ \sin 70^\circ}{2 \sin 10^\circ}$  is :
- (a) 1                      (b) 2                      (c)  $\sqrt{3}$                       (d)  $\frac{\sqrt{3}}{2}$
79. If  $x, y \in R$  and satisfy  $(x + 5)^2 + (y - 12)^2 = 14^2$ , then the minimum value of  $x^2 + y^2$  is :
- (a) 2                      (b) 1                      (c)  $\sqrt{3}$                       (d)  $\sqrt{2}$
80. If  $\theta_1, \theta_2$  and  $\theta_3$  are the three values of  $\theta \in [0, 2\pi]$  for which  $\tan \theta = \lambda$  then the value of  $\tan \frac{\theta_1}{3} \tan \frac{\theta_2}{3} + \tan \frac{\theta_2}{3} \tan \frac{\theta_3}{3} + \tan \frac{\theta_3}{3} \tan \frac{\theta_1}{3}$  is equal to ( $\lambda$  is a constant)
- (a) -3                      (b) -2                      (c) 2                      (d) 3
81. If  $\tan \alpha = \frac{b}{a}$ ,  $a > b > 0$  and if  $0 < \alpha < \frac{\pi}{4}$ , then  $\sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}}$  is equal to :
- (a)  $\frac{2 \sin \alpha}{\sqrt{\cos 2\alpha}}$                       (b)  $\frac{2 \cos \alpha}{\sqrt{\cos 2\alpha}}$                       (c)  $\frac{2 \sin \alpha}{\sqrt{\sin 2\alpha}}$                       (d)  $\frac{2 \cos \alpha}{\sqrt{\sin 2\alpha}}$
82. Minimum value of  $3 \sin \theta + 4 \cos \theta$  in the interval  $\left[0, \frac{\pi}{2}\right]$  is :
- (a) -5                      (b) 3                      (c) 4                      (d)  $\frac{7}{\sqrt{2}}$
83. If  $f(n) = \prod_{r=1}^n \cos r$ ,  $n \in N$ , then
- (a)  $|f(n)| > |f(n+1)|$                       (b)  $f(5) > 0$                       (c)  $f(4) > 0$                       (d)  $|f(n)| < |f(n+1)|$
84. If  $\tan A + \sin A = p$  and  $\tan A - \sin A = q$ , then the value of  $\frac{(p^2 - q^2)^2}{pq}$  is :
- (a) 16                      (b) 22                      (c) 18                      (d) 42
85. Let  $t_1 = (\sin \alpha)^{\cos \alpha}$ ,  $t_2 = (\sin \alpha)^{\sin \alpha}$ ,  $t_3 = (\cos \alpha)^{\cos \alpha}$ ,  $t_4 = (\cos \alpha)^{\sin \alpha}$ , where  $\alpha \in \left(0, \frac{\pi}{4}\right)$ , then which of the following is correct
- (a)  $t_3 > t_1 > t_2$                       (b)  $t_4 > t_2 > t_1$                       (c)  $t_4 > t_1 > t_2$                       (d)  $t_1 > t_3 > t_2$
86. If  $\cos A = \frac{3}{4}$ , then the value of expression  $32 \sin \frac{A}{2} \sin \frac{5A}{2}$  is equal to :
- (a) 11                      (b) -11                      (c) 12                      (d) 4
87. If  $\cos(\alpha + \beta) + \sin(\alpha - \beta) = 0$  and  $\tan \beta = \frac{1}{2009}$ ; then  $\tan 3\alpha$  is :
- (a) 2                      (b) 1                      (c) 3                      (d) 4
88. If  $2^x = 3^y = 6^{-z}$ , the value of  $\frac{1}{x} + \frac{1}{y} + \frac{1}{z}$  is equal to :
- (a) 0                      (b) 1                      (c) 2                      (d) 3

89. Let  $\alpha, \beta$  be such that  $\pi < \alpha - \beta < 3\pi$

If  $\sin \alpha + \sin \beta = -\frac{21}{65}$  and  $\cos \alpha + \cos \beta = -\frac{27}{65}$  then the value of  $\cos\left(\frac{\alpha - \beta}{2}\right)$  is :

- (a)  $\frac{-3}{\sqrt{130}}$       (b)  $\frac{3}{\sqrt{130}}$       (c)  $\frac{6}{65}$       (d)  $-\frac{6}{65}$

90. If  $\mu = \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} + \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$  then the difference between maximum and minimum values of  $\mu^2$  is :

- (a)  $2(a^2 + b^2)$       (b)  $(a + b)^2$       (c)  $2\sqrt{a^2 + b^2}$       (d)  $(a - b)^2$

91. If  $P = (\tan(3^{n+1} \theta) - \tan \theta)$  and  $Q = \sum_{r=0}^n \frac{\sin(3^r \theta)}{\cos(3^{r+1} \theta)}$ , then

- (a)  $P = 2Q$       (b)  $P = 3Q$       (c)  $2P = Q$       (d)  $3P = Q$

92. If  $270^\circ < \theta < 360^\circ$ , then find  $\sqrt{2 + \sqrt{2(1 + \cos \theta)}}$

- (a)  $-2 \sin\left(\frac{\theta}{4}\right)$       (b)  $2 \sin\left(\frac{\theta}{4}\right)$       (c)  $\pm 2 \sin \frac{\theta}{4}$       (d)  $2 \cos \frac{\theta}{4}$

93. If  $y = (\sin x + \cos x) + (\sin 4x + \cos 4x)^2$ , then :

- (a)  $y > 0 \forall x \in R$       (b)  $y \geq 0 \forall x \in R$   
(c)  $y < 2 + \sqrt{2} \forall x \in R$       (d)  $y = 2 + \sqrt{2}$  for some  $x \in R$

94. If  $\cos x + \cos y + \cos z = \sin x + \sin y + \sin z = 0$  then  $\cos(x - y) =$

- (a) 0      (b)  $-\frac{1}{2}$       (c) 2      (d) 1

95. The exact value of  $\operatorname{cosec} 10^\circ + \operatorname{cosec} 50^\circ - \operatorname{cosec} 70^\circ$  is :

- (a) 4      (b) 5      (c) 6      (d) 8

96. If  $270^\circ < \theta < 360^\circ$ , then find  $\sqrt{2 + \sqrt{2(1 + \cos \theta)}}$ :

- (a)  $-2 \sin\left(\frac{\theta}{4}\right)$       (b)  $2 \sin\left(\frac{\theta}{4}\right)$       (c)  $\pm 2 \sin \frac{\theta}{4}$       (d)  $2 \cos \frac{\theta}{4}$

## | Answers |

1.	(c)	2.	(d)	3.	(a)	4.	(c)	5.	(b)	6.	(c)	7.	(d)	8.	(d)	9.	(a)	10.	(c)
11.	(c)	12.	(b)	13.	(b)	14.	(d)	15.	(b)	16.	(d)	17.	(c)	18.	(a)	19.	(d)	20.	(a)
21.	(c)	22.	(b)	23.	(b)	24.	(d)	25.	(c)	26.	(b)	27.	(c)	28.	(b)	29.	(d)	30.	(a)
31.	(b)	32.	(c)	33.	(b)	34.	(c)	35.	(c)	36.	(a)	37.	(a)	38.	(a)	39.	(d)	40.	(d)
41.	(b)	42.	(b)	43.	(c)	44.	(d)	45.	(d)	46.	(b)	47.	(b)	48.	(c)	49.	(d)	50.	(a)
51.	(d)	52.	(b)	53.	(a)	54.	(a)	55.	(b)	56.	(b)	57.	(b)	58.	(d)	59.	(b)	60.	(a)
61.	(d)	62.	(a)	63.	(d)	64.	(a)	65.	(a)	66.	(b)	67.	(b)	68.	(c)	69.	(b)	70.	(b)
71.	(c)	72.	(a)	73.	(d)	74.	(a)	75.	(d)	76.	(b)	77.	(d)	78.	(a)	79.	(b)	80.	(a)
81.	(b)	82.	(b)	83.	(a)	84.	(a)	85.	(b)	86.	(a)	87.	(b)	88.	(a)	89.	(a)	90.	(d)
91.	(a)	92.	(b)	93.	(c)	94.	(b)	95.	(c)	96.	(b)								



**Exercise-2 : One or More than One Answer Is/are Correct**


- $\cot 12^\circ \cdot \cot 24^\circ \cdot \cot 28^\circ \cdot \cot 32^\circ \cdot \cot 48^\circ \cdot \cot 88^\circ = \dots$ 
  - $\tan 45^\circ$
  - 2
  - $2 \tan 15^\circ \cdot \tan 45^\circ \cdot \tan 75^\circ$
  - $\tan 15^\circ \cdot \tan 45^\circ \cdot \tan 75^\circ$
- If the equation  $\cot^4 x - 2 \operatorname{cosec}^2 x + a^2 = 0$  has at least one solution then possible integral values of  $a$  can be :
  - 1
  - 0
  - 1
  - 2
- Which of the following is/are true ?
  - $\tan 1 > \tan^{-1} 1$
  - $\sin 1 > \cos 1$
  - $\tan 1 < \sin 1$
  - $\cos(\cos 1) > \frac{1}{\sqrt{2}}$
- Which of the following is/are +ve ?
  - $\log_{\sin 1} \tan 1$
  - $\log_{\cos 1} (1 + \tan 3)$
  - $\log_{\log_{10} 5} (\cos \theta + \sec \theta)$
  - $\log_{\tan 15^\circ} (2 \sin 18^\circ)$
- If  $\sin \alpha + \cos \alpha = \frac{\sqrt{3} + 1}{2}$ ,  $0 < \alpha < 2\pi$ , then possible values  $\tan \frac{\alpha}{2}$  can take is/are :
  - $2 - \sqrt{3}$
  - $\frac{1}{\sqrt{3}}$
  - 1
  - $\sqrt{3}$
- If  $3 \sin \beta = \sin(2\alpha + \beta)$ , then :
  - $(\cot \alpha + \cot(\alpha + \beta))(\cot \beta - 3 \cot(2\alpha + \beta)) = 6$
  - $\sin \beta = \cos(\alpha + \beta) \sin \alpha$
  - $\tan(\alpha + \beta) = 2 \tan \alpha$
  - $2 \sin \beta = \sin(\alpha + \beta) \cos \alpha$
- If  $\sin(x + 20^\circ) = 2 \sin x \cos 40^\circ$  where  $x \in (0, 90^\circ)$ , then which of the following hold good ?
  - $\sec \frac{x}{2} = \sqrt{6} - \sqrt{2}$
  - $\cot \frac{x}{2} = 2 + \sqrt{3}$
  - $\tan 4x = \sqrt{3}$
  - $\operatorname{cosec} 4x = 2$
- If  $2(\cos(x - y) + \cos(y - z) + \cos(z - x)) = -3$ , then :
  - $\cos x \cos y \cos z = 1$
  - $\cos x + \cos y + \cos z = 0$
  - $\sin x + \sin y + \sin z = 1$
  - $\cos 3x + \cos 3y + \cos 3z = 12 \cos x \cos y \cos z$
- If  $0 < x < \frac{\pi}{2}$  and  $\sin^n x + \cos^n x \geq 1$ , then 'n' may belong to interval :
  - [1, 2]
  - [3, 4]
  - $(-\infty, 2]$
  - [-1, 1]
- If  $x = \sin(\alpha - \beta) \cdot \sin(\gamma - \delta)$ ,  $y = \sin(\beta - \gamma) \cdot \sin(\alpha - \delta)$ ,  $z = \sin(\gamma - \alpha) \cdot \sin(\beta - \delta)$ , then :
  - $x + y + z = 0$
  - $x^3 + y^3 + z^3 = 3xyz$
  - $x + y - z = 0$
  - $x^3 + y^3 - z^3 = 3xyz$

11. If  $X = x \cos \theta - y \sin \theta$ ,  $Y = x \sin \theta + y \cos \theta$  and  $X^2 + 4XY + Y^2 = Ax^2 + By^2$ ,  $0 \leq \theta \leq \pi/2$ , then :  
(where  $A$  and  $B$  are constants)
- (a)  $\theta = \frac{\pi}{6}$                       (b)  $\theta = \frac{\pi}{4}$                       (c)  $A = 3$                       (d)  $B = -1$
12. If  $2a = 2 \tan 10^\circ + \tan 50^\circ$ ;  $2b = \tan 20^\circ + \tan 50^\circ$   
 $2c = 2 \tan 10^\circ + \tan 70^\circ$ ;  $2d = \tan 20^\circ + \tan 70^\circ$   
Then which of the following is/are correct ?
- (a)  $a + d = b + c$               (b)  $a + b = c$                       (c)  $a > b < c > d$               (d)  $a < b < c < d$
13. Which of the following real numbers when simplified are neither terminating nor repeating decimal ?
- (a)  $\sin 75^\circ \cdot \cos 75^\circ$               (b)  $\log_2 28$                       (c)  $\log_3 5 \cdot \log_5 6$               (d)  $8^{-(\log_{27} 3)}$
14. If  $\alpha = \sin x \cos^3 x$  and  $\beta = \cos x \sin^3 x$ , then :
- (a)  $\alpha - \beta > 0$ ; for all  $x$  in  $\left(0, \frac{\pi}{4}\right)$                       (b)  $\alpha - \beta < 0$ ; for all  $x$  in  $\left(0, \frac{\pi}{4}\right)$   
(c)  $\alpha + \beta > 0$ ; for all  $x$  in  $\left(0, \frac{\pi}{2}\right)$                       (d)  $\alpha + \beta < 0$ ; for all  $x$  in  $\left(0, \frac{\pi}{2}\right)$
15. If  $\frac{\pi}{2} < \theta < \pi$ , then possible answers of  $\sqrt{2 + \sqrt{2 + 2 \cos 4\theta}}$  is/are :
- (a)  $2 \cos \theta$                       (b)  $2 \sin \theta$                       (c)  $-2 \sin \theta$                       (d)  $-2 \cos \theta$
16. If  $\cot^3 \alpha + \cot^2 \alpha + \cot \alpha = 1$  then which of the following is/are correct:
- (a)  $\cos 2\alpha \tan \alpha = 1$                       (b)  $\cos 2\alpha \cdot \tan \alpha = -1$   
(c)  $\cos 2\alpha - \tan 2\alpha = -1$                       (d)  $\cos 2\alpha - \tan 2\alpha = 1$
17. All values of  $x \in \left(0, \frac{\pi}{2}\right)$  such that  $\frac{\sqrt{3}-1}{\sin x} + \frac{\sqrt{3}+1}{\cos x} = 4\sqrt{2}$  are :
- (a)  $\frac{\pi}{15}$                       (b)  $\frac{\pi}{12}$                       (c)  $\frac{11\pi}{36}$                       (d)  $\frac{3\pi}{10}$
18. If  $\alpha > \frac{1}{\sin^6 x + \cos^6 x} \forall x \in R$ , then  $\alpha$  can be :
- (a) 3                      (b) 4                      (c) 5                      (d) 6
19. If  $x \in \left(0, \frac{\pi}{2}\right)$  and  $\sin x = \frac{3}{\sqrt{10}}$ ;  
Let  $k = \log_{10} \sin x + \log_{10} \cos x + 2 \log_{10} \cot x + \log_{10} \tan x$  then the value of  $k$  satisfies
- (a)  $k = 0$                       (b)  $k + 1 = 0$                       (c)  $k - 1 = 0$                       (d)  $k^2 - 1 = 0$
20. If  $A, B, C$  are angles of a triangle  $ABC$  and  $\tan A \tan C = 3$ ;  $\tan B \tan C = 6$  then which is(are) correct :
- (a)  $A = \frac{\pi}{4}$                       (b)  $\tan A \tan B = 2$                       (c)  $\frac{\tan A}{\tan C} = 3$                       (d)  $\tan B = 2 \tan A$








**Exercise-4 : Matching Type Problems**

1.

Column-I		Column-II
(A)	If $(1 + \tan 5^\circ)(1 + \tan 10^\circ) \dots (1 + \tan 45^\circ) = 2^{k+1}$ then 'k' equals	(P) 0
(B)	Sum of positive integral values of 'a' for which $a^2 - 6 \sin x - 5a \leq 0 \forall x \in R$ is	(Q) 2
(C)	The minimum value of $\frac{\left(a + \frac{1}{a}\right)^4 - \left(a^4 + \frac{1}{a^4}\right) - 2}{\left(a + \frac{1}{a}\right)^2 + a^2 + \frac{1}{a^2}}$ is	(R) 5
(D)	Number of real roots of the equation $\sum_{k=1}^3 (x-k)^2 = 0$ is	(S) 4
		(T) 5

2.

Column-I		Column-II
(A)	Maximum value of $y = \frac{1 - \tan^2(\pi/4 - x)}{1 + \tan^2(\pi/4 - x)}$	(P) 1
(B)	Minimum value of $\log_3 \left( \frac{5 \sin x - 12 \cos x + 26}{13} \right)$	(Q) 0
(C)	Minimum value of $y = -2 \sin^2 x + \cos x + 3$	(R) $\frac{7}{8}$
(D)	Maximum value of $y = 4 \sin^2 \theta + 4 \sin \theta \cos \theta + \cos^2 \theta$	(S) 5
		(T) 6

3.

Column-I		Column-II
(A)	The value of $\frac{\cos 68^\circ}{\sin 56^\circ \sin 34^\circ \tan 22^\circ}$ equals to	(P) 16
(B)	The value of $(\cos 65^\circ + \sqrt{3} \sin 5^\circ + \cos 5^\circ)^2 = \lambda \cos^2 25^\circ$ ; then value of $\lambda$ be	(Q) 3

(C)	If $\cos A = \frac{3}{4}$ ; then the value of $\frac{32}{11} \sin \frac{A}{2} \sin \frac{5A}{2}$ is equal to	(R)	4
(D)	If $7 \log_a \frac{16}{15} + 5 \log_a \frac{25}{24} + 3 \log_a \frac{81}{80} = 8$ then the value of $a^{16}$ equals to	(S)	2
		(T)	1

4.

Column-I		Column-II	
(A)	If $\sin x + \cos x = \frac{1}{5}$ ; then $ 12 \tan x $ is equal to	(P)	2
(B)	Number of values of $\theta$ lying in $(-2\pi, \pi)$ and satisfying $\cot \frac{\theta}{2} = (1 + \cot \theta)$ is	(Q)	6
(C)	If $2 - \sin^4 x + 8 \sin^2 x = \alpha$ has solution, then $\alpha$ can be	(R)	9
(D)	Number of integral values of $x$ satisfying $\log_4(2x^2 + 5x + 27) - \log_2(2x - 1) \geq 0$	(S)	14
		(T)	16

5. Match the function given in **Column-I** to the number of integers in its range given in **Column-II**.

Column-I		Column-II	
(A)	$f(x) = 2 \cos^2 x + \sin x - 8$	(P)	5
(B)	$f(x) = \sin^2 x + 3 \cos^2 x + 5$	(Q)	4
(C)	$f(x) = 4 \sin x \cos x - \sin^2 x + 3 \cos^2 x$	(R)	3
(D)	$f(x) = \cos(\sin x) + \sin(\sin x)$	(S)	2

### Answers

1. A  $\rightarrow$  S; B  $\rightarrow$  R; C  $\rightarrow$  Q; D  $\rightarrow$  P
2. A  $\rightarrow$  P; B  $\rightarrow$  Q; C  $\rightarrow$  R; D  $\rightarrow$  S
3. A  $\rightarrow$  S; B  $\rightarrow$  Q; C  $\rightarrow$  T; D  $\rightarrow$  R
4. A  $\rightarrow$  R, T; B  $\rightarrow$  P; C  $\rightarrow$  P, Q, R; D  $\rightarrow$  Q
5. A  $\rightarrow$  Q; B  $\rightarrow$  R; C  $\rightarrow$  P; D  $\rightarrow$  S

### Exercise-5 : Subjective Type Problems

- Let  $P = \frac{\sin 80^\circ \sin 65^\circ \sin 35^\circ}{\sin 20^\circ + \sin 50^\circ + \sin 110^\circ}$ , then the value of  $24P$  is :
- The value of expression  $(1 - \cot 23^\circ)(1 - \cot 22^\circ)$  is equal to :
- If  $\tan A$  and  $\tan B$  are the roots of the quadratic equation,  $4x^2 - 7x + 1 = 0$  then evaluate  $4\sin^2(A+B) - 7\sin(A+B) \cdot \cos(A+B) + \cos^2(A+B)$ .
- $A_1 A_2 A_3 \dots A_{18}$  is a regular 18 sided polygon.  $B$  is an external point such that  $A_1 A_2 B$  is an equilateral triangle. If  $A_{18} A_1$  and  $A_1 B$  are adjacent sides of a regular  $n$  sided polygon, then  $n =$
- If  $10\sin^4 \alpha + 15\cos^4 \alpha = 6$  and the value of  $9\operatorname{cosec}^4 \alpha + \beta \sec^4 \alpha$  is  $S$ , then find the value of  $\frac{S}{25}$ .
- The value of  $\left(1 + \tan \frac{3\pi}{8} \tan \frac{\pi}{8}\right) + \left(1 + \tan \frac{5\pi}{8} \tan \frac{3\pi}{8}\right) + \left(1 + \tan \frac{7\pi}{8} \tan \frac{5\pi}{8}\right) + \left(1 + \tan \frac{9\pi}{8} \tan \frac{7\pi}{8}\right)$
- If  $\alpha = \frac{\pi}{7}$  then find the value of  $\left(\frac{1}{\cos \alpha} + \frac{2\cos \alpha}{\cos 2\alpha}\right)$ .
- Given that for  $a, b, c, d \in R$ , if  $a \sec(200^\circ) - c \tan(200^\circ) = d$  and  $b \sec(200^\circ) + d \tan(200^\circ) = c$ , then find the value of  $\left(\frac{a^2 + b^2 + c^2 + d^2}{bd - ac}\right) \sin 20^\circ$ .
- The expression  $2 \cos \frac{\pi}{17} \cdot \cos \frac{9\pi}{17} + \cos \frac{7\pi}{17} + \cos \frac{9\pi}{17}$  simplifies to an integer  $P$ . Find the value of  $P$ .
- If the expression  $\frac{\sin \theta \sin 2\theta + \sin 3\theta \sin 6\theta + \sin 4\theta \sin 13\theta}{\sin \theta \cos 2\theta + \sin 3\theta \cos 6\theta + \sin 4\theta \cos 13\theta} = \tan k\theta$ , where  $k \in N$ . Find the value of  $k$ .
- Let  $a = \sin 10^\circ$ ,  $b = \sin 50^\circ$ ,  $c = \sin 70^\circ$ , then  $8abc \left(\frac{a+b}{c}\right) \left(\frac{1}{a} + \frac{1}{b} - \frac{1}{c}\right)$  is equal to
- If  $\sin^3 \theta + \sin^3 \left(\theta + \frac{2\pi}{3}\right) + \sin^3 \left(\theta + \frac{4\pi}{3}\right) = a \sin b\theta$ . Find the value of  $\left|\frac{b}{a}\right|$ .
- If  $\sum_{r=1}^n \left(\frac{\tan 2^{r-1}}{\cos 2^r}\right) = \tan p^n - \tan q$ , then find the value of  $(p+q)$ .
- If  $x = \sec \theta - \tan \theta$  and  $y = \operatorname{cosec} \theta + \cot \theta$ , then  $y - x - xy =$
- If  $\cos 18^\circ - \sin 18^\circ = \sqrt{n} \sin 27^\circ$ , then  $n =$
- The value of  $3(\sin 1 - \cos 1)^4 + 6(\sin 1 + \cos 1)^2 + 4(\sin^6 1 + \cos^6 1)$  is equal to
- If  $x = \alpha$  satisfy the equation  $3^{\sin 2x + 2\cos^2 x} + 3^{1 - \sin 2x + 2\sin^2 x} = 28$ , then  $(\sin 2\alpha - \cos 2\alpha)^2 + 8 \sin 4\alpha$  is equal to :
- The least value of the expression  $(\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2 \forall \theta \in R$  is
- If  $\tan 20^\circ + \tan 40^\circ + \tan 80^\circ - \tan 60^\circ = \lambda \sin 40^\circ$ , then  $\lambda$  is equal to




20. If  $K^\circ$  lies between  $360^\circ$  and  $540^\circ$  and  $K^\circ$  satisfies the equation  $1 + \cos 10x \cos 6x = 2 \cos^2 8x + \sin^2 8x$ , then  $\frac{K}{10} =$
21. If  $\cos 20^\circ + 2 \sin^2 55^\circ = 1 + \sqrt{2} \sin K^\circ$ ,  $K \in (0, 90)$ , then  $K =$
22. The exact value of  $\operatorname{cosec} 10^\circ + \operatorname{cosec} 50^\circ - \operatorname{cosec} 70^\circ$  is :
23. Let  $\alpha$  be the smallest integral value of  $x$ ,  $x > 0$  such that  $\tan 19x = \frac{\cos 96^\circ + \sin 96^\circ}{\cos 96^\circ - \sin 96^\circ}$ . The last digit of  $\alpha$  is :
24. Find the value of the expression  $\frac{\sin 20^\circ (4 \cos 20^\circ + 1)}{\cos 20^\circ \cos 30^\circ}$
25. If the value of  $\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} + \cos \frac{7\pi}{7} = -\frac{l}{2}$ . Find the value of  $l$ .
26. If  $\cos A = \frac{3}{4}$  and  $k \sin\left(\frac{A}{2}\right) \sin\left(\frac{5A}{2}\right) = \frac{11}{8}$ . Find  $k$ .
27. Find the least value of the expression  $3 \sin^2 x + 4 \cos^2 x$ .
28. If  $\tan \alpha$  and  $\tan \beta$  are the roots of equation  $x^2 - 12x - 3 = 0$ , then the value of  $\sin^2(\alpha + \beta) + 2 \sin(\alpha + \beta) \cos(\alpha + \beta) + 5 \cos^2(\alpha + \beta)$  is :
29. The value of  $\frac{\cos 24^\circ}{2 \tan 33^\circ \sin^2 57^\circ} + \frac{\sin 162^\circ}{\sin 18^\circ - \cos 18^\circ \tan 9^\circ} + \cos 162^\circ$  is equal to :
30. Find the value of  $\tan \theta (1 + \sec 2\theta)(1 + \sec 4\theta)(1 + \sec 8\theta)$ , when  $\theta = \frac{\pi}{32}$ .
31. If  $\lambda$  be the minimum value of  $y = (\sin x + \operatorname{cosec} x)^2 + (\cos x + \sec x)^2 + (\tan x + \cot x)^2$  where  $x \in \mathbb{R}$ . Find  $\lambda - 6$ .

### Answers

1.	6	2.	2	3.	1	4.	9	5.	3	6.	0	7.	4
8.	2	9.	0	10.	9	11.	6	12.	4	13.	3	14.	1
15.	2	16.	13	17.	1	18.	9	19.	8	20.	45	21.	65
22.	6	23.	9	24.	2	25.	3	26.	4	27.	3	28.	2
29.	2	30.	1	31.	7								

□□□


**Exercise-1 : Single Choice Problems**

1. Let  $x$  and  $y$  be 2 real numbers which satisfy the equations  $(\tan^2 x - \sec^2 y) = \frac{5a}{6} - 3$  and  $(-\sec^2 x + \tan^2 y) = a^2$ , then the product of all possible value's of  $a$  can be equal to :
- (a) 0                      (b)  $\frac{-2}{3}$                       (c) -1                      (d)  $\frac{-3}{2}$
2. The general solution of the equation  $\tan^2(x+y) + \cot^2(x+y) = 1 - 2x - x^2$  lie on the line is :
- (a)  $x = -1$                       (b)  $x = -2$                       (c)  $y = -1$                       (d)  $y = -2$
3. General solution of the equation  $\sin x + \cos x = \min_{a \in R} \{1, a^2 - 4a + 6\}$  is :
- (a)  $\frac{n\pi}{2} + (-1)^n \frac{\pi}{4}$                       (b)  $2n\pi + (-1)^n \frac{\pi}{4}$   
 (c)  $n\pi + (-1)^{n+1} \frac{\pi}{4}$                       (d)  $n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{4}$
- (where  $n \in I$ ,  $I$  represent set of integers)
4. The number of solutions of the equation  $\left(2 \sin\left(\frac{\sin x}{2}\right)\right) \left(\cos\left(\frac{\sin x}{2}\right)\right) \left(\sin\left(2 \tan \frac{x}{2} \cos^2 \frac{x}{2}\right) - 3\right) + 2 = 0$  in  $[0, 2\pi]$  is :
- (a) 0                      (b) 1                      (c) 2                      (d) 4
5. Number of solution of  $\tan(2x) = \tan(6x)$  in  $(0, 3\pi)$  is :
- (a) 4                      (b) 5                      (c) 3                      (d) None of these
6. The number of values of  $x$  in the interval  $[0, 5\pi]$  satisfying the equation  $3 \sin^2 x - 7 \sin x + 2 = 0$  is :
- (a) 0                      (b) 2                      (c) 6                      (d) 8

7. The number of different values of  $\theta$  satisfying the equation  $\cos \theta + \cos 2\theta = -1$ , and at the same time satisfying the condition  $0 < \theta < 360^\circ$  is :
- (a) 1 (b) 2 (c) 3 (d) 4
8. The total number of solutions of the equation  $\max(\sin x, \cos x) = \frac{1}{2}$  for  $x \in (-2\pi, 5\pi)$  is equal to:
- (a) 3 (b) 6 (c) 7 (d) 8
9. The general value of  $x$  satisfying the equation  $2 \cot^2 x + 2\sqrt{3} \cot x + 4 \operatorname{cosec} x + 8 = 0$  is : (where  $n \in I$ )
- (a)  $n\pi - \frac{\pi}{6}$  (b)  $n\pi + \frac{\pi}{6}$  (c)  $2n\pi - \frac{\pi}{6}$  (d)  $2n\pi + \frac{\pi}{6}$
10. The general solution of the equation  $\sin^2 x + \cos^2 3x = 1$  is equal to :
- (a)  $x = \frac{n\pi}{2}$  (b)  $x = n\pi + \frac{\pi}{4}$  (c)  $x = \frac{n\pi}{4}$  (d)  $x = n\pi + \frac{\pi}{2}$   
(where  $n \in I$ )
11. Values of  $x$  between 0 and  $2\pi$  which satisfy the equation  $\sin x \sqrt{8 \cos^2 x} = 1$  are in A.P. whose common difference is :
- (a)  $\frac{\pi}{4}$  (b)  $\frac{\pi}{3}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{2\pi}{3}$
12. Number of solutions of  $\sum_{r=1}^5 \cos rx = 5$  in the interval  $[0, 4\pi]$  is :
- (a) 0 (b) 2 (c) 3 (d) 7
13. General solution of  $4 \sin^2 x + \tan^2 x + \operatorname{cosec}^2 x + \cot^2 x - 6 = 0$  is :
- (a)  $n\pi \pm \frac{\pi}{4}$  (b)  $2n\pi \pm \frac{\pi}{4}$  (c)  $n\pi + \frac{\pi}{3}$  (d)  $n\pi - \frac{\pi}{6}$   
[where  $n \in I$ ]
14. Smallest positive  $x$  satisfying the equation  $\cos^3 3x + \cos^3 5x = 8 \cos^3 4x \cdot \cos^3 x$  is :
- (a)  $15^\circ$  (b)  $18^\circ$  (c)  $22.5^\circ$  (d)  $30^\circ$
15. The general solution of the equation  $\sin^{100} x - \cos^{100} x = 1$  is (where  $n \in I$ ):
- (a)  $2n\pi + \frac{\pi}{2}$  (b)  $n\pi + \frac{\pi}{2}$  (c)  $2n\pi - \frac{\pi}{2}$  (d)  $n\pi$
16. Number of solution(s) of equation  $\sin \theta = \sec^2 4\theta$  in  $[0, \pi]$  is/are:
- (a) 0 (b) 1 (c) 2 (d) 3
17. The number of solutions of the equation  $4 \sin^2 x + \tan^2 x + \cot^2 x + \operatorname{cosec}^2 x = 6$  in  $[0, 2\pi]$
- (a) 1 (b) 2 (c) 3 (d) 4
18. The number of solutions of the equation  $\sin^4 \theta - 2 \sin^2 \theta - 1 = 0$  which lie between 0 and  $2\pi$  is :
- (a) 0 (b) 2 (c) 4 (d) 8

19. The smallest positive value of  $p$  for which the equation  $\cos(p \sin x) = \sin(p \cos x)$  has solution in  $0 \leq x \leq 2\pi$  is :
- (a)  $\frac{\pi}{\sqrt{2}}$                       (b)  $\frac{\pi}{2}$                       (c)  $\frac{\pi}{2\sqrt{2}}$                       (d)  $\frac{3\pi}{2\sqrt{2}}$
20. The total number of ordered pairs  $(x, y)$  satisfying  $|x| + |y| = 2$  and  $\sin\left(\frac{\pi x^2}{3}\right) = 1$  is :
- (a) 2                      (b) 4                      (c) 6                      (d) 8
21. The complete set of values of  $x, x \in \left(-\frac{\pi}{2}, \pi\right)$  satisfying the inequality  $\cos 2x > |\sin x|$  is :
- (a)  $\left(-\frac{\pi}{6}, \frac{\pi}{6}\right)$                       (b)  $\left(-\frac{\pi}{2}, \frac{\pi}{6}\right) \cup \left(\frac{\pi}{6}, \frac{5\pi}{6}\right)$   
(c)  $\left(-\frac{\pi}{2}, -\frac{\pi}{6}\right) \cup \left(\frac{5\pi}{6}, \pi\right)$                       (d)  $\left(-\frac{\pi}{6}, \frac{\pi}{6}\right) \cup \left(\frac{5\pi}{6}, \pi\right)$
22. The total number of solution of the equation  $\sin^4 x + \cos^4 x = \sin x \cos x$  in  $[0, 2\pi]$  is :
- (a) 2                      (b) 4                      (c) 6                      (d) 8
23. Number of solution of the equation  $\sin \frac{5x}{2} - \sin \frac{x}{2} = 2$  in the interval  $[0, 2\pi]$ , is :
- (a) 1                      (b) 2                      (c) 0                      (d) Infinite
24. In the interval  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ . The equation  $\log_{\sin \theta} \cos 2\theta = 2$  has
- (a) No solution                      (b) One solution                      (c) Two solution                      (d) Infinite solution
25. If  $\alpha$  and  $\beta$  are 2 distinct roots of equation  $a \cos \theta + b \sin \theta = C$  then  $\cos(\alpha + \beta) =$
- (a)  $\frac{2ab}{a^2 + b^2}$                       (b)  $\frac{2ab}{a^2 - b^2}$                       (c)  $\frac{a^2 + b^2}{a^2 - b^2}$                       (d)  $\frac{a^2 - b^2}{a^2 + b^2}$

## Answers

1. (c)	2. (a)	3. (d)	4. (a)	5. (b)	6. (c)	7. (d)	8. (c)	9. (c)	10. (c)
11. (a)	12. (c)	13. (a)	14. (b)	15. ( )	16. (b)	17. (d)	18. (a)	19. (c)	20. (b)
21. (d)	22. (a)	23. (c)	24. (b)	25. (d)					

**Exercise-2 : One or More than One Answer is/are Correct**

1. If  $2 \cos \theta + 2\sqrt{2} = 3 \sec \theta$  where  $\theta \in (0, 2\pi)$  then which of the following can be correct ?  
 (a)  $\cos \theta = \frac{1}{\sqrt{2}}$  (b)  $\tan \theta = 1$  (c)  $\sin \theta = -\frac{1}{\sqrt{2}}$  (d)  $\cot \theta = -1$
2. In a triangle  $ABC$  if  $\tan C < 0$  then :  
 (a)  $\tan A \tan B < 1$  (b)  $\tan A \tan B > 1$   
 (c)  $\tan A + \tan B + \tan C < 0$  (d)  $\tan A + \tan B + \tan C > 0$
3. The inequality  $4 \sin 3x + 5 \geq 4 \cos 2x + 5 \sin x$  is true for  $x \in$   
 (a)  $\left[-\pi, \frac{3\pi}{2}\right]$  (b)  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$  (c)  $\left[\frac{5\pi}{8}, \frac{13\pi}{8}\right]$  (d)  $\left[\frac{23\pi}{14}, \frac{41\pi}{14}\right]$
4. The least difference between the roots of the equation  $4 \cos x(2 - 3 \sin^2 x) + \cos 2x + 1 = 0$   $\forall x \in R$  is :  
 (a) equal to  $\frac{\pi}{2}$  (b)  $> \frac{\pi}{10}$  (c)  $< \frac{\pi}{2}$  (d)  $< \frac{\pi}{3}$
5. The equation  $\cos x \cos 6x = -1$  :  
 (a) has 50 solutions in  $[0, 100\pi]$  (b) has 3 solutions in  $[0, 3\pi]$   
 (c) has even number of solutions in  $(3\pi, 13\pi)$  (d) has one solution in  $\left[\frac{\pi}{2}, \pi\right]$
6. Identify the correct options :  
 (a)  $\frac{\sin 3\alpha}{\cos 2\alpha} > 0$  for  $\alpha \in \left(\frac{3\pi}{8}, \frac{23\pi}{48}\right)$  (b)  $\frac{\sin 3\alpha}{\cos 2\alpha} < 0$  for  $\alpha \in \left(\frac{13\pi}{48}, \frac{14\pi}{48}\right)$   
 (c)  $\frac{\sin 2\alpha}{\cos \alpha} < 0$  for  $\alpha \in \left(-\frac{\pi}{2}, 0\right)$  (d)  $\frac{\sin 2\alpha}{\cos \alpha} > 0$  for  $\alpha \in \left(\frac{13\pi}{48}, \frac{14\pi}{48}\right)$
7. The equation  $\sin^4 x + \cos^4 x + \sin 2x + k = 0$  must have real solutions if :  
 (a)  $k = 0$  (b)  $|k| \leq \frac{1}{2}$   
 (c)  $-\frac{3}{2} \leq k \leq \frac{1}{2}$  (d)  $-\frac{1}{2} \leq k \leq \frac{3}{2}$
8. Let  $f(\theta) = \left(\cos \theta - \cos \frac{\pi}{8}\right)\left(\cos \theta - \cos \frac{3\pi}{8}\right)\left(\cos \theta - \cos \frac{5\pi}{8}\right)\left(\cos \theta - \cos \frac{7\pi}{8}\right)$  then :  
 (a) maximum value of  $f(\theta) \forall \theta \in R$  is  $\frac{1}{4}$   
 (b) maximum value of  $f(\theta) \forall \theta \in R$  is  $\frac{1}{8}$   
 (c)  $f(0) = \frac{1}{8}$   
 (d) Number of principle solutions of  $f(\theta) = 0$  is 8

9. If  $\frac{\sin^2 2x + 4\sin^4 x - 4\sin^2 x \cdot \cos^2 x}{4 - \sin^2 2x - 4\sin^2 x} = \frac{1}{9}$  and  $0 < x < \pi$ . Then the value of  $x$  is :

- (a)  $\frac{\pi}{3}$                       (b)  $\frac{\pi}{6}$                       (c)  $\frac{2\pi}{3}$                       (d)  $\frac{5\pi}{6}$

10. The possible value(s) of ' $\theta$ ' satisfying the equation

$$\sin^2 \theta \tan \theta + \cos^2 \theta \cot \theta - \sin 2\theta = 1 + \tan \theta + \cot \theta$$

where  $\theta \in [0, \pi]$  is/are :

- (a)  $\frac{\pi}{4}$                       (b)  $\pi$                       (c)  $\frac{7\pi}{12}$                       (d) None of these


11. If  $\sin \theta + \sqrt{3} \cos \theta = 6x - x^2 - 11$ ,  $0 \leq \theta \leq 4\pi$ ,  $x \in R$  holds for

- (a) no values of  $x$  and  $\theta$                       (b) one value of  $x$  and two values of  $\theta$   
 (c) two values of  $x$  and two values of  $\theta$                       (d) two pairs of values of  $(x, \theta)$

### Answers

1.	(a, b, c, d)	2.	(a, c)	3.	(a, b, c, d)	4.	(b, c, d)	5.	(a, c, d)	6.	(a, b, c, d)
7.	(a, b, c)	8.	(b, c, d)	9.	(b, d)	10.	(c)	11.	(b, d)		




**Exercise-4 : Matching Type Problems**

1.

Column-I		Column-II	
(A)	If $\sin x + \cos x = \frac{1}{5}$ ; then $ 12 \tan x $ is equal to	(P)	2
(B)	Number of values of $\theta$ lying in $(-2\pi, \pi)$ and satisfying $\cot \frac{\theta}{2} = (1 + \cot \theta)$ is	(Q)	6
(C)	If $2 - \sin^4 x + 8 \sin^2 x = \alpha$ has solution, then $\alpha$ can be	(R)	9
(D)	Number of integral values of $x$ satisfying $\log_4(2x^2 + 5x + 27) - \log_2(2x - 1) \geq 0$	(S)	14
		(T)	16

2.

Column-I		Column-II	
(A)	If $x, y \in [0, 2\pi]$ , then total number of ordered pair $(x, y)$ satisfying $\sin x \cos y = 1$ is	(P)	4
(B)	If $f(x) = \sin x - \cos x - kx + b$ decreases for all real values of $x$ , then $2\sqrt{2}k$ may be	(Q)	0
(C)	The number of solution of the equation $\sin^{-1}( x^2 - 1 ) + \cos^{-1}( 2x^2 - 5 ) = \frac{\pi}{2}$ is	(R)	2
(D)	The number of ordered pair $(x, y)$ satisfying the equation $\sin x + \sin y = \sin(x + y)$ and $ x  +  y  = 1$ is	(S)	3
		(T)	6

3.

Column-I		Column-II	
(A)	Minimum value of $y = 4 \sec^2 x + \cos^2 x$ for permissible real values of $x$ is equal to	(P)	2
(B)	If $m, n$ are positive integers and $m + n\sqrt{2} = \sqrt{41 + 24\sqrt{2}}$ then $(m + n)$ is equal to :	(Q)	7



<b>(C)</b>	Number of solutions of the equation : $\log\left(\frac{9x-x^2-14}{7}\right)(\sin 3x - \sin x) = \log\left(\frac{9x-x^2-14}{7}\right)\cos 2x$ is equal to :	<b>(R)</b>	4
<b>(D)</b>	Consider an arithmetic sequence of positive integers. If the sum of the first ten terms is equal to the 58th term, then the least possible value of the first term is equal to :	<b>(S)</b>	5
		<b>(T)</b>	3

### Answers

1. A → R, T; B → P; C → P, Q, R; D → Q
2. A → S; B → P, T; C → R; D → T
3. A → S; B → Q; C → P; D → R

### Exercise-5 : Subjective Type Problems

- Find the number of solutions of the equations  
 $(\sin x - 1)^3 + (\cos x - 1)^3 + (\sin x)^3 = (2 \sin x + \cos x - 2)^3$  in  $[0, 2\pi]$ .
- If  $x + \sin y = 2014$  and  $x + 2014 \cos y = 2013$ ,  $0 \leq y \leq \frac{\pi}{2}$ , then find the value of  $[x + y] - 2005$   
 (where  $[ ]$  denotes greatest integer function)
- The complete set of values of  $x$  satisfying  $\frac{2 \sin 6x}{\sin x - 1} < 0$  and  $\sec^2 x - 2\sqrt{2} \tan x \leq 0$  in  $\left(0, \frac{\pi}{2}\right)$  is  $[a, b) \cup (c, d]$ , then find the value of  $\left(\frac{cd}{ab}\right)$ .
- The range of value's of  $k$  for which the equation  $2 \cos^4 x - \sin^4 x + k = 0$  has atleast one solution is  $[\lambda, \mu]$ . Find the value of  $(9\mu + \lambda)$ .
- The number of points in interval  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ , where the graphs of the curves  $y = \cos x$  and  $y = \sin 3x$ ,  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$  intersects is
- The number of solutions of the system of equations :  

$$2 \sin^2 x + \sin^2 2x = 2$$

$$\sin 2x + \cos 2x = \tan x$$
 in  $[0, 4\pi]$  satisfying  $2 \cos^2 x + \sin x \leq 2$  is :
- If the sum of all the solutions of the equation  $3 \cot^2 \theta + 10 \cot \theta + 3 = 0$  in  $[0, 2\pi]$  is  $k\pi$  where  $k \in I$ , then find the value of  $k$ .
- If the sum of all values of  $\theta$ ,  $0 \leq \theta \leq 2\pi$  satisfying the equation  $(8 \cos 4\theta - 3)(\cot \theta + \tan \theta - 2)(\cot \theta + \tan \theta + 2) = 12$  is  $k\pi$ , then  $k$  is equal to :
- Find the number of solutions of the equation  $2 \sin^2 x + \sin^2 2x = 2$ ;  $\sin 2x + \cos 2x = \tan x$  in  $[0, 4\pi]$  satisfying the condition  $2 \cos^2 x + \sin x \leq 2$ .

### Answers

1.	5	2.	9	3.	6	4.	7	5.	3	6.	8	7.	5
8.	8	9.	8										

□□□



8. In a  $\Delta ABC$ ,  $\angle B = \frac{\pi}{3}$  and  $\angle C = \frac{\pi}{4}$  let  $D$  divide  $BC$  internally in the ratio  $1 : 3$ , then  $\frac{\sin(\angle BAD)}{\sin(\angle CAD)}$  is equal to :
- (a)  $\frac{1}{\sqrt{6}}$  (b)  $\frac{1}{3}$  (c)  $\frac{1}{\sqrt{3}}$  (d)  $\frac{\sqrt{2}}{3}$
9. Let  $AD, BE, CF$  be the lengths of internal bisectors of angles  $A, B, C$  respectively of triangle  $ABC$ . Then the harmonic mean of  $AD \sec \frac{A}{2}, BE \sec \frac{B}{2}, CF \sec \frac{C}{2}$  is equal to :
- (a) Harmonic mean of sides of  $\Delta ABC$  (b) Geometric mean of sides of  $\Delta ABC$   
 (c) Arithmetic mean of sides of  $\Delta ABC$  (d) Sum of reciprocals of the sides of  $\Delta ABC$
10. In triangle  $ABC$ , if  $2b = a + c$  and  $A - C = 90^\circ$ , then  $\sin B$  equals :
- [Note : All symbols used have usual meaning in triangle  $ABC$ .]
- (a)  $\frac{\sqrt{7}}{5}$  (b)  $\frac{\sqrt{5}}{8}$  (c)  $\frac{\sqrt{7}}{4}$  (d)  $\frac{\sqrt{5}}{3}$
11. In a triangle  $ABC$ , if  $2a \cos\left(\frac{B-C}{2}\right) = b + c$ , then  $\sec A$  is equal to :
- (All symbols used have usual meaning in a triangle.)
- (a)  $\frac{2}{\sqrt{3}}$  (b)  $\sqrt{2}$  (c) 2 (d) 3
12. Triangle  $ABC$  has  $BC = 1$  and  $AC = 2$ , then maximum possible value of  $\angle A$  is :
- (a)  $\frac{\pi}{6}$  (b)  $\frac{\pi}{4}$  (c)  $\frac{\pi}{3}$  (d)  $\frac{\pi}{2}$
13.  $\Delta I_1 I_2 I_3$  is an excentral triangle of an equilateral triangle  $\Delta ABC$  such that  $I_1 I_2 = 4$  unit, if  $\Delta DEF$  is pedal triangle of  $\Delta ABC$ , then  $\frac{Ar(\Delta I_1 I_2 I_3)}{Ar(\Delta DEF)} =$
- (a) 16 (b) 4 (c) 2 (d) 1
14. Let  $ABC$  be a triangle with  $\angle BAC = \frac{2\pi}{3}$  and  $AB = x$  such that  $(AB)(AC) = 1$ . If  $x$  varies then the longest possible length of the internal angle bisector  $AD$  equals :
- (a)  $\frac{1}{3}$  (b)  $\frac{1}{2}$   
 (c)  $\frac{2}{3}$  (d)  $\frac{\sqrt{2}}{3}$
15. In an equilateral triangle  $r, R$  and  $r_1$  form (where symbols used have usual meaning)
- (a) an A.P. (b) a G.P. (c) an H.P. (d) none of these
16. In  $\Delta ABC$  if  $\frac{\sin A}{\sin C} = \frac{\sin(A-B)}{\sin(B-C)}$ , then  $a^2, b^2, c^2$  are in :
- (a) A.P. (b) G.P. (c) H.P. (d) none of these

17. In  $\Delta ABC$ ,  $\tan A = 2$ ,  $\tan B = \frac{3}{2}$  and  $c = \sqrt{65}$ , then circumradius of the triangle is :
- (a) 65                      (b)  $\frac{65}{7}$                       (c)  $\frac{65}{14}$                       (d) none of these
18. If the sides  $a, b, c$  of a triangle  $ABC$  are the roots of the equation  $x^3 - 13x^2 + 54x - 72 = 0$ , then the value of  $\frac{\cos A}{a} + \frac{\cos B}{b} + \frac{\cos C}{c}$  is equal to :
- (a)  $\frac{61}{144}$                       (b)  $\frac{61}{72}$                       (c)  $\frac{169}{144}$                       (d)  $\frac{59}{144}$
19. In  $\Delta ABC$ , if  $\angle C = 90^\circ$ , then  $\frac{a+c}{b} + \frac{b+c}{a}$  is equal to :
- (a)  $\frac{c}{r}$                       (b)  $\frac{1}{2Rr}$                       (c) 2                      (d)  $\frac{R}{r}$
20. In a  $\Delta ABC$ , if  $a^2 \sin B = b^2 + c^2$ , then :
- (a)  $\angle A$  is obtuse                      (b)  $\angle A$  is acute                      (c)  $\angle B$  is obtuse                      (d)  $\angle A$  is right angle
21. If  $R$  and  $R'$  are the circumradii of triangles  $ABC$  and  $OBC$ , where  $O$  is the orthocenter of triangle  $ABC$ , then :
- (a)  $R' = \frac{R}{2}$                       (b)  $R' = 2R$                       (c)  $R' = R$                       (d)  $R' = 3R$
22. The acute angle of a rhombus whose side is geometric mean between its diagonals, is :
- (a)  $15^\circ$                       (b)  $20^\circ$                       (c)  $30^\circ$                       (d)  $60^\circ$
23. In a  $\Delta ABC$  right angled at  $A$ , a line is drawn through  $A$  to meet  $BC$  at  $D$  dividing  $BC$  in  $2 : 1$ . If  $\tan(\angle ADC) = 3$  then  $\angle BAD$  is :
- (a)  $30^\circ$                       (b)  $45^\circ$                       (c)  $60^\circ$                       (d)  $75^\circ$
24. A circle is circumscribed in an equilateral triangle of side ' $l$ '. The area of any square inscribed in the circle is :
- (a)  $\frac{4}{3}l^2$                       (b)  $\frac{2}{3}l^2$                       (c)  $\frac{1}{3}l^2$                       (d)  $l^2$
25. If the sides of a triangle are in the ratio  $2 : \sqrt{6} : (\sqrt{3} + 1)$ , then the largest angle of the triangle will be :
- (a)  $60^\circ$                       (b)  $72^\circ$                       (c)  $75^\circ$                       (d)  $90^\circ$
26. In a triangle  $ABC$  if  $a, b, c$  are in A.P. and  $C - A = 120^\circ$ , then  $\frac{s}{r} =$
- (where notations have their usual meaning)
- (a)  $\sqrt{15}$                       (b)  $2\sqrt{15}$                       (c)  $3\sqrt{15}$                       (d)  $6\sqrt{15}$
27. In a triangle  $ABC$ ,  $a = 5, b = 4$  and  $\cos(A - B) = \frac{31}{32}$ , then the third side is equal to :
- (where symbols used have usual meanings)
- (a)  $\sqrt{6}$                       (b)  $6\sqrt{6}$                       (c) 6                      (d)  $(216)^{1/4}$

28. If semiperimeter of a triangle is 15, then the value of  $(b + c) \cos(B + C) + (c + a) \cos(C + A) + (a + b) \cos(A + B)$  is equal to :  
(where symbols used have usual meanings)
- (a) -60 (b) -15  
(c) -30 (d) can not be determined
29. Let triangle  $ABC$  be an isosceles triangle with  $AB = AC$ . Suppose that the angle bisector of its angle  $B$  meets the side  $AC$  at a point  $D$  and that  $BC = BD + AD$ . Measure of the angle  $A$  in degrees, is :
- (a) 80 (b) 100 (c) 110 (d) 130
30. In triangle  $ABC$  if  $A : B : C = 1 : 2 : 4$ , then  $(a^2 - b^2)(b^2 - c^2)(c^2 - a^2) = \lambda a^2 b^2 c^2$ , where  $\lambda =$   
(where notations have their usual meaning)
- (a) 1 (b) 2 (c) 4 (d) 9
31. In a triangle  $ABC$  with altitude  $AD$ ,  $\angle BAC = 45^\circ$ ,  $DB = 3$  and  $CD = 2$ . The area of the triangle  $ABC$  is :
- (a) 6 (b) 15 (c)  $15/4$  (d) 12
32. A triangle has base 10 cm long and the base angles of  $50^\circ$  and  $70^\circ$ . If the perimeter of the triangle is  $x + y \cos z^\circ$  where  $z \in (0, 90)$  then the value of  $x + y + z$  equals :
- (a) 60 (b) 55 (c) 50 (d) 40
33. Let  $H$  be the orthocenter of triangle  $ABC$ , then angle subtended by side  $BC$  at the centre of incircle of  $\triangle CHB$  is :
- (a)  $\frac{A}{2} + \frac{\pi}{2}$  (b)  $\frac{B+C}{2} + \frac{\pi}{2}$  (c)  $\frac{B-C}{2} + \frac{\pi}{2}$  (d)  $\frac{B+C}{2} + \frac{\pi}{4}$
34. Triangle  $ABC$  is right angled at  $A$ . The points  $P$  and  $Q$  are on the hypotenuse  $BC$  such that  $BP = PQ = QC$ . If  $AP = 3$  and  $AQ = 4$  then the length  $BC$  is equal to :
- (a)  $\sqrt{27}$  (b)  $\sqrt{36}$  (c)  $\sqrt{45}$  (d)  $\sqrt{54}$
35. In a  $\triangle ABC$  if  $b = a(\sqrt{3} - 1)$  and  $\angle C = 30^\circ$  then the measure of the angle  $A$  is :
- (a)  $15^\circ$  (b)  $45^\circ$  (c)  $75^\circ$  (d)  $105^\circ$
36. Through the centroid of an equilateral triangle, a line parallel to the base is drawn. On this line, an arbitrary point  $P$  is taken inside the triangle. Let  $h$  denote the perpendicular distance of  $P$  from the base of the triangle. Let  $h_1$  and  $h_2$  be the perpendicular distance of  $P$  from the other two sides of the triangle. Then :
- (a)  $h = \frac{h_1 + h_2}{2}$  (b)  $h = \sqrt{h_1 h_2}$   
(c)  $h = \frac{2h_1 h_2}{h_1 + h_2}$  (d)  $h = \frac{(h_1 + h_2)\sqrt{3}}{4}$
37. The angles  $A, B$  and  $C$  of a triangle  $ABC$  are in arithmetic progression.  $AB = 6$  and  $BC = 7$ . Then  $AC$  is :
- (a)  $\sqrt{41}$  (b)  $\sqrt{39}$  (c)  $\sqrt{42}$  (d)  $\sqrt{43}$

38. In  $\Delta ABC$ , If  $A - B = 120^\circ$  and  $R = 8r$ , then the value of  $\frac{1 + \cos C}{1 - \cos C}$  equals :  
 (All symbols used have their usual meaning in a triangle)  
 (a) 12 (b) 15 (c) 21 (d) 31
39. The lengths of the sides  $CB$  and  $CA$  of a triangle  $ABC$  are given by  $a$  and  $b$  and the angle  $C$  is  $\frac{2\pi}{3}$ .  
 The line  $CD$  bisects the angle  $C$  and meets  $AB$  at  $D$ . Then the length of  $CD$  is :  
 (a)  $\frac{1}{a+b}$  (b)  $\frac{a^2 + b^2}{a+b}$  (c)  $\frac{ab}{2(a+b)}$  (d)  $\frac{ab}{a+b}$
40. In  $\Delta ABC$ , angle  $A$  is  $120^\circ$ ,  $BC + CA = 20$  and  $AB + BC = 21$ , then the length of the side  $BC$ , equals :  
 (a) 13 (b) 15 (c) 17 (d) 19
41. A triangle has sides 6, 7, 8. The line through its incentre parallel to the shortest side is drawn to meet the other two sides at  $P$  and  $Q$ . The length of the segment  $PQ$  is :  
 (a)  $\frac{12}{5}$  (b)  $\frac{15}{4}$  (c)  $\frac{30}{7}$  (d)  $\frac{33}{9}$
42. The perimeter of a  $\Delta ABC$  is 48 cm and one side is 20 cm. Then remaining sides of  $\Delta ABC$  must be greater than :  
 (a) 8 cm (b) 9 cm (c) 12 cm (d) 4 cm
43. In an equilateral  $\Delta ABC$ , (where symbols used have usual meanings), then  $r$ ,  $R$  and  $r_1$  form :  
 (a) an A.P. (b) a G.P.  
 (c) an H.P. (d) neither an A.P., G.P. nor H.P.
44. The expression  $\frac{(a+b+c)(b+c-a)(c+a-b)(a+b-c)}{4b^2c^2}$  is equal to :  
 (a)  $\cos^2 A$  (b)  $\sin^2 A$  (c)  $\cos A \cos B \cos C$  (d)  $\sin A \sin B \sin C$   
 (where symbols used have usual meanings)
45. Circumradius of an isosceles  $\Delta ABC$  with  $\angle A = \angle B$  is 4 times its in radius, then  $\cos A$  is root of the equation :  
 (a)  $x^2 - x - 8 = 0$  (b)  $8x^2 - 8x + 1 = 0$  (c)  $x^2 - x - 4 = 0$  (d)  $4x^2 - 4x + 1 = 0$
46.  $A$  is the orthocentre of  $\Delta ABC$  and  $D$  is reflection point of  $A$  w.r.t. perpendicular bisector of  $BC$ , then orthocenter of  $\Delta DBC$  is :  
 (a)  $D$  (b)  $C$  (c)  $B$  (d)  $A$
47. If  $a, b, c$  are sides of a scalene triangle, then the value of determinant  $\begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}$  is always:  
 (a)  $\geq 0$  (b)  $> 0$  (c)  $\leq -1$  (d)  $< 0$
48. In a triangle  $ABC$  if  $A : B : C = 1 : 2 : 4$ , then  $(a^2 - b^2)(b^2 - c^2)(c^2 - a^2) = \lambda a^2 b^2 c^2$ , where  $\lambda =$  :

- (a) 1                      (b) 2                      (c) 3                      (d)  $\frac{1}{3}$

49. The minimum value of  $\frac{r_1 r_2 r_3}{r^3}$  in a triangle is (symbols have their usual meaning)

- (a) 1                      (b) 3                      (c) 8                      (d) 27

50. In a triangle  $ABC$ ,  $BC = 3$ ,  $AC = 4$  and  $AB = 5$ . The value of  $\sin A + \sin 2B + \sin 3C$  equals

- (a)  $\frac{24}{25}$                       (b)  $\frac{14}{25}$                       (c)  $\frac{64}{25}$                       (d) None

51. In any triangle  $ABC$ , the value of  $\frac{r_1 + r_2}{1 + \cos C}$  is equal to (where notation have their usual meaning) :

- (a)  $2R$                       (b)  $2r$                       (c)  $R$                       (d)  $\frac{2R^2}{r}$

52. In a triangle  $ABC$ , medians  $AD$  and  $BE$  are drawn. If  $AD = 4$ ;  $\angle DAB = \frac{\pi}{6}$  and  $\angle ABE = \frac{\pi}{3}$  then the area of the triangle  $ABC$  is :

- (a)  $\frac{8}{3\sqrt{3}}$                       (b)  $\frac{16}{3\sqrt{3}}$                       (c)  $\frac{32}{3\sqrt{3}}$                       (d)  $\frac{64}{3\sqrt{3}}$

53. The sides of a triangle are  $\sin \alpha$ ,  $\cos \alpha$ ,  $\sqrt{1 + \sin \alpha \cos \alpha}$  for some  $0 < \alpha < \frac{\pi}{2}$  then the greatest angle of the triangle is :

- (a)  $\frac{\pi}{3}$                       (b)  $\frac{\pi}{2}$                       (c)  $\frac{2\pi}{3}$                       (d)  $\frac{5\pi}{6}$


54. Let  $ABC$  be a right triangle with  $\angle BAC = \frac{\pi}{2}$ , then  $\left(\frac{r^2}{2R^2} + \frac{r}{R}\right)$  is equal to :

(where symbols used have usual meaning in a triangle)

- (a)  $\sin B \sin C$                       (b)  $\tan B \tan C$                       (c)  $\sec B \sec C$                       (d)  $\cot B \cot C$

55. Find the radius of the circle escribed to the triangle  $ABC$  (Shown in the figure below) on the side  $BC$  if  $\angle NAB = 30^\circ$ ;  $\angle BAC = 30^\circ$ ;  $AB = AC = 5$ .

- (a)  $\frac{(10\sqrt{2} + 5\sqrt{3} - 5)(2 - \sqrt{3})}{2\sqrt{2}}$   
 (b)  $\frac{(10\sqrt{2} + 5\sqrt{3} + 5)(2 - \sqrt{3})}{2\sqrt{2}}$   
 (c)  $\frac{(10\sqrt{2} + 5\sqrt{3} - 5)(2 + \sqrt{3})}{2\sqrt{2}}$   
 (d)  $\frac{(10\sqrt{2} + 5\sqrt{2} + 1)(\sqrt{3} - 1)}{2\sqrt{3}}$





56. In a  $\triangle ABC$ , with usual notations, if  $b > c$  then distance between foot of median and foot of altitude both drawn from vertex  $A$  on  $BC$  is :

- (a)  $\frac{a^2 - b^2}{2c}$       (b)  $\frac{b^2 - c^2}{2a}$       (c)  $\frac{b^2 + c^2 - a^2}{2a}$       (d)  $\frac{b^2 + c^2 - a^2}{2c}$


57. In a triangle  $ABC$  the expression  $a \cos B \cos C + b \cos C \cos A + c \cos A \cos B$  equals to :

- (a)  $\frac{rs}{R}$       (b)  $\frac{r}{sR}$       (c)  $\frac{R}{rs}$       (d)  $\frac{Rs}{r}$

58. In an acute triangle  $ABC$ , altitudes from the vertices  $A, B$  and  $C$  meet the opposite sides at the points  $D, E$  and  $F$  respectively. If the radius of the circumcircle of  $\triangle AFE, \triangle BFD, \triangle CED, \triangle ABC$  be respectively  $R_1, R_2, R_3$  and  $R$ . Then the maximum value of  $R_1 + R_2 + R_3$  is :

- (a)  $\frac{3R}{8}$       (b)  $\frac{2R}{3}$       (c)  $\frac{4R}{3}$       (d)  $\frac{3R}{2}$


59. A circle of area 20 sq. units is centered at the point  $O$ . Suppose  $\triangle ABC$  is inscribed in that circle and has area 8 sq. units. The central angles  $\alpha, \beta$  and  $\gamma$  are as shown in the figure. The value of  $(\sin \alpha + \sin \beta + \sin \gamma)$  is equal to :



- (a)  $\frac{4\pi}{5}$       (b)  $\frac{3\pi}{4}$       (c)  $\frac{2\pi}{5}$       (d)  $\frac{\pi}{4}$

## Answers

1.	(d)	2.	(b)	3.	(b)	4.	(c)	5.	(b)	6.	(a)	7.	(d)	8.	(a)	9.	(a)	10.	(c)
11.	(c)	12.	(a)	13.	(a)	14.	(b)	15.	(a)	16.	(a)	17.	(c)	18.	(a)	19.	(a)	20.	(a)
21.	(c)	22.	(c)	23.	(b)	24.	(b)	25.	(c)	26.	(c)	27.	(c)	28.	(c)	29.	(b)	30.	(a)
31.	(b)	32.	(d)	33.	(b)	34.	(c)	35.	(d)	36.	(a)	37.	(d)	38.	(b)	39.	(d)	40.	(a)
41.	(c)	42.	(d)	43.	(a)	44.	(b)	45.	(b)	46.	(a)	47.	(d)	48.	(a)	49.	(d)	50.	(b)
51.	(a)	52.	(c)	53.	(c)	54.	(a)	55.	(a)	56.	(b)	57.	(a)	58.	(d)	59.	(a)		


**Exercise-2 : One or More than One Answer is/are Correct**

1. If  $r_1, r_2, r_3$  are radii of the escribed circles of a triangle  $ABC$  and  $r$  is the radius of its incircle, then the root(s) of the equation  $x^2 - r(r_1 r_2 + r_2 r_3 + r_3 r_1)x + (r_1 r_2 r_3 - 1) = 0$  is/are :
- (a)  $r_1$  (b)  $r_2 + r_3$  (c) 1 (d)  $r_1 r_2 r_3 - 1$
2. In  $\triangle ABC$ ,  $\angle A = 60^\circ$ ,  $\angle B = 90^\circ$ ,  $\angle C = 30^\circ$ . Let  $H$  be its orthocentre, then :  
(where symbols used have usual meanings)
- (a)  $AH = c$  (b)  $CH = a$  (c)  $AH = a$  (d)  $BH = 0$
3. In an equilateral triangle, if inradius is a rational number then which of the following is/are correct ?
- (a) circumradius is always rational (b) exradii are always rational  
(c) area is always ir-rational (d) perimeter is always rational
4. Let  $A, B, C$  be angles of a triangle  $ABC$  and let  $D = \frac{5\pi + A}{32}$ ,  $E = \frac{5\pi + B}{32}$ ,  $F = \frac{5\pi + C}{32}$ , then :  
(where  $D, E, F \neq \frac{n\pi}{2}$ ,  $n \in I$ ,  $I$  denote set of integers)
- (a)  $\cot D \cot E + \cot E \cot F + \cot D \cot F = 1$  (b)  $\cot D + \cot E + \cot F = \cot D \cot E \cot F$   
(c)  $\tan D \tan E + \tan E \tan F + \tan F \tan D = 1$  (d)  $\tan D + \tan E + \tan F = \tan D \tan E \tan F$
5. In a triangle  $ABC$ , if  $a = 4$ ,  $b = 8$  and  $\angle C = 60^\circ$ , then :  
(where symbols used have usual meanings)
- (a)  $c = 6$  (b)  $c = 4\sqrt{3}$  (c)  $\angle A = 30^\circ$  (d)  $\angle B = 90^\circ$
6. In a  $\triangle ABC$  if  $\frac{r}{r_1} = \frac{r_2}{r_3}$ , then which of the following is/are true ?  
(where symbols used have usual meanings)
- (a)  $a^2 + b^2 + c^2 = 8R^2$  (b)  $\sin^2 A + \sin^2 B + \sin^2 C = 2$   
(c)  $a^2 + b^2 = c^2$  (d)  $\Delta = s(s + c)$
7.  $ABC$  is a triangle whose circumcentre, incentre and orthocentre are  $O, I$  and  $H$  respectively which lie inside the triangle, then :
- (a)  $\angle BOC = A$  (b)  $\angle BIC = \frac{\pi}{2} + \frac{A}{2}$   
(c)  $\angle BHC = \pi - A$  (d)  $\angle BHC = \pi - \frac{A}{2}$
8. In a triangle  $ABC$ ,  $\tan A$  and  $\tan B$  satisfy the inequality  $\sqrt{3}x^2 - 4x + \sqrt{3} < 0$ , then which of the following must be correct ?  
(where symbols used have usual meanings)
- (a)  $a^2 + b^2 - ab < c^2$  (b)  $a^2 + b^2 > c^2$   
(c)  $a^2 + b^2 + ab > c^2$  (d)  $a^2 + b^2 < c^2$

9. If in a  $\triangle ABC$ ;  $\angle C = \frac{\pi}{8}$ ;  $a = \sqrt{2}$ ;  $b = \sqrt{2 + \sqrt{2}}$  then the measure of  $\angle A$  can be :
- (a)  $45^\circ$  (b)  $135^\circ$  (c)  $30^\circ$  (d)  $150^\circ$
10. In triangle  $ABC$ ,  $a = 3$ ,  $b = 4$ ,  $c = 2$ . Point  $D$  and  $E$  trisect the side  $BC$ . If  $\angle DAE = \theta$ , then  $\cot^2 \theta$  is divisible by :
- (a) 2 (b) 3 (c) 5 (d) 7
11. In a  $\triangle ABC$  if  $3 \sin A + 4 \cos B = 6$ ;  $4 \sin B + 3 \cos A = 1$  then possible value(s) of  $\angle C$  be:
- (a)  $\frac{\pi}{4}$  (b)  $\frac{\pi}{6}$  (c)  $\frac{\pi}{3}$  (d)  $\frac{5\pi}{6}$
12. If the line joining the incentre to the centroid of a triangle  $ABC$  is parallel to the side  $BC$ . Which of the following are correct ?
- (a)  $2b = a + c$  (b)  $2a = b + c$  (c)  $\cot \frac{A}{2} \cot \frac{C}{2} = 3$  (d)  $\cot \frac{B}{2} \cot \frac{C}{2} = 3$
13. In a triangle the length of two larger sides are 10 and 9 respectively. If the angles are in A.P., the length of third side can be :
- (a)  $5 - \sqrt{6}$  (b)  $5 + \sqrt{6}$  (c)  $6 - \sqrt{5}$  (d)  $6 + \sqrt{5}$
14. If area of  $\triangle ABC$ ,  $\Delta$  and angle  $C$  are given and if the side  $c$  opposite to given angle is minimum, then
- (a)  $a = \sqrt{\frac{2\Delta}{\sin C}}$  (b)  $b = \sqrt{\frac{2\Delta}{\sin C}}$  (c)  $a = \frac{4\Delta}{\sin C}$  (d)  $b = \frac{4\Delta}{\sin^2 C}$
15. In a triangle  $ABC$ , if  $\tan A = 2 \sin 2C$  and  $3 \cos A = 2 \sin B \sin C$  then possible values of  $C$  is/are
- (a)  $\frac{\pi}{8}$  (b)  $\frac{\pi}{6}$  (c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{3}$

### Answers

1.	(c, d)	2.	(a, b, d)	3.	(a, b, c)	4.	(b, c)	5.	(b, c, d)	6.	(a, b, c)
7.	(b, c)	8.	(a, c)	9.	(a)	10.	(b, c)	11.	(b)	12.	(b, d)
13.	(a, b)	14.	(a, b)	15.	(c, d)						

### Exercise-3 : Comprehension Type Problems

#### Paragraph for Question Nos. 1 to 2

Let  $\angle A = 23^\circ$ ,  $\angle B = 75^\circ$  and  $\angle C = 82^\circ$  be the angles of  $\triangle ABC$ .

The incircle of  $\triangle ABC$  touches the sides  $BC, CA, AB$  at points  $D, E, F$  respectively. Let  $r', r_1'$  respectively be the inradius, exradius opposite to vertex  $D$  of  $\triangle DEF$  and  $r$  be the inradius of  $\triangle ABC$ , then

1.  $\frac{r'}{r} =$

(a)  $\sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2} - 1$

(b)  $1 - \sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2}$

(c)  $\cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} - 1$

(d)  $1 - \cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2}$

2.  $\frac{r_1'}{r} =$

(a)  $\sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2} - 1$

(b)  $1 - \sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2}$

(c)  $\cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} - 1$

(d)  $1 - \cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2}$

#### Paragraph for Question Nos. 3 to 4

Internal angle bisectors of  $\triangle ABC$  meet its circumcircle at  $D, E$  and  $F$  where symbols have usual meaning.

3. Area of  $\triangle DEF$  is :

(a)  $2R^2 \cos^2\left(\frac{A}{2}\right) \cos^2\left(\frac{B}{2}\right) \cos^2\left(\frac{C}{2}\right)$

(b)  $2R^2 \sin\left(\frac{A}{2}\right) \sin\left(\frac{B}{2}\right) \sin\left(\frac{C}{2}\right)$

(c)  $2R^2 \sin^2\left(\frac{A}{2}\right) \sin^2\left(\frac{B}{2}\right) \sin^2\left(\frac{C}{2}\right)$

(d)  $2R^2 \cos\left(\frac{A}{2}\right) \cos\left(\frac{B}{2}\right) \cos\left(\frac{C}{2}\right)$

4. The ratio of area of triangle  $ABC$  and triangle  $DEF$  is :

(a)  $\geq 1$

(b)  $\leq 1$

(c)  $\geq 1/2$

(d)  $\leq 1/2$

#### Paragraph for Question Nos. 5 to 6

Let triangle  $ABC$  is right triangle right angled at  $C$  such that  $A < B$  and  $r = 8, R = 41$ .

5. Area of  $\triangle ABC$  is :

(a) 720

(b) 1440

(c) 360

(d) 480

6.  $\tan \frac{A}{2} =$

- (a)  $\frac{1}{18}$                       (b)  $\frac{1}{3}$                       (c)  $\frac{1}{6}$                       (d)  $\frac{1}{9}$

[where notations have their usual meaning]

### Paragraph for Question Nos. 7 to 8

Let the incircle of  $\Delta ABC$  touches the sides  $BC, CA, AB$  at  $A_1, B_1, C_1$  respectively. The incircle of  $\Delta A_1B_1C_1$  touches its sides of  $B_1C_1, C_1A_1$  and  $A_1B_1$  at  $A_2, B_2, C_2$  respectively and so on.

7.  $\lim_{n \rightarrow \infty} \angle A_n =$

- (a) 0                      (b)  $\frac{\pi}{6}$                       (c)  $\frac{\pi}{4}$                       (d)  $\frac{\pi}{3}$

8. In  $\Delta A_4B_4C_4$ , the value of  $\angle A_4$  is :

- (a)  $\frac{3\pi + A}{6}$                       (b)  $\frac{3\pi - A}{8}$                       (c)  $\frac{5\pi - A}{16}$                       (d)  $\frac{5\pi + A}{16}$

### Paragraph for Question Nos. 9 to 10

Let  $ABC$  be a given triangle. Points  $D$  and  $E$  are on sides  $AB$  and  $AC$  respectively and point  $F$  is on line segment  $DE$ . Let  $\frac{AD}{AB} = x, \frac{AE}{AC} = y, \frac{DF}{DE} = z$ . Let area of  $\Delta BDF = \Delta_1$ , area of  $\Delta CEF = \Delta_2$  and area of  $\Delta ABC = \Delta$ .

9.  $\frac{\Delta_1}{\Delta}$  is equal to :

- (a)  $xyz$                       (b)  $(1-x)y(1-z)$                       (c)  $(1-x)yz$                       (d)  $x(1-y)z$

10.  $\frac{\Delta_2}{\Delta}$  is equal to :

- (a)  $(1-x)y(1-z)$                       (b)  $(1-x)(1-y)z$                       (c)  $x(1-y)(1-z)$                       (d)  $(1-x)yz$

### Paragraph for Question Nos. 11 to 13

$a, b, c$  are the length of sides  $BC, CA, AB$  respectively of  $\Delta ABC$  satisfying  $\log\left(1 + \frac{c}{a}\right) + \log a - \log b = \log 2$ .

Also the quadratic equation  $a(1-x^2) + 2bx + c(1+x^2) = 0$  has two equal roots.

11.  $a, b, c$  are in :

- (a) A.P. (b) G.P. (c) H.P. (d) None

12. Measure of angle  $C$  is :

- (a)  $30^\circ$  (b)  $45^\circ$  (c)  $60^\circ$  (d)  $90^\circ$

13. The value of  $(\sin A + \sin B + \sin C)$  is equal to :

- (a)  $\frac{5}{2}$  (b)  $\frac{12}{5}$   
 (c)  $\frac{8}{3}$  (d) 2

**Paragraph for Question Nos. 14 to 16**

Let  $ABC$  be a triangle inscribed in a circle and let  $l_a = \frac{m_a}{M_a}$ ;  $l_b = \frac{m_b}{M_b}$ ;  $l_c = \frac{m_c}{M_c}$  where  $m_a, m_b, m_c$  are the lengths of the angle bisectors of angles  $A, B$  and  $C$  respectively, internal to the triangle and  $M_a, M_b$  and  $M_c$  are the lengths of these internal angle bisectors extended until they meet the circumcircle.

14.  $l_a$  equals :

- (a)  $\frac{\sin A}{\sin\left(B + \frac{A}{2}\right)}$  (b)  $\frac{\sin B \sin C}{\sin^2\left(\frac{B+C}{2}\right)}$  (c)  $\frac{\sin B \sin C}{\sin^2\left(B + \frac{A}{2}\right)}$  (d)  $\frac{\sin B + \sin C}{\sin^2\left(B + \frac{A}{2}\right)}$

15. The maximum value of the product  $(l_a l_b l_c) \times \cos^2\left(\frac{B-C}{2}\right) \times \cos^2\left(\frac{C-A}{2}\right) \times \cos^2\left(\frac{A-B}{2}\right)$  is equal to :

- (a)  $\frac{1}{8}$  (b)  $\frac{1}{64}$  (c)  $\frac{27}{64}$  (d)  $\frac{27}{32}$

16. The minimum value of the expression  $\frac{l_a}{\sin^2 A} + \frac{l_b}{\sin^2 B} + \frac{l_c}{\sin^2 C}$  is :

- (a) 2 (b) 3 (c) 4 (d) none of these

**Answers**

1.	(a)	2.	(b)	3.	(d)	4.	(b)	5.	(a)	6.	(d)	7.	(d)	8.	(d)	9.	(c)	10.	(c)
11.	(a)	12.	(d)	13.	(b)	14.	(c)	15.	(c)	16.	(b)								

### Exercise-4 : Matching Type Problems

1. Consider a right angled triangle **ABC** right angled at **C** with integer sides. List-I gives inradius. List-II gives the number of triangles.

	Column-I		Column-II
(A)	3	(P)	6
(B)	4	(Q)	7
(C)	6	(R)	8
(D)	9	(S)	10
		(T)	12

2.

	Column-I		Column-II
(A)	Find the sum of the series $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \frac{1}{9} + \frac{1}{12} + \dots \dots \infty$ , where the terms are the reciprocals of the positive integers whose only prime factors are two's and three's	(P)	7
(B)	The length of the sides of $\Delta ABC$ are $a, b$ and $c$ and $A$ is the angle opposite to side $a$ . If $b^2 + c^2 = a^2 + 54$ and $bc = \frac{a^3}{\cos A}$ then the value of $\left(\frac{b^2 + c^2}{9}\right)$ , is	(Q)	10
(C)	The equations of perpendicular bisectors of two sides $AB$ and $AC$ of a triangle $ABC$ are $x + y + 1 = 0$ and $x - y + 1 = 0$ respectively. If circumradius of $\Delta ABC$ is 2 units and the locus of vertex $A$ is $x^2 + y^2 + gx + c = 0$ , then $(g^2 + c^2)$ , is equal to	(R)	13
(D)	Number of solutions of the equation $\cos \theta \sin \theta + 6(\cos \theta - \sin \theta) + 6 = 0$ in $[0, 30]$ , is equal to	(S)	3

3. In  $\Delta ABC$ , if  $r_1 = 21, r_2 = 24, r_3 = 28$ , then

	Column-I		Column-II
(A)	$a =$	(P)	8
(B)	$b =$	(Q)	12
(C)	$s =$	(R)	26

(D)	$r =$	(S)	28
		(T)	42

(Where notations have their usual meaning)


4.

	Column-I		Column-II
(A)	$\frac{r_1(r_2 + r_3)}{\sqrt{r_2r_3 + r_3r_1 + r_1r_2}}$	(P)	$\sin \frac{A}{2}$
(B)	$\frac{r_1}{\sqrt{(r_1 + r_2)(r_1 + r_3)}}$	(Q)	$4R$
(C)	$r_1 + r_2 + r_3 - r$	(R)	$0$
(D)	$\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} - \frac{1}{r}$	(S)	$2R \sin A$

**Answers**


1.	A → P; B → P; C → T; D → S
2.	A → S; B → P; C → R; D → Q
3.	A → R; B → S; C → T; D → P
4.	A → S; B → P; C → Q; D → R




**Exercise-5 : Subjective Type Problems**

- If the median  $AD$  of  $\triangle ABC$  makes an angle  $\angle ADC = \frac{\pi}{4}$ . Find the value of  $|\cot B - \cot C|$ .
- In a  $\triangle ABC$ ,  $a = \sqrt{3}$ ,  $b = 3$  and  $\angle C = \frac{\pi}{3}$ . Let internal angle bisector of angle  $C$  intersects side  $AB$  at  $D$  and altitude from  $B$  meets the angle bisector  $CD$  at  $E$ . If  $O_1$  and  $O_2$  are incentres of  $\triangle BEC$  and  $\triangle BED$ . Find the distance between the vertex  $B$  and orthocentre of  $\triangle O_1EO_2$ .
- In a  $\triangle ABC$ ; inscribed circle with centre  $I$  touches sides  $AB, AC$  and  $BC$  at  $D, E, F$  respectively. Let area of quadrilateral  $ADIE$  is 5 units and area of quadrilateral  $BFID$  is 10 units. Find the value of  $\frac{\cos\left(\frac{C}{2}\right)}{\sin\left(\frac{A-B}{2}\right)}$ .
- If  $\Delta$  be area of incircle of a triangle  $ABC$  and  $\Delta_1, \Delta_2, \Delta_3$  be the area of excircles then find the least value of  $\frac{\Delta_1\Delta_2\Delta_3}{729\Delta^3}$ .
- In  $\triangle ABC$ ,  $b = c$ ,  $\angle A = 106^\circ$ ,  $M$  is an interior point such that  $\angle MBA = 7^\circ$ ,  $\angle MAB = 23^\circ$  and  $\angle MCA = \theta^\circ$ , then  $\frac{\theta}{2}$  is equal to  
(where notations have their usual meaning)
- In an acute angled triangle  $ABC$ ,  $\angle A = 20^\circ$ , let  $DEF$  be the feet of altitudes through  $A, B, C$  respectively and  $H$  is the orthocentre of  $\triangle ABC$ . Find  $\frac{AH}{AD} + \frac{BH}{BE} + \frac{CH}{CF}$ .
- Let  $\triangle ABC$  be inscribed in a circle having radius unity. The three internal bisectors of the angles  $A, B$  and  $C$  are extended to intersect the circumcircle of  $\triangle ABC$  at  $A_1, B_1$  and  $C_1$  respectively.  
Then  $\frac{AA_1 \cos \frac{A}{2} + BB_1 \cos \frac{B}{2} + CC_1 \cos \frac{C}{2}}{\sin A + \sin B + \sin C} =$
- If the quadratic equation  $ax^2 + bx + c = 0$  has equal roots where  $a, b, c$  denotes the lengths of the sides opposite to vertex  $A, B$  and  $C$  of the  $\triangle ABC$  respectively. Find the number of integers in the range of  $\frac{\sin A}{\sin C} + \frac{\sin C}{\sin A}$ .
- If in the triangle  $ABC$ ,  $\tan \frac{A}{2}$ ,  $\tan \frac{B}{2}$  and  $\tan \frac{C}{2}$  are in harmonic progression then the least value of  $\cot^2 \frac{B}{2}$  is equal to :
- In  $\triangle ABC$ , if circumradius ' $R$ ' and inradius ' $r$ ' are connected by relation  $R^2 - 4Rr + 8r^2 - 12r + 9 = 0$ , then the greatest integer which is less than the semiperimeter of  $\triangle ABC$  is :

11. Sides  $AB$  and  $AC$  in an equilateral triangle  $ABC$  with side length 3 is extended to form two rays from point  $A$  as shown in the figure. Point  $P$  is chosen outside the triangle  $ABC$  and between the two rays such that  $\angle ABP + \angle BCP = 180^\circ$ . If the maximum length of  $CP$  is  $M$ , then  $M^2/2$  is equal to :



12. Let  $a, b, c$  be sides of a triangle  $ABC$  and  $\Delta$  denotes its area.  
 If  $a = 2; \Delta = \sqrt{3}$  and  $a \cos C + \sqrt{3} a \sin C - b - c = 0$ ; then find the value of  $(b + c)$ .  
 (symbols used have usual meaning in  $\Delta ABC$ ).
13. If circumradius of  $\Delta ABC$  is 3 units and its area is 6 units and  $\Delta DEF$  is formed by joining foot of perpendiculars drawn from  $A, B, C$  on sides  $BC, CA, AB$  respectively. Find the perimeter of  $\Delta DEF$ .

Answers													
1.	2	2.	1	3.	3	4.	1	5.	7	6.	2	7.	2
8.	3	9.	3	10.	7	11.	6	12.	4	13.	4		





8. Number of solution(s) of the equation  $2 \tan^{-1}(2x - 1) = \cos^{-1}(x)$  is :  
 (a) 1 (b) 2 (c) 3 (d) infinitely many
9.  $\sin^{-1}\left(\frac{x^2}{4} + \frac{y^2}{9}\right) + \cos^{-1}\left(\frac{x}{2\sqrt{2}} + \frac{y}{3\sqrt{2}} - 2\right)$  equals to :  
 (a)  $\frac{\pi}{2}$  (b)  $\pi$  (c)  $\frac{\pi}{\sqrt{2}}$  (d)  $\frac{3\pi}{2}$
10. The complete solution set of the inequality  $(\cos^{-1} x)^2 - (\sin^{-1} x)^2 > 0$  is :  
 (a)  $\left[0, \frac{1}{\sqrt{2}}\right)$  (b)  $\left[-1, \frac{1}{\sqrt{2}}\right)$  (c)  $(-1, 1)$  (d)  $\left[-1, \frac{1}{2}\right)$
11. Let  $\alpha, \beta$  are the roots of the equation  $x^2 + 7x + k(k - 3) = 0$ , where  $k \in (0, 3)$  and  $k$  is a constant. Then the value of  $\tan^{-1} \alpha + \tan^{-1} \beta + \tan^{-1} \frac{1}{\alpha} + \tan^{-1} \frac{1}{\beta}$  is :  
 (a)  $\pi$  (b)  $\frac{\pi}{2}$  (c) 0 (d)  $-\frac{\pi}{2}$
12. Let  $f(x) = a + 2b \cos^{-1} x$ ,  $b > 0$ . If domain and range of  $f(x)$  are the same set, then  $(b - a)$  is equal to :  
 (a)  $1 - \frac{1}{\pi}$  (b)  $\frac{2}{\pi}$   
 (c)  $\frac{2}{\pi} + 1$  (d)  $1 + \frac{1}{\pi}$
13. If  $(\tan^{-1} x)^2 + (\cot^{-1} x)^2 = \frac{5\pi^2}{8}$ , then  $x$  equals to :  
 (a) -1 (b) 1 (c) 0 (d)  $\sqrt{3}$
14. The total number of ordered pairs  $(x, y)$  satisfying  $|y| = \cos x$  and  $y = \sin^{-1}(\sin x)$ , where  $x \in [-2\pi, 3\pi]$  is equal to :  
 (a) 2 (b) 4 (c) 5 (d) 6
15. If  $[\sin^{-1}(\cos^{-1}(\sin^{-1}(\tan^{-1} x)))] = 1$ , where  $[\cdot]$  denotes greatest integer function, then complete set of values of  $x$  is :  
 (a)  $[\tan(\sin(\cos 1)), \tan(\cos(\sin 1))]$  (b)  $[\tan(\sin(\cos 1)), \tan(\sin(\cos(\sin 1)))]$   
 (c)  $[\tan(\cos(\sin 1)), \tan(\sin(\cos(\sin 1)))]$  (d)  $[\tan(\sin(\cos 1)), 1]$
16. The number of ordered pair(s)  $(x, y)$  of real numbers satisfying the equation  $1 + x^2 + 2x \sin(\cos^{-1} y) = 0$ , is :  
 (a) 0 (b) 1 (c) 2 (d) 3
17. The value of  $\tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3$  is :  
 (a)  $\frac{\pi}{2}$  (b)  $\pi$  (c)  $\frac{3\pi}{4}$  (d)  $\frac{5\pi}{8}$

18. The complete set of values of  $x$  for which  $2 \tan^{-1} x + \cos^{-1} \left( \frac{1-x^2}{1+x^2} \right)$  is independent of  $x$  is :
- (a)  $(-\infty, 0]$                       (b)  $[0, \infty)$                       (c)  $(-\infty, -1]$                       (d)  $[1, \infty)$
19. The number of ordered pair(s)  $(x, y)$  which satisfy  $y = \tan^{-1} \tan x$  and  $16(x^2 + y^2) - 48\pi x + 16\pi y + 31\pi^2 = 0$ , is :
- (a) 0                                      (b) 1                                      (c) 2                                      (d) 3
20. Domain ( $D$ ) and range ( $R$ ) of  $f(x) = \sin^{-1}(\cos^{-1}[x])$  where  $[ ]$  denotes the greatest integer function is
- (a)  $D \equiv [1, 2), R \equiv \{0\}$                                       (b)  $D \equiv [0, 1), R \equiv \{-1, 0, 1\}$   
(c)  $D \equiv [-1, 1), R \equiv \left\{0, \frac{\pi}{2}, \pi\right\}$                                       (d)  $D \equiv [-1, 1), R \equiv \left\{-\frac{\pi}{2}, 0, \frac{\pi}{2}\right\}$
21. If  $2 \sin^{-1} x + \{\cos^{-1} x\} > \frac{\pi}{2} + \{\sin^{-1} x\}$ , then  $x \in$ : (where  $\{\cdot\}$  denotes fractional part function)
- (a)  $(\cos 1, 1]$                       (b)  $[\sin 1, 1]$                       (c)  $(\sin 1, 1]$                       (d) None of these
22. Let  $f(x) = x^{11} + x^9 - x^7 + x^3 + 1$  and  $f(\sin^{-1}(\sin 8)) = \alpha$ , ( $\alpha$  is constant). If  $f(\tan^{-1}(\tan 8)) = \lambda - \alpha$ , then the value of  $\lambda$  is :
- (a) 2                                      (b) 3                                      (c) 4                                      (d) 1
23. The number of real values of  $x$  satisfying the equation  $3 \sin^{-1} x + \pi x - \pi = 0$  is/are :
- (a) 0                                      (b) 1                                      (c) 2                                      (d) 3
24. Range of  $f(x) = \sin^{-1} x + x^2 + 4x + 1$  is :
- (a)  $\left[-\frac{\pi}{2} - 2, \frac{\pi}{2} + 6\right]$                       (b)  $\left[0, \frac{\pi}{2} + 6\right]$                       (c)  $\left[-\frac{\pi}{2} - 2, \infty\right)$                       (d)  $[-3, \infty)$
25. The solution set of the inequality  $(\operatorname{cosec}^{-1} x)^2 - 2 \operatorname{cosec}^{-1} x \geq \frac{\pi}{6}(\operatorname{cosec}^{-1} x - 2)$  is  $(-\infty, a] \cup [b, \infty)$ , then  $(a + b)$  equals :
- (a) 0                                      (b) 1                                      (c) 2                                      (d) -3
26. Number of solution of the equation  $2 \sin^{-1}(x + 2) = \cos^{-1}(x + 3)$  is :
- (a) 0                                      (b) 1                                      (c) 2                                      (d) None of these
27.  $\tan^{-1} \left( \frac{1}{3} \right) + \tan^{-1} \left( \frac{1}{7} \right) + \tan^{-1} \left( \frac{1}{13} \right) + \dots =$
- (a)  $\frac{\pi}{4}$                                       (b)  $\frac{\pi}{2}$                                       (c)  $\frac{\pi}{3}$                                       (d)  $\frac{\pi}{6}$
28. If  $\tan^{-1} \frac{1}{4} + \tan^{-1} \frac{2}{9} = \frac{1}{2} \cos^{-1} x$  then  $x$  is equal to :
- (a)  $\frac{1}{2}$                                       (b)  $\frac{2}{5}$                                       (c)  $\frac{3}{5}$                                       (d) none of these

29. The set of value of  $x$ , satisfying the equation  $\tan^2(\sin^{-1} x) > 1$  is :

- (a)  $(-1, 1)$  (b)  $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$   
 (c)  $[-1, 1] - \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  (d)  $(-1, 1) - \left[-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right]$

30. The sum of the series  $\cot^{-1}\left(\frac{9}{2}\right) + \cot^{-1}\left(\frac{33}{4}\right) + \cot^{-1}\left(\frac{129}{8}\right) + \dots \infty$  is equal to :

- (a)  $\cot^{-1}(2)$  (b)  $\cot^{-1}(3)$  (c)  $\cot^{-1}(-1)$  (d)  $\cot^{-1}(1)$

31. If  $\int \frac{\ln(\cot x)}{\sin x \cos x} dx = -\frac{1}{k} \ln^2(\cot x) + C$

(where  $C$  is a constant); then the value of  $k$  is :

- (a) 1 (b) 2 (c) 3 (d)  $\frac{1}{2}$

32. The number of solutions of  $\sin^{-1} x + \sin^{-1}(1+x) = \cos^{-1} x$  is/are :

- (a) 0 (b) 1 (c) 2 (d) infinite

33. The value of  $x$  satisfying the equation

$$(\sin^{-1} x)^3 - (\cos^{-1} x)^3 + (\sin^{-1} x)(\sin^{-1} x - \cos^{-1} x) = \frac{\pi^3}{16} \text{ is :}$$

- (a)  $\cos \frac{\pi}{5}$  (b)  $\cos \frac{\pi}{4}$  (c)  $\cos \frac{\pi}{8}$  (d)  $\cos \frac{\pi}{12}$

34. The complete solution set of the equation

$$\sin^{-1} \sqrt{\frac{1+x}{2}} - \sqrt{2-x} = \cot^{-1}(\tan \sqrt{2-x}) - \sin^{-1} \sqrt{\frac{1-x}{2}} \text{ is :}$$

- (a)  $\left[2 - \frac{\pi^2}{4}, 1\right]$  (b)  $\left[1 - \frac{\pi^2}{4}, 1\right]$  (c)  $\left[2 - \frac{\pi^2}{4}, 0\right]$  (d)  $[-1, 1]$

35. Let  $f(x) = \tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$  then which of the following is correct :

- (a)  $f(x)$  has only one integer in its range (b) Range of  $f(x)$  is  $\left(-\frac{\pi}{4}, \frac{\pi}{4}\right) - \{0\}$   
 (c) Range of  $f(x)$  is  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) - \{0\}$  (d) Range of  $f(x)$  is  $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right] - \{0\}$

36. If  $\tan^{-1} \frac{1}{4} + \tan^{-1} \frac{2}{9} = \frac{1}{2} \cos^{-1} x$  then  $x$  is equal to :

- (a)  $\frac{1}{2}$  (b)  $\frac{2}{5}$  (c)  $\frac{3}{5}$  (d) None of these

37. The set of values of  $x$ , satisfying the equation  $\tan^2(\sin^{-1} x) > 1$  is :
- (a)  $(-1, 1)$  (b)  $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$   
 (c)  $[-1, 1] - \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  (d)  $(-1, 1) - \left[-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right]$
38. The sum of the series  $\cot^{-1}\left(\frac{9}{2}\right) + \cot^{-1}\left(\frac{33}{4}\right) + \cot^{-1}\left(\frac{129}{8}\right) + \dots \infty$  is equal to
- (a)  $\cot^{-1}(2)$  (b)  $\cot^{-1}(3)$  (c)  $\cot^{-1}(-1)$  (d)  $\cot^{-1}(1)$
39. The number of real values of  $x$  satisfying  $\tan^{-1}\left(\frac{x}{1-x^2}\right) + \tan^{-1}\left(\frac{1}{x^3}\right) = \frac{3\pi}{4}$  is :
- (a) 0 (b) 1 (c) 2 (d) infinitely many
40. Number of integral values of  $\lambda$  such that the equation  $\cos^{-1} x + \cot^{-1} x = \lambda$  possesses solution is:
- (a) 2 (b) 8 (c) 5 (d) 10
41. If the equation  $x^3 + bx^2 + cx + 1 = 0$  ( $b < c$ ) has only one real root  $\alpha$ . Then the value of  $2 \tan^{-1}(\operatorname{cosec} \alpha) + \tan^{-1}(2 \sin \alpha \sec^2 \alpha)$  is :
- (a)  $-\frac{\pi}{2}$  (b)  $-\pi$  (c)  $\frac{\pi}{2}$  (d)  $\pi$
42. Range of the function  $f(x) = \cot^{-1}\{-x\} + \sin^{-1}\{x\} + \cos^{-1}\{x\}$ , where  $\{ \cdot \}$  denotes fractional part function
- (a)  $\left(\frac{3\pi}{4}, \pi\right)$  (b)  $\left[\frac{3\pi}{4}, \pi\right)$  (c)  $\left[\frac{3\pi}{4}, \pi\right]$  (d)  $\left(\frac{3\pi}{4}, \pi\right]$
43. If  $3 \leq a < 4$  then the value of  $\sin^{-1}(\sin[a]) + \tan^{-1}(\tan[a]) + \sec^{-1}(\sec[a])$ , where  $[x]$  denotes greatest integer function less than or equal to  $x$ , is equal to :
- (a) 3 (b)  $2\pi - 9$  (c)  $2\pi - 3$  (d)  $9 - 2\pi$
44. The number of real solutions of  $y + y^2 = \sin x$  and  $y + y^3 = \cos^{-1} \cos x$  is/are
- (a) 0 (b) 1 (c) 3 (d) Infinite
45. Range of  $f(x) = \sin^{-1}[x-1] + 2 \cos^{-1}[x-2]$  ( $[ \cdot ]$  denotes greatest integer function)
- (a)  $\left\{-\frac{\pi}{2}, 0\right\}$  (b)  $\left\{\frac{\pi}{2}, 2\pi\right\}$  (c)  $\left\{\frac{\pi}{4}, \frac{\pi}{2}\right\}$  (d)  $\left\{\frac{3\pi}{2}, 2\pi\right\}$






### Exercise-2 : One or More than One Answer is/are Correct

1.  $f(x) = \sin^{-1}(\sin x)$ ,  $g(x) = \cos^{-1}(\cos x)$ , then :
- (a)  $f(x) = g(x)$  if  $x \in \left(0, \frac{\pi}{4}\right)$                       (b)  $f(x) < g(x)$  if  $x \in \left(\frac{\pi}{2}, \frac{3\pi}{4}\right)$   
 (c)  $f(x) < g(x)$  if  $\left(\pi, \frac{5\pi}{4}\right)$                       (d)  $f(x) > g(x)$  if  $x \in \left(\pi, \frac{5\pi}{4}\right)$
2. The solution(s) of the equation  $\cos^{-1} x = \tan^{-1} x$  satisfy
- (a)  $x^2 = \frac{\sqrt{5}-1}{2}$                       (b)  $x^2 = \frac{\sqrt{5}+1}{2}$   
 (c)  $\sin(\cos^{-1} x) = \frac{\sqrt{5}-1}{2}$                       (d)  $\tan(\cos^{-1} x) = \frac{\sqrt{5}-1}{2}$
3. If the numerical value of  $\tan\left(\cos^{-1}\left(\frac{4}{5}\right) + \tan^{-1}\left(\frac{2}{3}\right)\right)$  is  $\left(\frac{a}{b}\right)$ , where  $a, b$  are two positive integers and their H.C.F. is 1
- (a)  $a + b = 23$                       (b)  $a - b = 11$                       (c)  $3b = a + 1$                       (d)  $2a = 3b$
4. A solution of the equation  $\cot^{-1} 2 = \cot^{-1} x + \cot^{-1}(10 - x)$  where  $1 < x < 9$  is :
- (a) 7                      (b) 3                      (c) 2                      (d) 5
5. Consider the equation  $\sin^{-1}\left(x^2 - 6x + \frac{17}{2}\right) + \cos^{-1} k = \frac{\pi}{2}$ , then :
- (a) the largest value of  $k$  for which equation has 2 distinct solution is 1  
 (b) the equation must have real root if  $k \in \left(-\frac{1}{2}, 1\right)$   
 (c) the equation must have real root if  $k \in \left(-1, \frac{1}{2}\right)$   
 (d) the equation has unique solution if  $k = -\frac{1}{2}$
6. The value of  $x$  satisfying the equation
- $$(\sin^{-1} x)^3 - (\cos^{-1} x)^3 + (\sin^{-1} x)(\cos^{-1} x)(\sin^{-1} x - \cos^{-1} x) = \frac{\pi^3}{16}$$
- can not be equal to :
- (a)  $\cos \frac{\pi}{5}$                       (b)  $\cos \frac{\pi}{4}$                       (c)  $\cos \frac{\pi}{8}$                       (d)  $\cos \frac{\pi}{12}$

## Answers

1.	(a, b, c)	2.	(a, c)	3.	(a, b, c)	4.	(a, b)	5.	(a, b, d)	6.	(a, b, d)
----	-----------	----	--------	----	-----------	----	--------	----	-----------	----	-----------




**Exercise-4 : Matching Type Problems**

1.

Column-I		Column-II	
(A)	$\sin^{-1} \frac{4}{5} + 2 \tan^{-1} \frac{1}{3} =$	(P)	$\frac{\pi}{6}$
(B)	$\sin^{-1} \frac{12}{13} + \cos^{-1} \frac{4}{5} + \tan^{-1} \frac{63}{16} =$	(Q)	$\frac{\pi}{2}$
(C)	If $A = \tan^{-1} \frac{x\sqrt{3}}{2\lambda - x}$ , $B = \tan^{-1} \left( \frac{2x - \lambda}{\lambda\sqrt{3}} \right)$ then $A - B$ can be equal to	(R)	$\frac{\pi}{4}$
(D)	$\tan^{-1} \frac{1}{7} + 2 \tan^{-1} \frac{1}{3} =$	(S)	$\pi$
		(T)	$\frac{\pi}{3}$

2.

Column-I		Column-II	
(P)	If $f(x) = \sin^{-1} x$ and $\lim_{x \rightarrow \frac{1}{2}^+} f(3x - 4x^3)$ $= l - 3 \left( \lim_{x \rightarrow \frac{1}{2}^+} f(x) \right)$ then $[l] =$ ( $[ \ ]$ denotes greatest integer function)	(P)	3
(Q)	If $x > 1$ , then the value of $\sin \left( \frac{1}{2} \tan^{-1} \frac{2x}{1-x^2} - \tan^{-1} x \right)$ is	(Q)	-1
(R)	Number of values of $x$ satisfying $\sin^{-1} x - \cos^{-1} x = \sin^{-1} (3x - 2)$	(R)	2
(S)	The value of $\sin \left( \tan^{-1} 3 + \tan^{-1} \frac{1}{3} \right)$	(S)	1

3.


Column-I		Column-II	
(A)	If the first term of an arithmetic progression is 1, its second term is $n$ , and the sum of the first $n$ terms is $33n$	(P)	3
(B)	If the equation $\cos^{-1} x + \cot^{-1} x = k$ possess solution, then the largest integral value of $k$ is	(Q)	4
(C)	The number of solution of equation $\cos \theta =  1 + \sin \theta $ in interval $[0, 3\pi]$ , is	(R)	5
(D)	If the quadratic equation $x^2 - x - a = 0$ has integral roots where $a \in N$ and $4 \leq a \leq 40$ , then the number of possible values of $a$ is	(S)	9

4.

Column-I		Column-II	
(A)	The value of $\tan^{-1}([\pi]) + \tan^{-1}([- \pi] + 1) =$ ( $[ \cdot ]$ denotes greatest integer function)	(P)	2
(B)	The number of solutions of the equation $\tan x + \sec x = 2 \cos x$ in the interval $[0, 2\pi]$ is	(Q)	3
(C)	The number of roots of the equation $x + 2 \tan x = \frac{\pi}{2}$ in the interval $[0, 2\pi]$ is	(R)	0
(D)	The number of solutions of the equation $x^3 + x^2 + 4x + 2 \sin x = 0$ in the interval $[0, 2\pi]$ is	(S)	1

### Answers

1. A  $\rightarrow$  Q; B  $\rightarrow$  S; C  $\rightarrow$  P; D  $\rightarrow$  R
2. A  $\rightarrow$  P; B  $\rightarrow$  Q; C  $\rightarrow$  R; D  $\rightarrow$  S
3. A  $\rightarrow$  S; B  $\rightarrow$  R; C  $\rightarrow$  P; D  $\rightarrow$  Q
4. A  $\rightarrow$  R; B  $\rightarrow$  P; C  $\rightarrow$  Q; D  $\rightarrow$  S

 **Exercise-5 : Subjective Type Problems**

1. The complete set of values of  $x$  satisfying the inequality  $\sin^{-1}(\sin 5) > x^2 - 4x$  is  $(2 - \sqrt{\lambda - 2\pi}, 2 + \sqrt{\lambda - 2\pi})$ , then  $\lambda =$
2. In a  $\Delta ABC$ ; if  $(II_1)^2 + (I_2I_3)^2 = \lambda R^2$ , where  $I$  denotes incentre;  $I_1, I_2$  and  $I_3$  denote centres of the circles escribed to the sides  $BC, CA$  and  $AB$  respectively and  $R$  be the radius of the circum circle of  $\Delta ABC$ . Find  $\lambda$ .
3. If  $2 \tan^{-1} \frac{1}{5} - \sin^{-1} \frac{3}{5} = -\cos^{-1} \frac{63}{\lambda}$ , then  $\lambda =$
4. If  $2 \tan^{-1} \frac{1}{5} - \sin^{-1} \frac{3}{5} = -\cos^{-1} \frac{9\lambda}{65}$ , then  $\lambda =$
5. If  $\sum_{n=0}^{\infty} 2 \cot^{-1} \left( \frac{n^2 + n + 4}{2} \right) = k\pi$ , then find the value of  $k$ .
6. Find number of solutions of the equation  $\sin^{-1}(|\log_6^2(\cos x) - 1|) + \cos^{-1}(|3 \log_6^2(\cos x) - 7|) = \frac{\pi}{2}$ , if  $x \in [0, 4\pi]$ .

 **Answers**

1.	9	2.	16	3.	65	4.	7	5.	1	6.	4		
----	---	----	----	----	----	----	---	----	---	----	---	--	--

□□□

# Vector & 3Dimensional Geometry

---

26. Vector and 3Dimensional Geometry

# VECTOR & 3D DIMENSIONAL GEOMETRY

## Exercise-1 : Single Choice Problems

- The minimum value of  $x^2 + y^2 + z^2$  if  $ax + by + cz = p$ , is :
 

(a)  $\left(\frac{p}{a+b+c}\right)^2$       (b)  $\frac{p^2}{a^2+b^2+c^2}$       (c)  $\frac{a^2+b^2+c^2}{p^2}$       (d) 0
- If the angle between the vectors  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{3}$  and the area of the triangle with adjacent sides equal to  $\vec{a}$  and  $\vec{b}$  is 3, then  $\vec{a} \cdot \vec{b}$  is equal to :
 

(a)  $\sqrt{3}$       (b)  $2\sqrt{3}$       (c)  $4\sqrt{3}$       (d)  $\frac{\sqrt{3}}{2}$
- A straight line cuts the sides  $AB, AC$  and  $AD$  of a parallelogram  $ABCD$  at points  $B_1, C_1$  and  $D_1$  respectively. If  $\vec{AB}_1 = \lambda_1 \vec{AB}, \vec{AD}_1 = \lambda_2 \vec{AD}$  and  $\vec{AC}_1 = \frac{\lambda_3}{2} \vec{AC}$ , where  $\lambda_1, \lambda_2$  and  $\lambda_3$  are positive real numbers, then :
 

(a)  $\lambda_1, \lambda_3$  and  $\lambda_2$  are in AP      (b)  $\lambda_1, \lambda_3$  and  $\lambda_2$  are in GP  
 (c)  $\lambda_1, \lambda_3$  and  $\lambda_2$  are in HP      (d)  $\lambda_1 + \lambda_2 + \lambda_3 = 0$
- Let  $\vec{a} = 2\hat{i} + \hat{j} - 2\hat{k}$  and  $\vec{b} = \hat{i} + \hat{j}$ . If  $\vec{c}$  is a vector such that  $|\vec{a} \cdot \vec{c}| = |\vec{c}|, |\vec{c} - \vec{a}| = 2\sqrt{2}$  and the angle between  $\vec{a} \times \vec{b}$  and  $\vec{c}$  is  $30^\circ$  then  $\left|(\vec{a} \times \vec{b}) \times \vec{c}\right|$  is equal to :
 

(a)  $\frac{2}{3}$       (b)  $\frac{3}{2}$       (c) 2      (d) 3
- If acute angle between the line  $\vec{r} = \hat{i} + 2\hat{j} + \lambda(4\hat{i} - 3\hat{k})$  and  $xy$ -plane is  $\theta_1$  and acute angle between the planes  $x + 2y = 0$  and  $2x + y = 0$  is  $\theta_2$ , then  $(\cos^2 \theta_1 + \sin^2 \theta_2)$  equals to :
 

(a) 1      (b)  $\frac{1}{4}$       (c)  $\frac{2}{3}$       (d)  $\frac{3}{4}$

6. If  $a, b, c, x, y, z$  are real and  $a^2 + b^2 + c^2 = 25$ ,  $x^2 + y^2 + z^2 = 36$  and  $ax + by + cz = 30$ , then  $\frac{a+b+c}{x+y+z}$  is equal to :
- (a) 1 (b)  $\frac{6}{5}$  (c)  $\frac{5}{6}$  (d)  $\frac{3}{4}$
7. If  $\vec{a}$  and  $\vec{b}$  are non-zero, non-collinear vectors such that  $|\vec{a}| = 2$ ,  $\vec{a} \cdot \vec{b} = 1$  and angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{3}$ . If  $\vec{r}$  is any vector such that  $\vec{r} \cdot \vec{a} = 2$ ,  $\vec{r} \cdot \vec{b} = 8$ ,  $(\vec{r} + 2\vec{a} - 10\vec{b}) \cdot (\vec{a} \times \vec{b}) = 4\sqrt{3}$  and satisfy to  $\vec{r} + 2\vec{a} - 10\vec{b} = \lambda(\vec{a} \times \vec{b})$ , then  $\lambda$  is equal to :
- (a)  $\frac{1}{2}$  (b) 2 (c)  $\frac{1}{4}$  (d) None of these
8. Let  $\vec{a} = 3\hat{i} + 2\hat{j} + 4\hat{k}$ ;  $\vec{b} = 2(\hat{i} + \hat{k})$  and  $\vec{c} = 4\hat{i} + 2\hat{j} + 3\hat{k}$ . Sum of the values of ' $\alpha$ ' for which the equation  $x\vec{a} + y\vec{b} + z\vec{c} = \alpha(x\hat{i} + y\hat{j} + z\hat{k})$  has non-trivial solution is :
- (a) -1 (b) 4 (c) 7 (d) 8
9. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$ ,  $\vec{c} = \hat{i} + 2\hat{j} - \hat{k}$ , then the value of  $\begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot \vec{a} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \\ \vec{c} \cdot \vec{a} & \vec{c} \cdot \vec{b} & \vec{c} \cdot \vec{c} \end{vmatrix}$  is equal to :
- (a) 2 (b) 4 (c) 16 (d) 64
10. If  $\vec{a}$  and  $\vec{b}$  are two vectors such that  $|\vec{a}| = 1$ ,  $|\vec{b}| = 4$ ,  $\vec{a} \cdot \vec{b} = 2$ . If  $\vec{c} = (2\vec{a} \times \vec{b}) - 3\vec{b}$ , then angle between  $\vec{b}$  and  $\vec{c}$  is :
- (a)  $\frac{\pi}{6}$  (b)  $\frac{\pi}{3}$  (c)  $\frac{2\pi}{3}$  (d)  $\frac{5\pi}{6}$
11. If  $\vec{a}, \vec{b}, \vec{c}$  are unit vectors, then the value of  $|\vec{a} - 2\vec{b}|^2 + |\vec{b} - 2\vec{c}|^2 + |\vec{c} - 2\vec{a}|^2$  does not exceed to:
- (a) 9 (b) 12 (c) 18 (d) 21
12. The adjacent side vectors  $\vec{OA}$  and  $\vec{OB}$  of a rectangle  $OACB$  are  $\vec{a}$  and  $\vec{b}$  respectively, where  $O$  is the origin. If  $16|\vec{a} \times \vec{b}| = 3(|\vec{a}| + |\vec{b}|)^2$  and  $\theta$  be the acute angle between the diagonals  $OC$  and  $AB$  then the value of  $\tan(\theta/2)$  is :
- (a)  $\frac{1}{\sqrt{2}}$  (b)  $\frac{1}{2}$  (c)  $\frac{1}{\sqrt{3}}$  (d)  $\frac{1}{3}$
13. The vector  $\vec{AB} = 3\hat{i} + 4\hat{k}$  and  $\vec{AC} = 5\hat{i} - 2\hat{j} + 4\hat{k}$  are the sides of a triangle  $ABC$ . The length of the median through  $A$  is :
- (a)  $\sqrt{288}$  (b)  $\sqrt{72}$  (c)  $\sqrt{33}$  (d)  $\sqrt{18}$



14. If  $\vec{a} = 2\hat{i} + \lambda\hat{j} + 3\hat{k}$ ;  $\vec{b} = 3\hat{i} + 3\hat{j} + 5\hat{k}$ ;  $\vec{c} = \lambda\hat{i} + 2\hat{j} + 2\hat{k}$  are linearly dependent vectors, then the number of possible values of  $\lambda$  is :  
 (a) 0 (b) 1 (c) 2 (d) More than 2
15. The scalar triple product  $[\vec{a} + \vec{b} - \vec{c} \quad \vec{b} + \vec{c} - \vec{a} \quad \vec{c} + \vec{a} - \vec{b}]$  is equal to :  
 (a) 0 (b)  $[\vec{a} \vec{b} \vec{c}]$  (c)  $2[\vec{a} \vec{b} \vec{c}]$  (d)  $4[\vec{a} \vec{b} \vec{c}]$
16. If  $\hat{a}$  and  $\hat{b}$  are unit vectors then the vector defined as  $\vec{V} = (\hat{a} \times \hat{b}) \times (\hat{a} + \hat{b})$  is collinear to the vector :  
 (a)  $\hat{a} + \hat{b}$  (b)  $\hat{b} - \hat{a}$  (c)  $2\hat{a} - \hat{b}$  (d)  $\hat{a} + 2\hat{b}$
17. The sine of angle formed by the lateral face  $ADC$  and plane of the base  $ABC$  of the tetrahedron  $ABCD$ , where  $A = (3, -2, 1)$ ;  $B = (3, 1, 5)$ ;  $C = (4, 0, 3)$  and  $D = (1, 0, 0)$ , is :  
 (a)  $\frac{2}{\sqrt{29}}$  (b)  $\frac{5}{\sqrt{29}}$  (c)  $\frac{3\sqrt{3}}{\sqrt{29}}$  (d)  $\frac{-2}{\sqrt{29}}$
18. Let  $\vec{a}_r = x_r\hat{i} + y_r\hat{j} + z_r\hat{k}$ ,  $r = 1, 2, 3$  be three mutually perpendicular unit vectors, then the value of  $\begin{vmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{vmatrix}$  is equal to :  
 (a) 0 (b)  $\pm 1$  (c)  $\pm 2$  (d)  $\pm 4$
19. Let  $\vec{a}, \vec{b}, \vec{c}$  be three non-coplanar vectors and  $\vec{r}$  be any arbitrary vector, then the expression  $(\vec{a} \times \vec{b}) \times (\vec{r} \times \vec{c}) + (\vec{b} \times \vec{c}) \times (\vec{r} \times \vec{a}) + (\vec{c} \times \vec{a}) \times (\vec{r} \times \vec{b})$  is always equal to :  
 (a)  $[\vec{a} \vec{b} \vec{c}] \vec{r}$  (b)  $2[\vec{a} \vec{b} \vec{c}] \vec{r}$  (c)  $4[\vec{a} \vec{b} \vec{c}] \vec{r}$  (d)  $\vec{0}$
20.  $E$  and  $F$  are the interior points on the sides  $BC$  and  $CD$  of a parallelogram  $ABCD$ . Let  $\vec{BE} = 4\vec{EC}$  and  $\vec{CF} = 4\vec{FD}$ . If the line  $EF$  meets the diagonal  $AC$  in  $G$ , then  $\vec{AG} = \lambda\vec{AC}$ , where  $\lambda$  is equal to :  
 (a)  $\frac{1}{3}$  (b)  $\frac{21}{25}$  (c)  $\frac{7}{13}$  (d)  $\frac{21}{5}$
21. If  $\hat{a}, \hat{b}$  are unit vectors and  $\vec{c}$  is such that  $\vec{c} = \vec{a} \times \vec{c} + \vec{b}$ , then the maximum value of  $[\vec{a} \vec{b} \vec{c}]$  is :  
 (a) 1 (b)  $\frac{1}{2}$  (c) 2 (d)  $\frac{3}{2}$
22. Consider matrices  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 1 & 2 \\ 1 & -1 & 1 \end{bmatrix}$ ;  $B = \begin{bmatrix} 2 & 1 & 3 \\ 4 & 1 & -1 \\ 2 & 2 & 3 \end{bmatrix}$ ;  $C = \begin{bmatrix} 14 \\ 12 \\ 2 \end{bmatrix}$ ;  $D = \begin{bmatrix} 13 \\ 11 \\ 14 \end{bmatrix}$ ;  $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$  such that solutions of equation  $AX = C$  and  $BX = D$  represents two points  $P(x_1, y_1, z_1)$  and  $Q(x_2, y_2, z_2)$  respectively in three dimensional space. If  $P'Q'$  is the reflection of the line  $PQ$  in the plane  $\Pi: x + y + z = 9$ , then the point which does not lie on  $P'Q'$  is :  
 (a) (3, 4, 2) (b) (5, 3, 4) (c) (7, 2, 3) (d) (1, 5, 6)

23. The value of  $\alpha$  for which point  $M(\alpha\hat{i} + 2\hat{j} + \hat{k})$ , lies in the plane containing three points  $A(\hat{i} + \hat{j} + \hat{k})$ ,  $B(2\hat{i} + 2\hat{j} + \hat{k})$  and  $C(3\hat{i} - \hat{k})$  is :
- (a) 1 (b) 2 (c)  $\frac{1}{2}$  (d)  $-\frac{1}{2}$
24.  $Q$  is the image of point  $P(1, -2, 3)$  with respect to the plane  $x - y + z = 7$ . The distance of  $Q$  from the origin is :
- (a)  $\sqrt{\frac{70}{3}}$  (b)  $\frac{1}{2}\sqrt{\frac{70}{3}}$  (c)  $\sqrt{\frac{35}{3}}$  (d)  $\sqrt{\frac{15}{2}}$
25.  $\hat{a}$ ,  $\hat{b}$  and  $\hat{a} - \hat{b}$  are unit vectors. The volume of the parallelepiped, formed with  $\hat{a}$ ,  $\hat{b}$  and  $\hat{a} \times \hat{b}$  as coterminal edges is :
- (a) 1 (b)  $\frac{1}{4}$  (c)  $\frac{2}{3}$  (d)  $\frac{3}{4}$
26. A line passing through  $P(3, 7, 1)$  and  $R(2, 5, 7)$  meet the plane  $3x + 2y + 11z - 9 = 0$  at  $Q$ . Then  $PQ$  is equal to :
- (a)  $\frac{5\sqrt{41}}{59}$  (b)  $\frac{\sqrt{41}}{59}$  (c)  $\frac{50\sqrt{41}}{59}$  (d)  $\frac{25\sqrt{41}}{59}$
27. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are three non-zero non-coplanar vectors and  $\vec{p} = \vec{a} + \vec{b} - 2\vec{c}$ ;  $\vec{q} = 3\vec{a} - 2\vec{b} + \vec{c}$  and  $\vec{r} = \vec{a} - 4\vec{b} + 2\vec{c}$  are three vectors such that the volumes of the parallelepiped formed by  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  and  $\vec{p}$ ,  $\vec{q}$ ,  $\vec{r}$  as their coterminal edges are  $V_1$  and  $V_2$  respectively. Then  $\frac{V_2}{V_1}$  is equal to :
- (a) 10 (b) 15 (c) 20 (d) None of these
28. If the two lines represented by  $x + ay = b$ ;  $z + cy = d$  and  $x = a'y + b'$ ;  $z = c'y + d'$  be perpendicular to each other, then the value of  $aa' + cc'$  is :
- (a) 1 (b) 2 (c) 3 (d) 4
29. The distance between the line  $\vec{r} = 2\hat{i} - 2\hat{j} + 3\hat{k} + \lambda(\hat{i} - \hat{j} + 4\hat{k})$  and the plane  $\vec{r} \cdot (\hat{i} + 5\hat{j} + \hat{k}) = 5$  is :
- (a)  $\frac{10}{9}$  (b)  $\frac{10}{3\sqrt{3}}$  (c)  $\frac{3}{10}$  (d)  $\frac{10}{3}$
30. If  $(\vec{a} \times \vec{b}) \times \vec{c} = \vec{a} \times (\vec{b} \times \vec{c})$ , where  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are any three vectors such that  $\vec{a} \cdot \vec{b} \neq 0$ ,  $\vec{b} \cdot \vec{c} \neq 0$ , then  $\vec{a}$  and  $\vec{c}$  are :
- (a) Inclined at an angle of  $\frac{\pi}{3}$  (b) Inclined at an angle of  $\frac{\pi}{6}$   
 (c) Perpendicular (d) Parallel

31. Let  $\vec{r}$  be position vector of variable point in cartesian plane  $OXY$  such that  $\vec{r} \cdot (\vec{r} + 6\hat{j}) = 7$  cuts the co-ordinate axes at four distinct points, then the area of the quadrilateral formed by joining these points is :

- (a)  $4\sqrt{7}$  (b)  $6\sqrt{7}$  (c)  $7\sqrt{7}$  (d)  $8\sqrt{7}$

32. If  $|\vec{a}| = 2, |\vec{b}| = 5$  and  $\vec{a} \cdot \vec{b} = 0$ , then  $\vec{a} \times (\vec{a} \times (\vec{a} \times (\vec{a} \times (\vec{a} \times (\vec{a} \times \vec{b}))))$  is equal to :

- (a)  $64\vec{a}$  (b)  $64\vec{b}$  (c)  $-64\vec{a}$  (d)  $-64\vec{b}$

33. If  $O$  (origin) is a point inside the triangle  $PQR$  such that  $\vec{OP} + k_1 \vec{OQ} + k_2 \vec{OR} = 0$ , where  $k_1, k_2$  are constants such that  $\frac{\text{Area}(\Delta PQR)}{\text{Area}(\Delta OQR)} = 4$ , then the value of  $k_1 + k_2$  is :

- (a) 2 (b) 3 (c) 4 (d) 5

34. Let  $PQ$  and  $QR$  be diagonals of adjacent faces of a rectangular box, with its centre at  $O$ . If  $\angle QOR, \angle ROP$  and  $\angle POQ$  are  $\theta, \phi$  and  $\Psi$  respectively then the value of ' $\cos \theta + \cos \phi + \cos \Psi$ ' is :

- (a)  $-2$  (b)  $-\sqrt{3}$  (c)  $-1$  (d) 0

35. The value of  $\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \\ \vec{a} \cdot \vec{p} & \vec{b} \cdot \vec{p} & \vec{c} \cdot \vec{p} \\ \vec{a} \cdot \vec{q} & \vec{b} \cdot \vec{q} & \vec{c} \cdot \vec{q} \end{vmatrix}$  is equal to :

- (a)  $(\vec{p} \times \vec{q}) [\vec{a} \times \vec{b} \vec{b} \times \vec{c} \vec{c} \times \vec{a}]$  (b)  $2(\vec{p} \times \vec{q}) [\vec{a} \times \vec{b} \vec{b} \times \vec{c} \vec{c} \times \vec{a}]$   
 (c)  $4(\vec{p} \times \vec{q}) [\vec{a} \times \vec{b} \vec{b} \times \vec{c} \vec{c} \times \vec{a}]$  (d)  $(\vec{p} \times \vec{q}) \sqrt{[\vec{a} \times \vec{b} \vec{b} \times \vec{c} \vec{c} \times \vec{a}]}$

36. If  $\vec{r} = a(\vec{m} \times \vec{n}) + b(\vec{n} \times \vec{l}) + c(\vec{l} \times \vec{m})$  and  $[\vec{l} \vec{m} \vec{n}] = 4$ , find  $\frac{a+b+c}{\vec{r} \cdot (\vec{l} + \vec{m} + \vec{n})}$  :

- (a)  $\frac{1}{4}$  (b)  $\frac{1}{2}$  (c) 1 (d) 2

37. The volume of tetrahedron, for which three co-terminus edges are  $\vec{a}, \vec{b}$  and  $\vec{c}$ , is  $k$  units. Then, the volume of a parallelepiped formed by  $\vec{a} - \vec{b}, \vec{b} + 2\vec{c}$  and  $3\vec{a} - \vec{c}$  is :

- (a)  $6k$  (b)  $7k$  (c)  $30k$  (d)  $42k$

38. The equation of a plane passing through the line of intersection of the planes :  $x + 2y + z - 10 = 0$  and  $3x + y - z = 5$  and passing through the origin is :

- (a)  $5x + 3z = 0$  (b)  $5x - 3z = 0$   
 (c)  $5x + 4y + 3z = 0$  (d)  $5x - 4y + 3z = 0$

39. Find the locus of a point whose distance from  $x$ -axis is twice the distance from the point  $(1, -1, 2)$ :

(a)  $y^2 + 2x - 2y - 4z + 6 = 0$

(b)  $x^2 + 2x - 2y - 4z + 6 = 0$

(c)  $x^2 - 2x + 2y - 4z + 6 = 0$

(d)  $z^2 - 2x + 2y - 4z + 6 = 0$

### Answers

1.	(b)	2.	(b)	3.	(c)	4.	(b)	5.	(a)	6.	(c)	7.	(d)	8.	(c)	9.	(c)	10.	(d)
11.	(d)	12.	(d)	13.	(c)	14.	(c)	15.	(d)	16.	(b)	17.	(b)	18.	(b)	19.	(b)	20.	(b)
21.	(b)	22.	(a)	23.	(b)	24.	(a)	25.	(d)	26.	(d)	27.	(b)	28.	(a)	29.	(b)	30.	(d)
31.	(d)	32.	(d)	33.	(b)	34.	(c)	35.	(d)	36.	(a)	37.	(d)	38.	(b)	39.	(c)		

**Exercise-2 : One or More than One Answer is/are Correct**

1. If equation of three lines are :

$$\frac{x}{1} = \frac{y}{2} = \frac{z}{3}; \frac{x}{2} = \frac{y}{1} = \frac{z}{3} \text{ and } \frac{x-1}{1} = \frac{2-y}{1} = \frac{z-3}{0}, \text{ then}$$

which of the following statement(s) is/are correct ?

- (a) Triangle formed by the line is equilateral
- (b) Triangle formed by the lines is isosceles
- (c) Equation of the plane containing the lines is  $x + y = z$
- (d) Area of the triangle formed by the lines is  $\frac{3\sqrt{3}}{2}$

2. If  $\vec{a} = \hat{i} + 6\hat{j} + 3\hat{k}$ ;  $\vec{b} = 3\hat{i} + 2\hat{j} + \hat{k}$  and  $\vec{c} = (\alpha + 1)\hat{i} + (\beta - 1)\hat{j} + \hat{k}$  are linearly dependent vectors and  $|\vec{c}| = \sqrt{6}$ ; then the possible value(s) of  $(\alpha + \beta)$  can be :

- (a) 1
- (b) 2
- (c) 3
- (d) 4

3. Consider the lines :

$$L_1 : \frac{x-2}{1} = \frac{y-1}{7} = \frac{z+2}{-5}$$

$$L_2 : x - 4 = y + 3 = -z$$

Then which of the following is/are correct ?

- (a) Point of intersection of  $L_1$  and  $L_2$  is  $(1, -6, 3)$
- (b) Equation of plane containing  $L_1$  and  $L_2$  is  $x + 2y + 3z + 2 = 0$
- (c) Acute angle between  $L_1$  and  $L_2$  is  $\cot^{-1}\left(\frac{13}{15}\right)$
- (d) Equation of plane containing  $L_1$  and  $L_2$  is  $x + 2y + 2z + 3 = 0$

4. Let  $\hat{a}, \hat{b}$  and  $\hat{c}$  be three unit vectors such that  $\hat{a} = \hat{b} + (\hat{b} \times \hat{c})$ , then the possible value(s) of  $|\hat{a} + \hat{b} + \hat{c}|^2$  can be :

- (a) 1
- (b) 4
- (c) 16
- (d) 9

5. The value(s) of  $\mu$  for which the straight lines  $\vec{r} = 3\hat{i} - 2\hat{j} - 4\hat{k} + \lambda_1(\hat{i} - \hat{j} + \mu\hat{k})$  and  $\vec{r} = 5\hat{i} - 2\hat{j} + \hat{k} + \lambda_2(\hat{i} + \mu\hat{j} + 2\hat{k})$  are coplanar is/are :

- (a)  $\frac{5 + \sqrt{33}}{4}$
- (b)  $\frac{-5 + \sqrt{33}}{4}$
- (c)  $\frac{5 - \sqrt{33}}{4}$
- (d)  $\frac{-5 - \sqrt{33}}{4}$

6. If  $\hat{i} \times [(\vec{a} - \hat{j}) \times \hat{i}] + \hat{j} \times [(\vec{a} - \hat{k}) \times \hat{j}] + \hat{k} \times [(\vec{a} - \hat{i}) \times \hat{k}] = 0$  and  $\vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$ , then :

- (a)  $x + y = 1$
- (b)  $y + z = \frac{1}{2}$
- (c)  $x + z = 1$
- (d) None of these

7. The value of expression  $[\vec{a} \times \vec{b} \ \vec{c} \times \vec{d} \ \vec{e} \times \vec{f}]$  is equal to :

- (a)  $[\vec{a} \vec{b} \vec{d}][\vec{c} \vec{e} \vec{f}] - [\vec{a} \vec{b} \vec{c}][\vec{d} \vec{e} \vec{f}]$       (b)  $[\vec{a} \vec{b} \vec{e}][\vec{f} \vec{c} \vec{d}] - [\vec{a} \vec{b} \vec{f}][\vec{e} \vec{c} \vec{d}]$   
 (c)  $[\vec{c} \vec{d} \vec{a}][\vec{b} \vec{e} \vec{f}] - [\vec{c} \vec{d} \vec{b}][\vec{a} \vec{e} \vec{f}]$       (d)  $[\vec{b} \vec{c} \vec{d}][\vec{a} \vec{e} \vec{f}] - [\vec{b} \vec{c} \vec{f}][\vec{a} \vec{e} \vec{d}]$

8. If  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$  are the position vectors of the points  $A, B, C$  and  $D$  respectively in three dimensional space and satisfy the relation  $3\vec{a} - 2\vec{b} + \vec{c} - 2\vec{d} = 0$ , then :

- (a)  $A, B, C$  and  $D$  are coplanar  
 (b) The line joining the points  $B$  and  $D$  divides the line joining the point  $A$  and  $C$  in the ratio of 2:1  
 (c) The line joining the points  $A$  and  $C$  divides the line joining the points  $B$  and  $D$  in the ratio of 1:1  
 (d) The four vectors  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$  are linearly dependent.

9. If  $OABC$  is a tetrahedron with equal edges and  $\hat{p}, \hat{q}, \hat{r}$  are unit vectors along bisectors of

$\vec{OA}, \vec{OB} : \vec{OB}, \vec{OC} : \vec{OC}, \vec{OA}$  respectively and  $\hat{a} = \frac{\vec{OA}}{|\vec{OA}|}, \hat{b} = \frac{\vec{OB}}{|\vec{OB}|}, \hat{c} = \frac{\vec{OC}}{|\vec{OC}|}$ , then :

- (a)  $\frac{[\hat{a} \hat{b} \hat{c}]}{[\hat{p} \hat{q} \hat{r}]} = \frac{3\sqrt{3}}{2}$       (b)  $\frac{[\hat{a} + \hat{b} \ \hat{b} + \hat{c} \ \hat{c} + \hat{a}]}{[\hat{p} + \hat{q} \ \hat{q} + \hat{r} \ \hat{r} + \hat{p}]} = \frac{3\sqrt{3}}{4}$   
 (c)  $\frac{[\hat{a} + \hat{b} \ \hat{b} + \hat{c} \ \hat{c} + \hat{a}]}{[\hat{p} \hat{q} \hat{r}]} = \frac{3\sqrt{3}}{2}$       (d)  $\frac{[\hat{a} \hat{b} \hat{c}]}{[\hat{p} + \hat{q} \ \hat{q} + \hat{r} \ \hat{r} + \hat{p}]} = \frac{3\sqrt{3}}{4}$

10. Let  $\hat{a}$  and  $\hat{c}$  are unit vectors and  $|\vec{b}| = 4$ . If the angle between  $\hat{a}$  and  $\hat{c}$  is  $\cos^{-1}\left(\frac{1}{4}\right)$ ; and

$\vec{b} - 2\hat{c} = \lambda\hat{a}$ , then the value of  $\lambda$  can be :

- (a) 2      (b) -3  
 (c) 3      (d) -4

11. Consider the line  $L_1: x = y = z$  and the line  $L_2: 2x + y + z - 1 = 0 = 3x + y + 2z - 2$ , then :

- (a) The shortest distance between the two lines is  $\frac{1}{\sqrt{2}}$   
 (b) The shortest distance between the two lines is  $\sqrt{2}$   
 (c) Plane containing the line  $L_2$  and parallel to line  $L_1$  is  $z - x + 1 = 0$   
 (d) Perpendicular distance of origin from plane containing line  $L_2$  and parallel to line  $L_1$  is  $\frac{1}{\sqrt{2}}$

12. Let  $\vec{r} = \sin x (\vec{a} \times \vec{b}) + \cos y (\vec{b} \times \vec{c}) + 2(\vec{c} \times \vec{a})$ , where  $\vec{a}, \vec{b}$  and  $\vec{c}$  are three non-coplanar vectors. It is given that  $\vec{r}$  is perpendicular to  $\vec{a} + \vec{b} + \vec{c}$ . The possible value(s) of  $x^2 + y^2$  is/are :
- (a)  $\pi^2$  (b)  $\frac{5\pi^2}{4}$   
 (c)  $\frac{35\pi^2}{4}$  (d)  $\frac{37\pi^2}{4}$
13. If  $(\vec{a} \times \vec{b}) \times (\vec{c} \times \vec{d}) = h \vec{a} + k \vec{b} = r \vec{c} + s \vec{d}$ , where  $\vec{a}, \vec{b}$  are non-collinear and  $\vec{c}, \vec{d}$  are also non-collinear then :
- (a)  $h = [\vec{b} \vec{c} \vec{d}]$  (b)  $k = [\vec{a} \vec{c} \vec{d}]$   
 (c)  $r = [\vec{a} \vec{b} \vec{d}]$  (d)  $s = -[\vec{a} \vec{b} \vec{c}]$
14. Let  $a$  be a real number and  $\vec{\alpha} = \hat{i} + 2\hat{j}$ ,  $\vec{\beta} = 2\hat{i} + a\hat{j} + 10\hat{k}$ ,  $\vec{\gamma} = 12\hat{i} + 20\hat{j} + a\hat{k}$  be three vectors, then  $\vec{\alpha}, \vec{\beta}$  and  $\vec{\gamma}$  are linearly independent for :
- (a)  $a > 0$  (b)  $a < 0$   
 (c)  $a = 0$  (d) No value of  $a$
15. The volume of a right triangular prism  $ABCA_1B_1C_1$  is equal to 3. If the position vectors of the vertices of the base  $ABC$  are  $A(1, 0, 1); B(2, 0, 0)$  and  $C(0, 1, 0)$ , then the position vectors of the vertex  $A_1$  can be :
- (a)  $(2, 2, 2)$  (b)  $(0, 2, 0)$   
 (c)  $(0, -2, 2)$  (d)  $(0, -2, 0)$
16. If  $\vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$ ,  $\vec{b} = y\hat{i} + z\hat{j} + x\hat{k}$ , and  $\vec{c} = z\hat{i} + x\hat{j} + y\hat{k}$ , then  $\vec{a} \times (\vec{b} \times \vec{c})$  is :
- (a) Parallel to  $(y - z)\hat{i} + (z - x)\hat{j} + (x - y)\hat{k}$   
 (b) Orthogonal to  $\hat{i} + \hat{j} + \hat{k}$   
 (c) Orthogonal to  $(y + z)\hat{i} + (z + x)\hat{j} + (x + y)\hat{k}$ ,  
 (d) Orthogonal to  $x\hat{i} + y\hat{j} + z\hat{k}$
17. If a line has a vector equation,  $\vec{r} = 2\hat{i} + 6\hat{j} + \lambda(\hat{i} - 3\hat{j})$  then which of the following statements holds good ?
- (a) the line is parallel to  $2\hat{i} + 6\hat{j}$   
 (b) the line passes through the point  $3\hat{i} + 3\hat{j}$   
 (c) the line passes through the point  $\hat{i} + 9\hat{j}$   
 (d) the line is parallel to  $xy$  plane

18. Let  $M, N, P$  and  $Q$  be the mid points of the edges  $AB, CD, AC$  and  $BD$  respectively of the tetrahedron  $ABCD$ . Further,  $MN$  is perpendicular to both  $AB$  and  $CD$  and  $PQ$  is perpendicular to both  $AC$  and  $BD$ . Then which of the following is/are correct :
- (a)  $AB = CD$  (b)  $BC = DA$   
 (c)  $AC = BD$  (d)  $AN = BN$
19. The solution vectors  $\vec{r}$  of the equation  $\vec{r} \times \hat{i} = \hat{j} + \hat{k}$  and  $\vec{r} \times \hat{j} = \hat{k} + \hat{i}$  represent two straight lines which are :
- (a) Intersecting (b) Non coplanar (c) Coplanar (d) Non intersecting
20. Which of the following statement(s) is/are incorrect ?
- (a) The lines  $\frac{x-4}{-3} = \frac{y+6}{-1} = \frac{z+6}{-1}$  and  $\frac{x-1}{-1} = \frac{y-2}{-2} = \frac{z-3}{2}$  are orthogonal  
 (b) The planes  $3x - 2y - 4z = 3$  and the plane  $x - y - z = 3$  are orthogonal  
 (c) The function  $f(x) = \ln(e^{-2} + e^x)$  is monotonic increasing  $\forall x \in \mathbb{R}$   
 (d) If  $g$  is the inverse of the function,  $f(x) = \ln(e^{-2} + e^x)$  then  $g(x) = \ln(e^x - e^{-2})$
21. The lines with vector equations are;  $\vec{r}_1 = -3\hat{i} + 6\hat{j} + \lambda(-4\hat{i} + 3\hat{j} + 2\hat{k})$  and  $\vec{r}_2 = -2\hat{i} + 7\hat{j} + \mu(-4\hat{i} + \hat{j} + \hat{k})$  are such that :
- (a) they are coplanar  
 (b) they do not intersect  
 (c) they are skew  
 (d) the angle between them is  $\tan^{-1}(3/7)$

### Answers

1.	(b, c, d)	2.	(a, c)	3.	(a, b, c)	4.	(a, d)	5.	(a, c)	6.	(a, c)
7.	(a, b, c)	8.	(a, c, d)	9.	(a, d)	10.	(c, d)	11.	(a, d)	12.	(b, d)
13.	(b, c, d)	14.	(a, b, c)	15.	(a, d)	16.	(a, b, c, d)	17.	(b, c, d)	18.	(a, b, c, d)
19.	(b, d)	20.	(a, b)	21.	(b, c, d)						




**Exercise-3 : Comprehension Type Problems**
**Paragraph for Question Nos. 1 to 3**

The vertices of  $\Delta ABC$  are  $A(2, 0, 0)$ ,  $B(0, 1, 0)$ ,  $C(0, 0, 2)$ . Its orthocentre is  $H$  and circumcentre is  $S$ .  $P$  is a point equidistant from  $A, B, C$  and the origin  $O$ .

- The  $z$ -coordinate of  $H$  is :  
 (a) 1                      (b)  $1/2$                       (c)  $1/6$                       (d)  $1/3$
- The  $y$ -coordinate of  $S$  is :  
 (a)  $5/6$                       (b)  $1/3$                       (c)  $1/6$                       (d)  $1/2$
- $PA$  is equal to :  
 (a) 1                      (b)  $\sqrt{2}$                       (c)  $\sqrt{\frac{3}{2}}$                       (d)  $\frac{3}{2}$

**Paragraph for Question Nos. 4 to 6**

Consider a plane  $\pi: \vec{r} \cdot \vec{n} = d$  (where  $\vec{n}$  is not a unit vector). There are two points  $A(\vec{a})$  and  $B(\vec{b})$  lying on the same side of the plane.

- If foot of perpendicular from  $A$  and  $B$  to the plane  $\pi$  are  $P$  and  $Q$  respectively, then length of  $PQ$  be:  
 (a)  $\frac{|(\vec{b} - \vec{a}) \cdot \vec{n}|}{|\vec{n}|}$                       (b)  $|(\vec{b} - \vec{a}) \cdot \vec{n}|$                       (c)  $\frac{|(\vec{b} - \vec{a}) \times \vec{n}|}{|\vec{n}|}$                       (d)  $|(\vec{b} - \vec{a}) \times \vec{n}|$
- Reflection of  $A(\vec{a})$  in the plane  $\pi$  has the position vector :  
 (a)  $\vec{a} + \frac{2}{(\vec{n})^2} (d - \vec{a} \cdot \vec{n}) \vec{n}$                       (b)  $\vec{a} - \frac{1}{(\vec{n})^2} (d - \vec{a} \cdot \vec{n}) \vec{n}$   
 (c)  $\vec{a} + \frac{2}{(\vec{n})^2} (d + \vec{a} \cdot \vec{n}) \vec{n}$                       (d)  $\vec{a} + \frac{2}{(\vec{n})^2} \vec{n}$
- If a plane  $\pi_1$  is drawn from the point  $A(\vec{a})$  and another plane  $\pi_2$  is drawn from point  $B(\vec{b})$  parallel to  $\pi$ , then the distance between the planes  $\pi_1$  and  $\pi_2$  is :  
 (a)  $\frac{|(\vec{a} - \vec{b}) \cdot \vec{n}|}{|\vec{n}|}$                       (b)  $|(\vec{a} - \vec{b}) \cdot \vec{n}|$                       (c)  $|(\vec{a} - \vec{b}) \times \vec{n}|$                       (d)  $\frac{|(\vec{a} - \vec{b}) \times \vec{n}|}{|\vec{n}|}$

### Paragraph for Question Nos. 7 to 9

Consider a plane  $\Pi: \vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) = 5$ , a line  $L_1: \vec{r} = (3\hat{i} - \hat{j} + 2\hat{k}) + \lambda(2\hat{i} - 3\hat{j} - \hat{k})$  and a point  $A(3, -4, 1)$ .  $L_2$  is a line passing through  $A$  intersecting  $L_1$  and parallel to plane  $\Pi$ .

7. Equation of  $L_2$  is :

- (a)  $\vec{r} = (1 + \lambda)\hat{i} + (2 - 3\lambda)\hat{j} + (1 - \lambda)\hat{k}; \lambda \in R$   
 (b)  $\vec{r} = (3 + \lambda)\hat{i} - (4 - 2\lambda)\hat{j} + (1 + 3\lambda)\hat{k}; \lambda \in R$   
 (c)  $\vec{r} = (3 + \lambda)\hat{i} - (4 + 3\lambda)\hat{j} + (1 - \lambda)\hat{k}; \lambda \in R$   
 (d) None of the above

8. Plane containing  $L_1$  and  $L_2$  is :

- (a) parallel to  $yz$ -plane  
 (b) parallel to  $x$ -axis  
 (c) parallel to  $y$ -axis  
 (d) passing through origin

9. Line  $L_1$  intersects plane  $\Pi$  at  $Q$  and  $xy$ -plane at  $R$  the volume of tetrahedron  $OAQR$  is :  
 (where 'O' is origin)

- (a) 0  
 (b)  $\frac{14}{3}$   
 (c)  $\frac{3}{7}$   
 (d)  $\frac{7}{3}$

### Paragraph for Question Nos. 10 to 11

Consider three planes :

$$2x + py + 6z = 8; x + 2y + qz = 5 \text{ and } x + y + 3z = 4$$

10. Three planes intersect at a point if :

- (a)  $p = 2, q \neq 3$       (b)  $p \neq 2, q \neq 3$       (c)  $p \neq 2, q = 3$       (d)  $p = 2, q = 3$

11. Three planes do not have any common point of intersection if :

- (a)  $p = 2, q \neq 3$       (b)  $p \neq 2, q \neq 3$       (c)  $p \neq 2, q = 3$       (d)  $p = 2, q = 3$

### Paragraph for Question Nos. 12 to 14

The points  $A, B$  and  $C$  with position vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$  respectively lie on a circle centered at origin  $O$ . Let  $G$  and  $E$  be the centroid of  $\triangle ABC$  and  $\triangle ACD$  respectively where  $D$  is mid point of  $AB$ .

12. If  $OE$  and  $CD$  are mutually perpendicular, then which of the following will be necessarily true ?

- (a)  $|\vec{b} - \vec{a}| = |\vec{c} - \vec{a}|$   
 (b)  $|\vec{b} - \vec{a}| = |\vec{b} - \vec{c}|$   
 (c)  $|\vec{c} - \vec{a}| = |\vec{c} - \vec{b}|$   
 (d)  $|\vec{b} - \vec{a}| = |\vec{c} - \vec{a}| = |\vec{b} - \vec{c}|$

13. If  $GE$  and  $CD$  are mutually perpendicular, then orthocenter of  $\Delta ABC$  must lie on :  
 (a) median through  $A$  (b) median through  $C$   
 (c) angle bisector through  $A$  (d) angle bisector through  $B$
14. If  $[\vec{AB} \ \vec{AC} \ \vec{AB} \times \vec{AC}] = \lambda [\vec{AE} \ \vec{AG} \ \vec{AE} \times \vec{AG}]$ , then the value of  $\lambda$  is :  
 (a)  $-18$  (b)  $18$  (c)  $-324$  (d)  $324$

**Paragraph for Question Nos. 15 to 16**

Consider a tetrahedron  $D-ABC$  with position vectors if its angular points as  
 $A(1, 1, 1); B(1, 2, 3); C(1, 1, 2)$   
 and centre of tetrahedron  $\left(\frac{3}{2}, \frac{3}{4}, 2\right)$ .

15. Shortest distance between the skew lines  $AB$  and  $CD$  :  
 (a)  $\frac{1}{2}$  (b)  $\frac{1}{3}$  (c)  $\frac{1}{4}$  (d)  $\frac{1}{5}$
16. If  $N$  be the foot of the perpendicular from point  $D$  on the plane face  $ABC$  then the position vector of  $N$  are :  
 (a)  $(-1, 1, 2)$  (b)  $(1, -1, 2)$  (c)  $(1, 1, -2)$  (d)  $(-1, -1, 2)$

**Paragraph for Question Nos. 17 to 18**

In a triangle  $AOB$ ,  $R$  and  $Q$  are the points on the side  $OB$  and  $AB$  respectively such that  $3OR = 2RB$  and  $2AQ = 3QB$ . Let  $OQ$  and  $AR$  intersect at the point  $P$  (where  $O$  is origin).

17. If the point  $P$  divides  $OQ$  in the ratio of  $\mu : 1$ , then  $\mu$  is :  
 (a)  $\frac{2}{19}$  (b)  $\frac{2}{17}$  (c)  $\frac{2}{15}$  (d)  $\frac{10}{9}$
18. If the ratio of area of quadrilateral  $PQBR$  and area of  $\Delta OPA$  is  $\frac{\alpha}{\beta}$  then  $(\beta - \alpha)$  is (where  $\alpha$  and  $\beta$  are coprime numbers) :  
 (a)  $1$  (b)  $9$  (c)  $7$  (d)  $0$

**Answers**

1.	(d)	2.	(c)	3.	(d)	4.	(c)	5.	(a)	6.	(a)	7.	(c)	8.	(b)	9.	(d)	10.	(b)
11.	(c)	12.	(a)	13.	(b)	14.	(d)	15.	(b)	16.	(b)	17.	(d)	18.	(d)				

### Exercise-4 : Matching Type Problems

1.

	Column-I		Column-II
(A)	Lines $\frac{x-1}{-2} = \frac{y+2}{3} = \frac{z}{-1}$ and $\vec{r} = (3\hat{i} - \hat{j} + \hat{k}) + t(\hat{i} + \hat{j} + \hat{k})$ are	(P)	Intersecting
(B)	Lines $\frac{x+5}{1} = \frac{y-3}{7} = \frac{z+3}{3}$ and $x - y + 2z - 4 = 0 = 2x + y - 3z + 5$ are	(Q)	Perpendicular
(C)	Lines $(x = t - 3, y = -2t + 1, z = -3t - 2)$ and $\vec{r} = (t + 1)\hat{i} + (2t + 3)\hat{j} + (-t - 9)\hat{k}$ are	(R)	Parallel
(D)	Lines $\vec{r} = (\hat{i} + 3\hat{j} - \hat{k}) + t(2\hat{i} - \hat{j} - \hat{k})$ and $\vec{r} = (-\hat{i} - 2\hat{j} + 5\hat{k}) + s\left(\hat{i} - 2\hat{j} + \frac{3}{4}\hat{k}\right)$ are	(S)	Skew
		(T)	Coincident

2.

	Column-I		Column-II
(A)	If $\vec{a}, \vec{b}$ and $\vec{c}$ are three mutually perpendicular vectors where $ \vec{a}  =  \vec{b}  = 2,  \vec{c}  = 1$ , then $[\vec{a} \times \vec{b}, \vec{b} \times \vec{c}, \vec{c} \times \vec{a}]$ is	(P)	-12
(B)	If $\vec{a}$ and $\vec{b}$ are two unit vectors inclined at $\frac{\pi}{3}$ , then $16[\vec{a}, \vec{b} + (\vec{a} \times \vec{b}), \vec{b}]$ is	(Q)	0
(C)	If $\vec{b}$ and $\vec{c}$ are orthogonal unit vectors and $\vec{b} \times \vec{c} = \vec{a}$ then $[\vec{a} + \vec{b} + \vec{c}, \vec{a} + \vec{b}, \vec{b} + \vec{c}]$ is	(R)	16
(D)	If $[\vec{x}, \vec{y}, \vec{a}] = [\vec{x}, \vec{y}, \vec{b}] = [\vec{a}, \vec{b}, \vec{c}] = 0$ , each vector being a non-zero vector, then $[\vec{x}, \vec{y}, \vec{c}]$ is	(S)	1
		(T)	4

3.

Column-I		Column-II	
(A)	The number of real roots of equation $2^x + 3^x + 4^x - 9^x = 0$ is $\lambda$ , then $\lambda^2 + 7$ is divisible by	(P)	2
(B)	Let $ABC$ be a triangle whose centroid is $G$ , orthocenter is $H$ and circumcentre is the origin ' $O$ '. If $D$ is any point in the plane of the triangle such that not three of $O, A, B, C$ and $D$ are collinear satisfying the relation $\vec{AD} + \vec{BD} + \vec{CH} + 3\vec{HG} = \lambda\vec{HD}$ , then $\lambda + 4$ is divisible by	(Q)	3
(C)	If $A$ $(\text{adj } A) = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$ , then $5 A  - 2$ is divisible by	(R)	4
(D)	$\vec{a}, \vec{b}, \vec{c}$ are three unit vector such that $\vec{a} + \vec{b} = \sqrt{2}\vec{c}$ , then $ 6\vec{a} - 8\vec{b} $ is divisible by	(S)	6
		(T)	10

**Answers**

1.	$A \rightarrow Q, S; B \rightarrow R; C \rightarrow P, Q; D \rightarrow P$
2.	$A \rightarrow R; B \rightarrow P; C \rightarrow S; D \rightarrow Q$
3.	$A \rightarrow P, R; B \rightarrow P, Q, S; C \rightarrow P, Q, R, S; D \rightarrow P, T$

### Exercise-5 : Subjective Type Problems

1. A straight line  $L$  intersects perpendicularly both the lines :

$$\frac{x+2}{2} = \frac{y+6}{3} = \frac{z-34}{-10} \text{ and } \frac{x+6}{4} = \frac{y-7}{-3} = \frac{z-7}{-2},$$

then the square of perpendicular distance of origin from  $L$  is

2. If  $\hat{\mathbf{a}}, \hat{\mathbf{b}}$  and  $\hat{\mathbf{c}}$  are non-coplanar unit vectors such that  $[\hat{\mathbf{a}} \hat{\mathbf{b}} \hat{\mathbf{c}}] = [\hat{\mathbf{b}} \times \hat{\mathbf{c}} \hat{\mathbf{c}} \times \hat{\mathbf{a}} \hat{\mathbf{a}} \times \hat{\mathbf{b}}]$ , then find the projection of  $\hat{\mathbf{b}} + \hat{\mathbf{c}}$  on  $\hat{\mathbf{a}} \times \hat{\mathbf{b}}$ .

3. Let  $OA, OB, OC$  be coterminal edges of a cuboid. If  $l, m, n$  be the shortest distances between the sides  $OA, OB, OC$  and their respective skew body diagonals to them, respectively, then find

$$\frac{\left(\frac{1}{l^2} + \frac{1}{m^2} + \frac{1}{n^2}\right)}{\left(\frac{1}{OA^2} + \frac{1}{OB^2} + \frac{1}{OC^2}\right)}.$$

4. Let  $OABC$  be a tetrahedron whose edges are of unit length. If  $\vec{OA} = \vec{\mathbf{a}}, \vec{OB} = \vec{\mathbf{b}}$  and  $\vec{OC} = \alpha(\vec{\mathbf{a}} + \vec{\mathbf{b}}) + \beta(\vec{\mathbf{a}} \times \vec{\mathbf{b}})$ , then  $(\alpha\beta)^2 = \frac{p}{q}$  where  $p$  and  $q$  are relatively prime to each other.

Find the value of  $\left[\frac{q}{2p}\right]$  where  $[\cdot]$  denotes greatest integer function.

5. Let  $\vec{\mathbf{v}}_0$  be a fixed vector and  $\vec{\mathbf{v}}_0 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ . Then for  $n \geq 0$  a sequence is defined

$$\vec{\mathbf{v}}_{n+1} = \vec{\mathbf{v}}_n + \left(\frac{1}{2}\right)^{n+1} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^{n+1} \vec{\mathbf{v}}_0 \text{ then } \lim_{n \rightarrow \infty} \vec{\mathbf{v}}_n = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}. \text{ Find } \frac{\alpha}{\beta}.$$

6. If  $A$  is the matrix  $\begin{bmatrix} 1 & -3 \\ -1 & 1 \end{bmatrix}$ , then  $A - \frac{1}{3}A^2 + \frac{1}{9}A^3 - \dots + \left(-\frac{1}{3}\right)^n A^{n+1} + \dots = \frac{3}{13} \begin{bmatrix} 1 & a \\ b & 1 \end{bmatrix}$ .

Find  $\left|\frac{a}{b}\right|$ .

7. A sequence of  $2 \times 2$  matrices  $\{M_n\}$  is defined as follows  $M_n = \begin{bmatrix} \frac{1}{(2n+1)!} & \frac{1}{(2n+2)!} \\ \sum_{k=0}^n \frac{(2n+2)!}{(2k+2)!} & \sum_{k=0}^n \frac{(2n+1)!}{(2k+1)!} \end{bmatrix}$

then  $\lim_{n \rightarrow \infty} \det. (M_n) = \lambda - e^{-1}$ . Find  $\lambda$ .

8. Let  $|\vec{\mathbf{a}}| = 1, |\vec{\mathbf{b}}| = 1$  and  $|\vec{\mathbf{a}} + \vec{\mathbf{b}}| = \sqrt{3}$ . If  $\vec{\mathbf{c}}$  be a vector such that  $\vec{\mathbf{c}} = \vec{\mathbf{a}} + 2\vec{\mathbf{b}} - 3(\vec{\mathbf{a}} \times \vec{\mathbf{b}})$  and  $p = |(\vec{\mathbf{a}} \times \vec{\mathbf{b}}) \times \vec{\mathbf{c}}|$ , then find  $[p^2]$ . (where  $[\cdot]$  represents greatest integer function).

9. Let  $\vec{r} = (\vec{a} \times \vec{b}) \sin x + (\vec{b} \times \vec{c}) \cos y + 2(\vec{c} \times \vec{a})$ , where  $\vec{a}, \vec{b}, \vec{c}$  are non-zero and non-coplanar vectors. If  $\vec{r}$  is orthogonal to  $\vec{a} + \vec{b} + \vec{c}$ , then find the minimum value of  $\frac{4}{\pi^2}(x^2 + y^2)$ .
10. The plane denoted by  $\Pi_1 : 4x + 7y + 4z + 81 = 0$  is rotated through a right angle about its line of intersection with the plane  $\Pi_2 : 5x + 3y + 10z = 25$ . If the plane in its new position be denoted by  $\Pi$ , and the distance of this plane from the origin is  $\sqrt{53} k$  where  $k \in N$ , then find  $k$ .
11.  $ABCD$  is a regular tetrahedron,  $A$  is the origin and  $B$  lies on  $x$ -axis.  $ABC$  lies in the  $xy$ -plane  $|\vec{AB}| = 2$ . Under these conditions, the number of possible tetrahedrons is :
12.  $A, B, C, D$  are four points in the space and satisfy  $|\vec{AB}| = 3, |\vec{BC}| = 7, |\vec{CD}| = 11$  and  $|\vec{DA}| = 9$ . Then find the value of  $\vec{AC} \cdot \vec{BD}$ .
13. Let  $OABC$  be a regular tetrahedron of edge length unity. Its volume be  $V$  and  $6V = \sqrt{p/q}$  where  $p$  and  $q$  are relatively prime. The find the value of  $(p + q)$  :
14. If  $\vec{a}$  and  $\vec{b}$  are non zero, non collinear vectors and  $\vec{a}_1 = \lambda \vec{a} + 3\vec{b}; \vec{b}_1 = 2\vec{a} + \lambda \vec{b}; \vec{c}_1 = \vec{a} + \vec{b}$ . Find the sum of all possible real values of  $\lambda$  so that points  $A_1, B_1, C_1$  whose position vectors are  $\vec{a}_1, \vec{b}_1, \vec{c}_1$  respectively are collinear is equal to .
15. Let  $P$  and  $Q$  are two points on curve  $y = \log_{\frac{1}{2}}\left(x - \frac{1}{2}\right) + \log_2 \sqrt{4x^2 - 4x + 1}$  and  $P$  is also on  $x^2 + y^2 = 10$ .  $Q$  lies inside the given circle such that its abscissa is integer. Find the smallest possible value of  $\vec{OP} \cdot \vec{OQ}$  where 'O' being origin.
16. In above problem find the largest possible value of  $|\vec{PQ}|$ .
17. If  $a, b, c, l, m, n \in R - \{0\}$  such that  $al + bm + cn = 0, bl + cm + an = 0, cl + am + bn = 0$ . If  $a, b, c$  are distinct and  $f(x) = ax^3 + bx^2 + cx + 2$ . Find  $f(1)$ :
18. Let  $\vec{\mu}$  and  $\vec{v}$  are unit vectors and  $\vec{\omega}$  is vector such that  $\vec{\mu} \times \vec{v} + \vec{\mu} = \vec{\omega}$  and  $\vec{\omega} \times \vec{\mu} = \vec{v}$ . The find the value of  $[\vec{\mu} \ \vec{v} \ \vec{\omega}]$ .

Answers

1.	5	2.	1	3.	2	4.	5	5.	2	6.	3	7.	1
8.	5	9.	5	10.	4	11.	8	12.	0	13.	0	14.	2
15.	4	16.	2	17.	2	18.	1						